OTIC FILE CURY



AD _____

Single-Dose Absorption and Pharmacokinetics of WR 6026

TASK ORDER #3 FINAL REPORT

Brent G. Petty, M.D.
David M. Kornhauser, M.D.
Theresa B. Shapiro, M.D., Ph.D.
Paul S. Lietman, M.D., Ph.D.

01 August 1988

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND Fort Detrick, Maryland 21701-5012

Contract No. DAMD17-85-C-5133

S AUG 2 9 1988

Division of Clinical Pharmacology
The Johns Hopkins University School of Medicine
600 N. Wolfe Street
Baltimore, Maryland 21205

Approved for public release; distribution is unlimited.

The views, opinions, and/or findings contained in this report are those of the authors and should not be construed as an official Department of Army position, policy, or decision, unless so designated by other documents.

88 8 28 04 9

| | SECURITY CLASSIFICATION OF THIS PAGE | | | | | | | |
|--|---|--------------------------------------|---|-------------------------------|------------------------------------|----------------|--|--|
| | REPORT | N PAGE | | | Form Approved OMB No. 0704-0188 | | | |
| | 1a. REPORT SECURITY CLASSIFICATION Unclassified | | 1b. RESTRICTIVE | MARKINGS | | | | |
| 3 2 | 2a. SECURITY CLASSIFICATION AUTHORITY | | /AVAILABILITY OF | | | | | |
| | 2b. DECLASSIFICATION / DOWNGRADING SCHEDU | ILE | | d for Public ution Unlimi | | | | |
| | 4. PERFORMING ORGANIZATION REPORT NUMBER | ER(S) | 5. MONITORING | ORGANIZĂTION REF | ORT NUMBER | R(S) | | |
| X | 6. NAME OF PERFORMING ORGANIZATION Div. of Clinical Pharmacology The Johns Hopkins II. Sch. of Med | 6b. OFFICE SYMBOL (If applicable) | 7a. NAME OF MONITORING ORGANIZATION U.S. Army Medical National Development Center | | | | | |
| | 6c. ADDRESS (City, State, and ZIP Code) 600 N. Wolfe Street - Osler 52 Baltimore, Maryland 21205 | | 7b. ADDRESS (Cit | y, State, and ZIP Co | ide) | | | |
| | 8a. NAME OF FUNDING / SPONSORING ORGANIZATION USAMRDC | 8b. OFFICE SYMBOL (If applicable) | 9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAMD17-85-C-5133 | | | | | |
| Ø. | 8c. ADDRESS (City, State, and ZIP Code) | | 10. SOURCE OF FUNDING NUMBERS PROGRAM PROJECT TASK WORK UN | | | | | |
| - | Fort Detrick Frederick, Maryland 21701-501 | .2 | ELEMENT NO. NO. NO. | | NO. AE | ACCESSION NO. | | |
| 8 | 11. TITLE (Include Security Classification) | | _ 03/31R | 2773 | TLL . | 103 | | |
| 2533 | Phase I Clinical Pharmacology Studies 12. PERSONAL AUTHOR(S) Paul S. Lietman, M.D., Ph.D. Brent G. Petty, M.D.: David M. Kornhauser, M.D.: Theresa B. Shapiro, M.D., Ph.D. 13a. TYPE OF REPORT 13b. TIME COVERED 14. DATE OF REPORT (Year, Month, Day) 15. PAGE COUNT Final - Phase I 1988 August 01 | | | | | | | |
| | Subtitle: Single-Dose Absorp | tion and Pharmac | cokinetics of | WR 6026 | | | | |
| X | 17. COSATI CODES FIELD GROUP SUB-GROUP | 18. SUBJECT TERMS (| Continue on reverse if necessary and identify by block number) | | | | | |
| 78 | 06 01 06 15 | | is, Infection | us Disease, H | luman Stu | dy, Drugs | | |
| 255 | 19. ABSTRACT (Continue on reverse if necessary WR 6026 (8-(6-diethylaminohexy promising agent for the treats | lamino)-6-metho | xy-4-methylqu | inoline dihy is based on e | drochlor experimen | ide) is a | | |
| ************************************** | promising agent for the treatment of visceral leishmaniasis based on experiments in both an animal model and an in vitro test system. This human study was performed in order to increase our understanding of the pharmacokinetics, safety, and tolerance of a single oral 60 mg dose of WR 6026 in healthy male volunteers. | | | | | | | |
| 253 | Hopkins Hospital and which was approved by the Joint Committee on Clinical Investigations | | | | | | | |
| 222 | of The Johns Hopkins Medical Institutions and the Human Use Review Office of the U.S. Army. Following the administration of a single dose of 60 mg of WR 6026, serial blood specimens and urine collections were obtained in order to assess the pharmacokinetics of this compound. The volunteers were monitored for subjective tolerance by daily interview and for | | | | | | | |
| 2 | objective toxicity with clinic | | | | | | | |
| | 20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED SAME AS I | RPT. DTIC USERS | Unclassi | | | -25 en | | |
| | 22a. NAME OF RESPONSIBLE INDIVIDUAL Schuster, Brian, Col., MC | | 22b TELEPHONE ((301) 427 | Include Area Code) -5346 | 22c. OFFICE SCRI | SYMBOL)-UW | | |

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE



19 ABSTRACT (continued)

electrocardiograms, urinalysis, and methemoglobin determinations.

All of the subjects tolerated WR 6026 very well with no adverse symptoms. Two subjects had an increase in the serum aspartate aminotransferase (AST, SGOT) on the fourth day after drug administration and one had a corresponding increase in the serum alanine aminotransferase (ALT, SGPT). Four other subjects had elevations of the serum lactic dehydrogenase (LDH), three on the fourth and one on the second day following drug administration. Whether these elevations were related to laboratory variability or to a delayed effect of the drug is not clear. One subject had an increase in fasting serum triglycerides on the second day following drug administration. No subject had a significant change in hematological parameters, electrocardiograms, methemoglobin, creatine kinase or urinalysis.

The pharmacokinetic results demonstrated that there was approximately a 30-minute lag time between administration and detectable drug absorption, with peak WR 6026 concentrations occurring approximately three hours after drug administration. The areas under the plasma concentration-time curves varied approximately fourfold between the eight subjects. The data were not well described by compartmental analysis. However, using a one-compartment model the mean elimination half-time was about 11 hours, with a relatively wide range between subjects of 5.2 to 17.3 hours.

The urinary excretion of the parent drug and two metabolites over 6 days after dosing with 60 mg WR 6026 was quantified. The average amount of drug recovered in the urine as these three compounds was 14.1% of the dose of WR 6026 administered, with a range of 6.2-30.0%.

In the first two subjects, whole blood as well as plasma concentrations of WR 6026 were measured after dosing. Concentrations in whole blood were lower than those in plasma, indicating that the drug was not concentrated in the cellular components of blood.



| Acces | sion For | |
|-------|------------------------|-------|
| NTIS | GRA&I | 12 |
| DTIC | TAB | |
| Unann | ounced | |
| Justi | fication_ | |
| | ibution/ lability (| Codes |
| A V | Avail and | |
| Dist | Special | • |
| A-1 | | |



| AD | | |
|----|--|--|
| | | |

Single-Dose Absorption and Pharmacokinetics of WR 6026

TASK ORDER #3 FINAL REPORT

Brent G. Petty, M.D.
David M. Kornhauser, M.D.
Theresa B. Shapiro, M.D., Ph.D.
Paul S. Lietman, M.D., Ph.D.

01 August 1988

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND Fort Detrick, Maryland 21701-5012

Contract No. DAMD17-85-C-5133

Division of Clinical Pharmacology
The Johns Hopkins University School of Medicine
600 N. Wolfe Street
Baltimore, Maryland 21205

Approved for public release; distribution is unlimited.

The views, opinions, and/or findings contained in this report are those of the authors and should not be construed as an official Department of Army position, policy, or decision, unless so designated by other documents.

C

SUMMARY

WR 6026 (8-(6-diethylaminohexylamino)-6-methoxy-4-methylquinoline dihydrochloride) is a promising agent for the treatment of visceral leishmaniasis, based on experiments in both an animal model and an <u>in vitro</u> test system. This human study was performed in order to increase our understanding of the pharmacokinetics, safety, and tolerance of a single oral 60 mg dose of WR 6026 in healthy male volunteers.

Eight subjects, who gave written informed consent, participated in this study which was conducted in the Clinical Pharmacology Research Unit, an inpatient service at The Johns Hopkins Hospital, and which was approved by the Joint Committee on Clinical Investigations of The Johns Hopkins Medical Institutions and the Human Use Review Office of the U. S. Army. Following the administration of a single dose of 60 mg of WR 6026, serial blood specimens and urine collections were obtained in order to assess the pharmacokinetics of this compound. The volunteers were monitored for subjective tolerance by daily interview and for objective toxicity with clinical laboratory tests of hematology and chemistry variables, electrocardiograms, urinalysis, and methemoglobin determinations.

All of the subjects tolerated WR 6026 very well with no adverse symptoms. Two subjects had an increase in the serum aspartate aminotransferase (AST, SGOT) on the fourth day after drug administration and one had a corresponding increase in the serum alanine aminotransferase (ALT, SGPT). Four other subjects had elevations of the serum lactic dehydrogenase (LDH), three on the fourth and one on the second day following drug administration. Whether these elevations were related to laboratory variability or to a delayed effect

of the drug is not clear. One subject had an increase in fasting serum triglycerides on the second day following drug administration. No subject had a significant change in hematological parameters, electrocardiograms, methemoglobin, creatine kinase or urinalysis.

The pharmacokinetic results demonstrated that there was approximately a 30-minute lag time between administration and detectable drug absorption, with peak WR 6026 concentrations occurring approximately three hours after drug administration. The areas under the plasma concentration-time curves varied approximately fourfold between the eight subjects. The data were not well described by compartmental analysis. However, using a one-compartment model, the mean elimination half-time was about 11 hours, with a relatively wide range between subjects of 5.2 to 17.3 hours.

The urinary excretion of the parent drug and two metabolites over 6 days after dosing with 60 mg WR 6026 was quantified. The average amount of drug recovered in the urine as these three compounds was 14.1% of the dose of WR 6026 administered, with a range of 6.2-30.0%.

In the first two subjects, whole blood as well as plasma concentrations of WR 6026 were measured after dosing. Concentrations in whole blood were lower than those in plasma, indicating that the drug was not concentrated in the cellular components of blood.

FOREWORD

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

For the protection of human subjects the investigators have adhered to the policies of applicable Federal Law 45 CFR 46.

TABLE OF CONTENTS

| | | | | | Page | | |
|----|-----|------------------------|-------------------------|--|-----------------------|--|--|
| I. | REP | ORT | | | | | |
| | 1. | INTRODUCTION | | | | | |
| | 2. | MATERIALS AND METHODS | | | | | |
| | | 2.1 | WR 6026 | 6 | 4 | | |
| | | 2.2 | Subject | ts | 5 | | |
| | | | 2.2.2 2.2.3 2.2.4 | Inclusion Criteria Exclusion Criteria Recruitment Informed Consent Compensation | 5 5 6 6 7 | | |
| | | 2.3 | Experi | mental Protocol | 7 | | |
| | | | | Objectives Design | 7 7 | | |
| | | 2.4 | Clinica | al Laboratory Evaluations | 8 | | |
| | | | 2.4.2 2.4.3 2.4.4 | Hematology Serum Chemistry Urine Analysis Electrocardiography Chest X-rays | 8 8 9 9 | | |
| | | 2.5 | Specime | en Handling | 10 | | |
| | | | | Blood/Plasma Collection and Storage Urine Collection and Storage Specimen Shipment | 10 10 10 | | |
| | | 2.6 | Assay (| of WR 6026 and Metabolites | 11 | | |
| | | | 2.6.1 2.6.2 | Assay of WR 6026 in Plasma and Blood Assay of WR 6026 and its Metabolites in Urine | 11 12 | | |
| | | 2.7 | Pharma | cokinetic Analysis | 13 | | |
| | 3. | RESULTS AND DISCUSSION | | | | | |
| | | 3.1 | Amendm | ents and Compliance | 15 | | |
| | | | | Amendments | 15 17 | | |

| | 3.2 | Description of Subject Population | 19 | | | |
|-----|-------------|--|----------|--|--|--|
| 3.3 | | Clinical Results | | | | |
| | | 3.3.1 Symptomatic 3.3.2 Laboratory | 19 19 | | | |
| | | 3.3.2.1 Liver Function Tests | 19 | | | |
| | | 3.3.2.2 Creatine Kinase | 20 | | | |
| | | 3.3.2.3 Triglycerides | 21 | | | |
| | | 3.3.2.4 Hematological Tests | 21 | | | |
| | | 3.3.2.5 Electrocardiograms | 21 | | | |
| | | 3.3.2.6 Methemoglobin | 22 | | | |
| | | 3.3.3 Clinical Conclusions | 22 | | | |
| | 3.4 | Pharmacokinetic Results and Discussion | 23 | | | |
| | 4. CONC | LUSIONS | 32 | | | |
| | 5. REFE | RENCES | 34 | | | |
| II. | TABLES AN | D FIGURES | | | | |
| | Table 1 | Individual Subject Characteristics | 36 | | | |
| | Table 2 | Peak Clinical Laboratory Abnormalities | 37 | | | |
| | Table 3 | Plasma Concentrations of WR 6026 | 38 | | | |
| | Table 4 | Pharmacokinetic Parameters for Individual Subjects | 39 | | | |
| | Table 5 | Urinary Excretion of WR 6026 and Metabolites | 40 | | | |
| | Table 6 | Derived Pharmacokinetic Parameters for | | | | |
| | | Individual Subjects | 41 | | | |
| | Table 7 | Comparison of Areas Under the Curve by Trapezoidal | | | | |
| | | Rule and Fitted Equation | 42 | | | |
| | Table 8 | Elimination Rate Constant and Plasma Half-Life of | | | | |
| | | WK 6026 | 43 | | | |
| | Table 9 | Pharmacokinetic Data for Individual Subjects | 44 | | | |
| | Table 10 | Pharmacokinetic Parameters for Individual Subjects | 45 | | | |
| | Figure 1 | Chemical Structure of WR 6026 | 46 | | | |
| | Figure 2 | Plasma WR 6026 Levels for Subjects 1, 2, and 3 | 47 | | | |
| | Figure 3 | Plasma WR 6026 Levels for Subjects 4, 5, and 6 | 48 | | | |
| | Figure 4 | Plasma WR 6026 Levels for Subjects 7 and 8 | 49 | | | |
| | Figure 5 | Average Plasma WR 6026 Concentrations | 50 | | | |
| | Figure 6 | Blood WR 6026 for Subjects 1 and 2 | 51 | | | |
| | Figure 7 | Plotted Plasma WR 6026 Levels for Subject 1 | 52 | | | |
| | Figure 8 | Plotted Plasma WR 6026 Levels for Subject 2 | 53 | | | |
| | Figure 9 | Plotted Plasma WR 6026 Levels for Subject 3 | 54 | | | |
| | Figure 10 | | 55 | | | |
| | Figure 11 | | 56 | | | |
| | Figure 12 | | 57 | | | |
| | Figure 13 | | 58 | | | |
| | Figure 14 | • | 59 60 | | | |
| | לו פתווסוים | MODELING FOR SUDJECT I | h() | | | |

COSC | COSC | COSC |

| | Figure | 16 | Subject | | Calculated | Plasma | Concentrations | for 61 |
|------|--------|-------|-----------------|------|-----------------------------|---------|----------------|--------|
| | Figure | 17 | • | vs. | Calculated | Plasma | Concentrations | |
| | Figure | 18 | • | vs. | Calculated | Plasma | Concentrations | |
| | Figure | 19 | - | vs. | Calculated | Plasma | Concentrations | |
| | Figure | 20 | • | vs. | Calculated | Plasma | Concentrations | |
| | Figure | 21 | _ | vs. | Calculated | Plasma | Concentrations | |
| | Figure | 22 | • | vs. | Calculated | Plasma | Concentrations | |
| | Figure | 23 | _ | vs. | Calculated | Plasma | Concentrations | |
| III. | APPENI | DICES | | | | | | |
| | Append | | - | | | | | 69 |
| | Append | díx B | Subject | Cons | sent Form | | | 72 |
| | Append | dix C | Measured and Bl | | ncentrations | s of WR | 6026 in Plasma | 73 |
| | Append | dix D | | | ncentrations es in Urine | s of WR | 6026 and | 79 |
| | Append | dix E | | | | | | 83 |
| | Append | | - | | | | | 195 |
| | | | | | | | | |

88

8

23

Ħ

22

ñ

% &

Ž.

33

É

8

1. INTRODUCTION

Visceral leishmaniasis is a serious disease caused by the parasite

Leishmania donovani (1). Though it is ordinarily a zoonosis transmitted

between animals, especially rodents and canines through the bites of infected

sandflies, humans may also be infected. It is found in parts of Europe,

Africa, Asia, Central America and much of South America. The disease causes

prolonged debility, and left untreated it causes death in 75% to 90% of

patients. Cutaneous leishmaniasis is not lethal, but may cause substantial

morbidity (1). Not only is leishmaniasis a major world health problem, it is

also a problem for U.S. military forces stationed in areas where transmission

of the disease occurs.

The mainstay of therapy for leishmaniasis is pentavalent antimony. To produce a cure, this must be administered in repeated doses over a period of up to one month. While side effects are rare, their incidence is doserelated. Up to 15% of patients may relapse after antimony therapy and must be given extended antimony treatment or be treated with other drugs which are toxic (2). Pentamidine causes significant side effects including fatigue, anorexia, nausea, abdominal pain and prolonged hypoglycemia. Ten percent of patients treated with this drug develop permanent diabetes. Amphotericin B causes fever, fatigue, nausea, vomiting, anemia, uremia, and thrombophlebitis. Most significantly, it also produces permanent impairment of renal function in many patients. Given the significant failure rate of antimony compounds and the toxicity of other effective drugs, there is a clear need for development of alternative drugs.

WR 6026 (8-(6-diethylaminohexylamino)-6-methoxy-4-methylquinoline

dihydrochloride) (Figure 1) has been shown to be a highly active anti-leishmanial drug, both in an animal and an <u>in vitro</u> test system. Like primaquine, it is an 8-aminoquinoline. In hamsters infected with <u>L. donovani</u>, it produced significant suppression of infection in a dose of 0.025 mg/kg administered orally twice daily for four days, making it approximately 700 times more active than the reference antimonial compound pentostam (3). In the human macrophage-<u>L. tropica</u> test system, it was approximately 7 times more active than pentostam (4).

Subchronic toxicity studies of WR 6026 in rats given daily oral doses of 4.5, 9 or 18 mg/kg for 28 days revealed minor changes in blood counts and serum chemistry values including minor increases in methemoglobin levels (5). Target organs identified in these studies by histopathologic examination were lung and uterus at intermediate and high doses, and spleen, heart and kidneys at high doses. The abnormalities encountered in these organs included increased numbers of alveolar macrophages and eosinophilic granular material in alveolar spaces in the lung; hydrometra in the uterus; extramedullary hematopoiesis in the spleen; myocardial fibrosis in the heart; and protein-aceous casts along with abnormalities of tubular epithelium, including necrosis, in the kidney.

X

در

10 to

X

Subchronic toxicity studies of WR 6026 were conducted in dogs given daily oral maintenance doses of 0.3, 1 or 3 mg/kg for 25 days after loading doses of four times the daily maintenance dose on Day 1, three times the daily maintenance dose on Day 2, and twice the daily maintenance dose on Day 3. These studies revealed no significant changes in blood counts apart from a mild decrease in platelets after the first week of dosing with subsequent

return toward pre-treatment levels, and a mild increase in reticulocyte counts at mid- and high-dose levels (6). Methemoglobin levels increased with increasing drug dose. Control dogs had methemoglobin levels of less than 1%; the mean value for low-dose dogs was 1.4%, for mid-dose dogs was 8.8% and for high-dose dogs was 35%. Serum albumin levels were decreased mildly in mid-and high-dose dogs. Mild to moderate increases were also observed in serum globulin, SGOT, LDH, cholesterol, and triglycerides, and there was a 2-3 fold increase of haptoglobin levels. These animals also developed dose-related weight loss at the mid- and high-dose levels.

20

8

Histomorphologic abnormalities in mid- and high-dose dogs were found in the liver, spleen, heart, kidney and gallbladder. In mid- and high-dose dogs, there was increased extramedullary hematopoiesis in the spleen. In spleens of high-dose dogs, there was also congestion of the red pulp, as well as plasma cell infiltration and lymphoid depletion. In addition, high-dose dogs had bile duct hyperplasia, pleocellular periportal infiltrates, and vacuolated reticuloendothelial cells in the liver. Two dogs in the high-dose group had cytoplasmic degeneration of myocardium. In the kidneys of one high-dose dog there was tubular necrosis as well as brown granular pigment in tubular cells. The gallbladder in mid- and high-dose dogs showed mucosal hyperplasia associated with inspissated bile.

In a study comparing the cardiovascular and pulmonary effects of WR 6026 and primaquine in anesthetized dogs during intravenous infusion of the drugs, the major effects of WR 6026 at infusions of 1.0, 2.5, and 4.0 micromoles/kg/min were a weakening of ventricular contractility and a constriction of the pulmonary vasculature (7). These effects were more

significant at increasing infusion rates. In addition, there was short-lived prolongation of the P-R and Q-T intervals at the higher doses. No cardiac arrhythmias were noted.

Phase I testing of WR 6026 in 44 healthy male subjects given increasing single oral doses of up to 60 mg (60 mg given to only two subjects) revealed no significant drug-related symptoms or physical or laboratory abnormalities (8). Laboratory studies designed to detect the occurrence of hemolytic anemia revealed no significant differences in comparison with subjects treated with placebo. Methemoglobinemia did not occur in any subject.

8

B

In efficacy and toxicity studies in man conducted during World War II among volunteers infected with <u>vivax</u> malaria (Chesson strain), WR 6026, administered in an oral dose of up to 30 mg base daily concurrently with quinine (2.0 grams salt per day) for 14 days, was associated with 2.0-3.1% methemoglobinemia, low grade leukocytosis (WBC 12,000-14,000) and non-specific T wave changes on EKG (9). No other significant drug-related side effects were noted.

The purpose of this study was to determine the pharmacokinetics, safety and tolerance of a single 60 mg dose of WR 6026 in a larger group of healthy volunteers than previously studied at this dose. These data could then be used to design appropriate multiple-dose studies.

2. MATERIALS AND METHODS

2.1 WR 6026

WR 6026 (manufacturer's code WRA-20-05186, bottle #BK01845) was supplied by the Army as 15 mg capsules and was delivered to The Johns Hopkins Hospital Adult Medicine Pharmacy. Capsules were

prepackaged by the pharmacy study monitor in unit packets containing four 15 mg capsules, labelled by dose and subject name. Each subject was administered a single 60 mg (4 x 15 mg) dose.

2.2 SUBJECTS

A group of eight healthy male volunteers capable of providing written informed consent was recruited for study participation.

Institutional Review Board approval was obtained from both the Joint Committee on Clinical Investigations of The Johns Hopkins Medical Institutions and the Human Use Review Office of the U.S. Army.

2.2.1 Inclusion Criteria

Participants in this study were to be males between 18 and 35 years of age and within the Army's weight limit for height according to AR 600-9. Subjects were required to have no clinically significant medical condition as determined by a detailed medical history and physical examination performed by a physician. Serum chemistry profile, hematology and urine analysis were required to be within the normal ranges as defined by The Johns Hopkins Hospital Department of Laboratory Medicine. Chest x-ray within six months of entry and an electrocardiogram had to be normal. Prospective volunteers were required to be available and agreeable to be confined to the Clinical Pharmacology Research Unit for the entire study period.

2.2.2 Exclusion Criteria

Women were excluded from this study. Men were excluded if they did not meet the entrance criteria listed above (2.2.1) or if

they had a known or suspected allergy to antimalarial compounds or related drugs. Candidates requiring systemic medication were not eligible for study participation. Subjects with documented glucose-6-phosphate dehydrogenase (G6PD) deficiency, determined by a quantitative assay of enzyme activity with levels less than 7.4 IU/per gram of hemoglobin, or abnormal methemoglobin levels were excluded from study participation. Once accepted as a candidate for the study, subjects were not permitted to take any medication for 72 hours prior to admission into the study.

Χc

Ŋ

2.2.3 Recruitment

Advertisements were placed in the Help Wanted classified sections of the metropolitan Baltimore Sun newspaper. A special telephone line was dedicated to volunteer recruitment. Interested candidates were pre-screened on the telephone by research personnel who described the details of the study, took a brief medical history, and scheduled the candidates for additional screening examinations.

2.2.4 Informed Consent

Written informed consent was obtained from each subject prior to entering the study and was made a part of the subject's permanent study record. The informed consent described in detail the purpose of the study, the research protocol, and the associated potential risk. Each subject was advised that study participation was voluntary and that he could withdraw at any time. Each subject was afforded ample opportunity to ask

questions of the investigator prior to and after entering the study.

2.2.5 Compensation

Based on the duration of time the subject stayed in the hospital and the numerous venipunctures and urine collections required during the study, subject compensation was computed at \$250 per subject for completion of the study.

2.3 EXPERIMENTAL PROTOCOL

2.3.1 Objectives

The primary objective of this study was to determine the pharmacokinetics of a single oral 60 mg dose of WR 6026 in healthy adult male subjects. The information obtained from this study could then be used in the design and development of multiple-dose studies for WR 6026. Additionally, we evaluated the tolerance and toxicity of a single oral 60 mg dose of WR 6026 in these subjects.

2.3.2 Design

This study was an open-label design with each subject receiving a single 60 mg oral dose of WR 6026. The dose was given after an eight-hour fast and the subjects were allowed to resume eating four hours after the dose was given. The subjects swallowed the dose under the observation of study personnel.

The study examined each subject's tolerance of a single 60 mg dose of WR 6026 through regular, non-directed questioning regarding symptoms of adverse effects. Clinical laboratory

studies, including chemistry profiles, hematology tests, urine analysis, and methemoglobin levels, were monitored at regular intervals. Electrocardiographic measurements were also obtained at regular intervals before and after drug administration. The Study Flow Chart, showing the scheduled time of each event in the protocol, is Appendix A.

All subjects were screened as outpatients. Drug administration, blood and urine specimen collection, and post-drug tolerance and toxicity evaluations were performed during the inpatient hospital phase.

N.

X

2.4 CLINICAL LABORATORY EVALUATIONS

All laboratory evaluations, with the exception of assays for WR 6026 and its metabolites, were done within The Johns Hopkins Medical Institutions. Hematology and chemistry determinations were performed by the Department of Laboratory Medicine (Clinical Laboratory License No. 19-1054). Clinical laboratory tests were conducted at screening, two days prior to drug administration, immediately before drug administration and at 24, 48 and 96 hours post-dose.

2.4.1 Hematology

Routine hematologic determinations including hematocrit, hemoglobin, red blood cell count, white blood cell count with differential count, and platelet count were done.

2.4.2 Serum Chemistry

Serum was assayed for sodium, potassium, chloride, CO₂, urea nitrogen, creatinine, glucose, uric acid, calcium, phosphate,

total protein, albumin, direct and total bilirubin, aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase, lactic dehydrogenase (LDH), creatine kinase (CK), and methemoglobin levels. Cholesterol and fasting triglyceride tests were done 24 hours pre-dose and at 24, 48 and 96 hours post-dose.

2.4.3 Urine Analysis

Urine analysis was performed by personnel in the Division of Clinical Pharmacology. Protein, ketones, glucose, and bilirubin were measured qualitatively, and pH and specific gravity were determined. A microscopic examination of the sediment was also performed.

2.4.4 Electrocardiography

A standard 12-lead electrocardiogram (ECG) was performed on each subject at screening, just prior to drug administration, and at 4 hours, 1 day, 2 days, and 4 days after drug administration. All ECG's were reviewed and interpreted by a physician on the staff of The Johns Hopkins Hospital in the Division of Internal Medicine.

2.4.5 Chest X-rays

Standard postero-anterior and lateral chest x-rays were performed on each subject within six months of entry into the study. All x-rays were reviewed and formal reports provided by The Johns Hopkins Hospital Department of Radiology.

2.5 SPECIMEN HANDLING

2.5.1 Blood/Plasma Collection and Storage

Venous blood specimens were collected into heparinized glass tubes using an indwelling catheter fitted with a heparin lock. Blood specimens were centrifuged for 10 minutes and the plasma was decanted into plastic storage vials and frozen at -80°C. Whole blood samples were also obtained from Subjects #1 and #2, transferred to plastic vials, and frozen at -80°C. All blood and plasma specimens were labelled with the subject's initials and number, and the date and time of collection.

2.5.2 Urine Collection and Storage

For each urine collection period (as specified by protocol), the total urine volume was carefully measured and recorded. Aliquots from each collection were stored in three 60 ml plastic containers at -80°C. All urine specimens were labelled with the subject's initials and number, and the date and time period of collection.

2.5.3 Specimen Shipment

The frozen plasma and blood specimens were shipped in a sealed insulated container packed with dry ice. Shipment was made by an overnight carrier to Dr. Emil Lin at the University of California at San Francisco. The frozen urine specimens were transported in a sealed insulated container packed with dry ice. The urine samples were picked up by Army personnel and delivered to the laboratory of Dr. Anthony Theoharides at the Division of Experimental Therapeutics at WRAIR.

2.6 ASSAY OF WR 6026 AND METABOLITES

2.6.1 Assay of WR 6026 in Plasma and Blood

Concentrations of WR 6026 in plasma were assayed according to the method utilized by Dr. Emil Lin (10). Briefly, 0.5 ml of plasma was precipitated with 1.2 ml of CH3CN containing the internal standard WR 223658. The mixture was centrifuged and the supernatant applied to C2 Bond Elut cartridges which had been previously washed with 3 ml of 98% CH3CN containing 0.1% sodium lauryl sulfate (SDS), 6 ml of water and 3 ml of CH3CN. The cartridges were washed with 2 ml of 100% CH₃CN and 1 ml of CH₃CN containing 0.1% SDS. The WR 6026 was eluted with 3 ml of 98% CH₃CN containing 0.1% SDS into silanized tubes. Eluates were evaporated to dryness under nitrogen at room temperature. The residues were reconstituted in 200 ul of 80% CH3CN. Aliquots were chromatographed on an Altex C8 column using a mobile phase of acetonitrile:water (60:40) containing 0.2% SDS (wt/vol) and 0.2% glacial acetic acid (vol/vol) with the final volume adjusted to pH 5.5 with ammonium sodium lauryl sulfate hydroxide at a flow rate of 1.2 ml/min. Drug was detected by measuring absorbance at 263 nanometers.

The assay had a level of detection for the WR 6026 base of 6.44 ng/ml. The desethyl metabolite of WR 6026 (8-(6-ethylamino-hexylamino)-6-methoxy-4-methylquinoline dihydrochloride hemihydrate--WR 211789) can also be quantitated with this same method with a quantitation limit of 8.0 ng/ml. Precision of the assay, measured as the percent coefficient of variation of replicate spiked samples of WR 6026 in plasma, varied from

1.23% to 5.10% for inter-day and 2.78 to 10.2% intra-day analysis (10). Blood was also assayed, although the procedure was not validated. One-half ml of water was added to 0.5 ml of blood. The mixture was vortexed and then 1.2 ml of CH₃CN containing the internal standard and 1.0 ml of CH₃CN were added. The resultant mixture was vortexed for 20 seconds, sonicated for 10 minutes and centrifuged. The supernatant was treated identically to the plasma supernatants (10). Results of the analysis of blood samples must be regarded as preliminary since the method was not validated at the time the assays were performed.

X

Ø

69

2.6.2 Assay of WR 6026 and its Metabolites in Urine Urine was assayed for WR 6026 and its desethyl (WR 211789) and 4-hydroxymethyl (8-(6-diethylaminohexylamino)-6-methoxy-4hydroxymethylquinoline dihydrochloride hydrate--WR 254421) metabolites by Dr. Anthony Theoharides, Department of Pharmacology, Division of Experimental Therapeutics at WRAIR. The assay is detailed in his report (11). Briefly, the internal standard, WR 223658, was added to ten ml of urine. The mixture was applied to Baker 3 ml silica gel solid phase extraction columns which had been prewashed with 6 ml each of deionized water, methanol and deionized water. The colum. s were washed with 6 ml of 50% ethanol/deionized water. compounds of interest were eluted with 6 ml of 50% ethanol/0.1M ammonium acetate, pH 7.5, into clean extraction tubes and evaporated to dryness under nitrogen at 35°C. The residue was reconstituted in 200 microliters of methanol and

analyzed by HPLC.

Chromatography was performed using a 10 micron micro bond pack C-18 steel column from Water Associates with electrochemical detection using a glassy carbon electrode operated in the oxidative mode at an applied potential of +0.80 volts (Model TL-6A, BAS). The mobile phase used for the chromatography was 0.1M ammonium acetate, pH 4.5:acetonitrile (68:32) at a flow rate of 1.5 ml/minute. The method is accurate to within 7% for each of the three compounds, while the coefficients of variation range from 7% to 27%, depending upon both the compound and the concentration (11).

2.7 PHARMACOKINETIC ANALYSIS

Visual analysis of plots of the concentration of WR 6026 against time demonstrated that after a short lag period, plasma concentrations of WR 6026 rose and then gradually declined. In most subjects, the data appeared to be most consistent with a one-compartment model with first-order absorption following a time lag for absorption to begin, and with first-order elimination. In some instances, the last one or two measured concentrations appeared to lie above the expected concentrations if a single elimination phase were present, suggesting the possibility of biphasic elimination. PC NONLIN^R, a commercially available curve fitting program (12), was used to estimate the pharmacokinetic parameters for each patient assuming a single compartment with first-order absorption after a time lag and first-order elimination. Analyses of the data from several patients were performed weighting the concentration at each time point equally.

This method estimated poorly the low plasma concentrations, i.e., those occurring early after dosing and those greater than 24 hours after dosing. Accordingly, analyses were performed which weighted the data proportionally to the inverse of the square of the concentration. This technique provided better estimates of both low and high plasma concentrations. However, the peak concentrations were consistently underestimated. A two-compartment model was tried to see whether the data could be fit better with this model. Using this model, peak concentrations were estimated poorly, and the standard errors of the parameter estimates were very large, suggesting that there were insufficient data for the model. Attempts were made to fit the data to a two-compartment model with MK MODELR, another commercial curve-fitting program (13). Large standard errors of the estimates were obtained using this program as well. RSTRIPR, a pharmacokinetic data stripping program with least squares parameter optimization (14), was also tried. RSTRIPR produced parameter estimates to approximate the data with equations containing two and three exponential terms (equivalent to one- and two-compartment models with first-order absorption). Once again, error estimates of the variables were large for the two-compartment fits, indicating that the data were insufficient to describe a model of this complexity. Accordingly, PC NONLINR was used to fit the plasma and blood concentrations to a one-compartment model with data weighted to the inverse square of the concentration since this model provided the best approximation of the observed data. The amounts of WR 6026 excreted into the urine unchanged were used to estimate the elimination rate constant of WR 6026 from the plasma. WR 6026 was

蒙

3

3

K

Z

33

8

X

not detected in the urine collected from 84 to 96 hours after dosing except in Subject #5, in whom less than 2% of the total drug excreted unchanged was present. This suggested that the excretion of unchanged drug was complete at the end of the collection period. The natural logarithm of the amount of unchanged drug remaining to be excreted (X_{U}^{ω} - X_{U}^{t}) was plotted against time (15). Unweighted linear regressions were performed to estimate the slope of the line, which is equal to the negative of the elimination rate constant of WR 6026 from the plasma.

The single exponential decline of the plot of the urinary data suggested that all the data, the amount of unchanged WR 6026 excreted in the urine and the plasma concentrations of WR 6026, could be used together to model the pharmacokinetics of WR 6026. PC NONLINR was used to estimate the parameters of a one-compartment model with first-order absorption after a lag time and first-order elimination (including both urinary excretion and metabolism) which best described the plasma concentrations and urinary excretion of WR 6026 for each subject. Estimates were obtained when weighting each observation equally and also using the reciprocals of the observations as the weights for each data point.

3. RESULTS AND DISCUSSION

3.1 AMENDMENTS AND COMPLIANCE

3.1.1 Amendments

The consent form was amended prior to enrolling subjects to provide information concerning:

- 1. The fact that WR 6026 is not currently licensed.
- Follow-up on significant changes in clinical laboratory tests.
- 3. The risks of a heparin lock.
- 4. The right of Army inspectors to review relevant records.
- The treatment of any injuries incurred in this project.
 The amended consent form is provided as Appendix B.

3)

M

3

 Ξ

3

7

N

The original protocol scheduled clinical chemistry and hematology evaluations at screening, on admission to the hospital (Day 1) and on Days 4, 5, and 6. In reviewing this schedule we realized that we did not have measurements immediately prior to WR 6026 administration to use for comparison with the post-drug samples, and therefore added a zero time point for a more reliable baseline. There was a discrepancy between the text of the protocol and the flow sheet regarding the timing of CK measurements. We decided, after conferring with Army personnel, that CK should be measured at screening and at time points specified on the flow sheet.

In the original protocol the repeated urine analyses eliminated some urine from the timed collections. For this reason we stopped doing urine analyses after drug administration until the final urine collection was completed. This change was made while the second volunteer was in the midst of the protocol. Therefore, urine analyses were done per original protocol on the first subject. The second volunteer had a urine analysis on Day 4, but not on Day 5.

Eliminating the urine analysis scheduled for Days 4 and 5 allowed us to keep the continuous sequential urine collections intact after WR 6026 administration without sacrificing even a small part for urine analysis. We obtained a urine analysis after the final 12-hour urine collection was completed on Day 7 for safety data.

The lipid profile was changed from Day 2 to zero hour on Day 3. This was only a time change, and still occurred prior to drug administration.

It was initially intended to measure WR 6026 concentration in whole blood and plasma in all of our subjects. However, whole blood samples, in addition to plasma and urine, were obtained and saved only from the first two subjects. At the request of Army representatives, only plasma and urine specimens were collected for WR 6026 concentration from the remainder of our subjects.

All of the amendments mentioned above were authorized by Army personnel prior to implementation.

3.1.2 Compliance

Subjects were enrolled into the study even if screening and/or admission CK and triglyceride levels were elevated. We believe the CK elevations do not reflect a myopathic process, but rather the active physical lifestyles of the subjects. Creatine kinase levels increase with exercise, and athletes have been reported to have higher baseline levels of serum CK as their "normal" (16-23). The elevated triglyceride levels

were not far enough above the "normal range" to be considered significant. We are satisfied that these subjects were healthy and that those with elevated CK and triglycerides did not have a pathological process.

One subject (#5) who was entered into the protocol exceeded the AR 600-9 weight limit for height. He was 27 years old, 6'5" tall and weighed 231 pounds. The upper limit according to AR 600-9 for individuals age 26-39 of that height is 229 pounds. Another weight criterion often used in our healthy volunteer studies is that subjects must be within 10% of the ideal body weight specified by the Metropolitan Life Insurance table. Our Subject #5 did fall within 10% of his ideal body weight for height as listed in that table. Furthermore, this individual was very large and well-muscled, and his "excessive weight" was not due to obesity.

X

V

Three subjects were discharged but not recalled when the discharge laboratory tests returned showing that the LDH had risen above the upper limit of normal. In two of these individuals the elevation was deemed trivial and not pursued. In the other subject the LDH had risen to nearly twice the upper limit of normal. Although the subject was asymptomatic and this was an isolated abnormality in his clinical laboratory tests, the test should have been repeated and our failure to repeat the test after discharge was an oversight and deviation from the protocol.

3.2 DESCRIPTION OF SUBJECT POPULATION

During the volunteer recruitment period, 49 men responded to the newspaper advertisements. Of these, 6 failed to keep scheduled appointments, while 32 were disqualified for not meeting one or more of the inclusion/exclusion criteria. Eleven subjects were examined by the physician and qualified for study participation. Three of these were extras and were not used. The study group was comprised of eight healthy black males. The average age was 26.3 years, ranging from 20 to 34 years. The average height and weight for the study group were 179.4 cm (range 165-195) and 75.5 kg (range 54.5-105), respectively. Relevant characteristics for individual subjects are presented in Table 1.

3.3 CLINICAL RESULTS

3.3.1 Symptomatic

All of the volunteers tolerated the administration of WR 6026 very well. There were no symptomatic complaints.

3.3.2 Laboratory (Table 2)

3.3.2.1 Liver Function Tests

Two subjects had an increase in serum AST to a level less than twice the upper limit of normal on post-drug day 4, having had normal levels at each time tested prior to that day. These elevations were not associated with symptoms and only one of the two had a corresponding increase in serum ALT on the same day. In both of these subjects the subsequent serum AST levels on days 5 and 7 were within normal limits.

Two other subjects had a minimal elevation of AST or ALT (less than twice the upper limit of normal) just before the drug was administered, even though the screening levels had been normal. In these two cases the transferases remained minimally elevated, but because the initial elevation preceded the administration of the drug it was not attributable to a drug effect. There were no elevations in alkaline phosphatase. Four subjects had increases in LDH above the normal limit, three just slightly over the upper limit of normal and one reaching nearly twice the upper limit of normal. Three of the elevations were noted on post-drug day 4, while the other was present on post-drug day 2 and was back within the normal range on post-drug day 4. These modest LDH elevations were unassociated with symptoms. Inasmuch as these LDH and transferase elevations were rather mild and usually occurred four days after drug administration, it is not clear whether they were related to the drug or whether this simply represented laboratory variability. Because the elevations were an isolated finding and unassociated with symptoms, or were only slightly elevated, no follow-up blood tests for LDH were performed in the three subjects whose levels were increased on the day of discharge.

K

7

3.3.2.2 Creatine Kinase

Several of the subjects had an elevated CK at

screening and at entry into the hospital, but in no case did the CK level increase after the administration of WR 6026 to a level higher than the admission value. This is consistent with previous experience in our subjects, where CK was found to be elevated with no symptoms whatever, and, as mentioned in 3.1.2, probably represents release of this enzyme from muscle tissues as a result of a very active, physical life style rather than from a myopathic process.

3.3.2.3 Triglycerides

Only one subject developed an increase in measured serum triglycerides above the upper limit of normal. This was on the second day following drug administration. Because of a laboratory instrument malfunction, the test was not repeated on the fourth day following drug administration.

3.3.2.4 Hematological Tests

No subject had a significant change in hematocrit, white blood count, differential count or platelet count during the study period following administration of WR 6026.

3.3.2.5 Electrocardiograms

Electrocardiograms in these subjects did not show a significant change at four hours after dosing (presumably near maximum plasma levels) or during the

period of four days after drug administration.

Minor, usually non-specific abnormalities were

sometimes present prior to drug administration; nonspecific T-wave changes developed in two subjects

after drug administration, possibly attributable to

lead placement in one of the two. In neither case

was the magnitude of change felt to be of any

significance. Furthermore, there were no symptoms of
cardiac dysfunction.

3.3.2.6 Methemoglobin

No subject in this study had an increase in methemoglobin above the normal range for our laboratory after administration of WR 6026.

N

3.3.3 Clinical Conclusions

The subjects appeared to tolerate the administration of WR 6026 very well. No symptoms whatever were reported and none of the minimal abnormalities of the serum chemistries noted above were of clinical significance. We believe that the single dose of WR 6026 was well tolerated and that further testing can be pursued. Nevertheless, during future studies it is important to continue close monitoring of the liver function tests, triglycerides, LDH, methemoglobin and electrocardiograms, not only by virtue of the minimal abnormalities seen in this project, but also because of good clinical research practice and the experience in laboratory animals.

3.4 PHARMACOKINETIC RESULTS AND DISCUSSION

The plasma and blood concentrations of WR 6026 measured in each of the subjects at each sampling time are listed in Appendix C (10). The plasma concentrations also are tabulated according to the scheduled time of sampling in Table 3. The excretion of the parent drug and two metabolites, 4-OH WR 6026 (WR 254421) and desethyl WR 6026 (WR 211789) into the urine was quantified in the eight subjects by Dr. Theoharides at WRAIR (11). Tables copied from his report are contained in Appendix D of this report. Table 1 of Appendix D details the accuracy and precision of the assays for WR 254421, WR 211789, and WR 6026 in urine. Tables 2, 3 and 4 of Appendix D list the concentrations in each urinary sample, the volume of each sample and the amount excreted in each time period. All of the clinical results and drug concentrations for each subject are included in the Case Report Forms provided in Appendix E.

Plasma concentrations of WR 6026 measured in the eight subjects, as well as simple descriptive summary statistics, are listed in Table 3. Plasma concentrations remained below the minimal detectable concentration of 6.44 ng base/ml for a variable period of time after dosing and then rose gradually before declining. In one subject, #5, drug was not measurable until the 1.25 hour sample, while in five of the subjects drug was measurable at 0.5 hours. The time at which the peak concentration occurred also varied, ranging from 1.5 to 5.0 hours, with the average being 3.0 hours (Table 4). Peak concentrations varied from 52.5 to 116.0 ng/ml with a mean of 73.5 ng/ml. After peaking, concentrations fell gradually. Plasma concentrations remained above the minimum detectable level at 24 hours in seven sub-

jects; in two, #6 and #7, drug was still measurable at 48 hours (Table 3). Drug was not measurable in the plasma of any of the subjects 60 hours after dosing or later. Graphs of the plasma concentrations in the individual subjects are provided in Figures 2-4.

The average WR 6026 plasma concentration at each sampling time is listed in Table 3 and shown in Figure 5. The average concentration rose slowly after a short lag time. A plateau was reached at about 60 ng/ml lasting from two hours after dosing until five hours after dosing before drug levels declined. Of note, no desethyl WR 6026 was detected in the plasma specimens (10).

X

V

Whole blood as well as plasma concentrations of WR 6026 were measured after dosing Subjects #1 and #2 and are listed in Appendix C. No statistics have been calculated since values were obtained in only two subjects. Concentrations in blood were lower than those in plasma (Figure 6 compared to Figure 2) indicating that the concentration of drug in the cellular components of blood, primarily red cells, is lower than the concentration of drug in the plasma.

Concentrations in both were low, only nanograms per milliliter, after a 60 mg dose, suggesting that one or more of the following occurs: poor absorption, widespread distribution, or presystemic elimination. The data were insufficient to distinguish among these possibilities.

The fraction of the dose of WR 6026 recovered in the urine in 96 hours as unchanged drug was small and variable between subjects (Table 5). Hydroxymethyl WR 6026 (WR 254421) and desethyl WR 6026 were also recovered in the urine (Table 5). In each subject, more hydroxymethyl WR 6026 was recovered than WR 6026, which in turn was

more than the amount of desethyl WR 6026. In fact, of the total amount of drug recovered as these three moieties, 76 to 96% was recovered as hydroxymethyl WR 6026. There was at least a fivefold variation in the total amount of each compound (in micromoles) excreted in the urine in 96 hours. In addition, the amounts of these compounds present in the urine collected 84 to 96 hours after dosing were small, indicating that an insignificant amount of these compounds remained to be excreted; i.e., urinary excretion was virtually complete by 96 hours. The mean fraction of the dose recovered as unchanged drug and the two metabolites was 0.141 with a standard deviation of 0.075. Recovery ranged from 6.2 to 30.0 percent of the dose (Table 5). Thus, the majority of the administered drug cannot be accounted for by the drug and the two metabolites measured in the urine in the 96 hours after dosing.

A standard pharmacokinetic model which produced good estimates of the observed plasma data could not be found. Although the rise of plasma concentrations to a peak followed by a decline suggested a one-compartment model with first-order absorption and elimination, this model did not accurately estimate the peak plasma concentrations. Changes in the weighting of the data failed to correct severe underestimates of the peak concentrations, although when the data were weighted to the inverse square of the concentrations, the low concentrations were estimated rather well. A two-compartment model did not describe the data better; large standard errors of the model's parameter estimates were obtained, suggesting that an insufficient number of data points were present to describe the model. Therefore, a one-compartment model with data weighted to the inverse square of

the concentration was used to describe the data, realizing that the difficulties in measuring low concentrations of drug (those less than 6.4 ng/ml) and in estimating peak values serve to produce pharmacokinetic constants for each patient that are, at best, approximate. The pharmacokinetic constants do, however, provide a way to discuss quantitatively the pharmacokinetic processes and the inter-subject differences. The measured concentrations and the plasma concentrations predicted by the curve-fitting process for each subject are shown in Figures 7 through 14.

Measured peak plasma concentrations are tabulated in Table 4 while the estimates of the peaks derived from the curve fitting are listed in Table 6. The measured maximum concentrations varied by a factor of two in the eight subjects. The observed peaks were severely underestimated by the model. Furthermore, the calculated times to peak did not agree with the times at which the maximum concentrations of WR 6026 were actually observed.

The areas under the plasma concentration-time curve (AUC) also varied widely between subjects (Table 7). As measured by the trapezoidal rule, there was a fourfold difference in the AUC in the eight subjects; the mean AUC was 1066 ng·hr/ml with a range of 512 to 2179 ng·hr/ml. The coefficient of variation was 0.47. The curve fitting process produced areas under the concentration-time curves which closely approximated those obtained by the trapezoidal rule (Table 7). In 7 of the 8 subjects, the estimate of the area under the concentration-time curve was within 5% of the value calculated by trapezoidal rule. In the other subject, #8, a 12% difference

Ŋ

K

-

occurred.

The model parameters estimated from the plasma data are listed in Table 4. Each parameter had at least a twofold range in the eight subjects; coefficients of variation for the mean estimates were 27% to 57%. The average time before absorption was initiated was 0.55 hours with a standard deviation of 0.26 hours. The time to onset of absorption varied from 0.30 to 0.89 hours (Table 4). The mean rate constant for drug absorption was 0.88/hour with a coefficient of variation of 0.40; estimates ranged from 0.44/hour to 1.36/hour. These extremes are equivalent to calculated absorption half-times of 1.56 and 0.51 hours, respectively (Tables 4 and 6).

Estimates of the rate at which WR 6026 disappeared from the plasma also differed between subjects. The mean rate constant of elimination, K10, was 0.082/hr with coefficient of variation of 0.57. The greatest elimination rate constant was 0.192/hour in Subject #2, equivalent to an elimination half-time of 3.61 hours, while the slowest elimination occurred in Subject #3; his rate constant of elimination was 0.048/hour which translates to an elimination half-life of 14.52 hours (Tables 4 and 6).

Analysis of the data on unchanged WR 6026 in the urine provided another way of estimating the rate of elimination of WR 6026 from the plasma. Although the amount of unchanged WR 6026 recovered in the urine was small, ranging from 0.53 to 4.57 micromoles, or 0.4% to 3.2% of the administered dose, the time course of excretion was used to estimate the rate constant of elimination of WR 6026 from the plasma. The "amount of drug remaining to be excreted in the urine"

method was used to estimate the elimination rate constant, K10, and the half-life of WR 6026 in each subject. The plot demonstrated a single exponential decline, providing only the terminal elimination rate constant. The values obtained for each subject and the summary statistics are listed in Table 8. The rate constants are smaller and the half-lives longer than the estimates obtained from the modeling of the plasma concentration data in each subject except for Subject #7. Nearly a threefold variation in the parameter estimates occurred in the eight subjects. The mean rate constant of elimination, estimated from the urinary data, is 0.060/hour compared to the mean estimate of 0.082/hour from the plasma data. The mean (geometric) WR 6026 half-life is 11.6 hours when using the urinary data and 8.4 hours when estimated from the plasma concentration data.

Cal

7

3

3

8

[] :->

The plasma and urine WR 6026 concentration data in combination were then used to estimate the parameters for the one-compartment model. Since the magnitude of the amounts of drug in the urine was higher than the plasma concentrations, three different methods for weighting the observed data were tried. Data were weighted equally (wt = 0), to the reciprocal of the value (wt = -1), and to the square of the reciprocal (wt = -2). The parameter estimates for the one-compartment model are displayed in Table 9 (in conjunction with the estimates obtained from the plasma data and from the urine data alone). Each method of weighting estimated the total amount of the drug recovered in the urine accurately (Table 9) as well as the time course of urinary excretion (see Figure 15 for example of fit). However, all three consistently underestimated the peak plasma concentrations. Using equal weights for each observation produced

the highest estimates of the peaks. These, however, were still much lower than the observed peaks (Table 9). Furthermore, the low concentrations were estimated least well with this weighting technique (see Figure 15), and the calculated areas under the plasma concentration-time curve had the greatest deviation from the areas under the concentration-time curve calculated with the trapezoidal rule using the plasma concentration data with extrapolation to infinite time by dividing the last measurable concentration by the estimate of the elimination rate constant (Table 9). The best estimates of the low concentrations were produced by weighting each observation by the square of its reciprocal (see Figure 15 for example); calculated areas generally were in good agreement with the areas determined by trapezoidal rule (Table 9). The observed data and calculated estimates using this "wt = -2" method are depicted in Figures 16 through 23.

As indicated in Table 9, using a one-compartment model and weighting each data point by the square of its reciprocal produced estimates of the area under the concentration-time curve which were within 10% of the AUC calculated using the trapezoidal rule. In addition, the model estimated accurately the amount of WR 6026 excreted unchanged into the urine.

The pharmacokinetic parameter estimates for the one-compartment model for each subject and the summary statistics are listed in Table 10.

The calculated maximum concentrations and the calculated times at which the maximum concentration occurs differed widely from the measured peak concentrations and the times these peaks were actually

observed (Table 9). The plots of the measured plasma concentrations and the estimated concentrations demonstrate large discrepancies near the time when the peak occurred, while both the initial rise in the concentration and the later declines in concentration appear to be estimated quite closely. This suggests that the one-compartment model is not sufficiently complex to describe the late absorption of WR 6026 and its distribution throughout the subject. The mean estimates of the absorption rate constant, K10, time before onset of absorption, TLAG, and volume of distribution divided by the bioavailability, VOLUME/F, are similar to those obtained using only the plasma data (Table 4). As before, estimates of each parameter varied widely among the eight subjects.

The elimination rate constants (K10) obtained with the "weight ~ -2" procedure were similar to those obtained using the urinary data alone (Table 9) and were smaller than the estimates obtained from the plasma data alone in seven of the eight subjects. The mean rate constant of elimination, estimated using plasma and urinary data together, was 0.065/hour, equivalent to an elimination half-life of 10.7 hours. Six subjects had half-lives longer than 10 hours; the other two eliminated drug more rapidly. There was a threefold difference between the smallest and largest estimate. The plasma concentrations at 24 hours and later and the amounts of WR 6026 excreted into the urine for each subject are estimated closely by the individual K10's, suggesting that the terminal rate of elimination of WR 6026 from the body is described well by the model, in contradistinction to the late absorptive and distributive phases.

Thus, in the eight subjects studied, WR 6026 absorption began after a modest delay which averaged about 0.4 hours and proceeded slowly. Peak drug concentrations were not reached until 1.5 to 5 hours after dosing and drug was eliminated with a mean half-life of about 10.7 hours. Inter-subject differences in both the absorption and elimination rates resulted in large variations in the peak concentrations achieved and in the areas under the concentration-time curves. The parameter estimates, based on a compartmental analysis of the data, must be interpreted loosely since the compartment model did not accurately quantify the late absorption and early disposition, i.e., distribution, of the drug. Peak plasma concentrations were systematically underestimated by the model, suggesting that the absorption rate may be more rapid than estimated and that a rapid, distribution phase may be present as well. The sensitivity of the plasma assay limited the number of data points obtained at late sampling times, precluding analysis using a more complicated model. However, hints of a slow, terminal elimination phase appeared in the lowest concentrations measured. The simple model employed did estimate accurately the area under the plasma concentration-time curve, plasma concentrations after 24 hours and the amount of WR 6026 excreted into the urine unchanged.

Quantification of the amounts of parent drug and of two metabolites excreted in the urine over 96 hours accounted for a minor fraction of the administered oral dosage. Incomplete absorption, non-renal elimination or the presence of other metabolites, not measured using the techniques employed in this study, could account for the low

recovery of drug in the urine.

4. <u>CONCLUSIONS</u>

The eight volunteers studied in this project tolerated a single oral dose of 60 mg of WR 6026 very well with no symptomatic complaints. Only minimal and clinically insignificant changes in the serum chemistries were noted. Detectable WR 6026 absorption was delayed for a short period of time, about one-half hour, and proceeded slowly; peak concentrations occurred at about three hours, and varied by more than twofold in just 8 subjects. There was approximately a fourfold variability in the areas under the concentration-time curves. Compartmental analyses of the data were not optimal, but suggested that the terminal elimination half-life averaged about 11 hours and might be longer if a more sensitive assay were available.

These data suggest that further evaluation of WR 6026 in human subjects is appropriate. We believe that the next step should be multiple-dose testing in healthy subjects with close monitoring of symptoms, clinical Paboratory tests, and plasma and urine concentrations of drug. Dosing once a day, or about every two half-lives, would appear to be a reasonable approach. Higher plasma levels which might be achieved during a multiple-dose investigation and later sampling times would enable a more accurate determination of the pharmacokinetics of the compound. In particular, a multiple-dose study could determine whether a single-compartment model is sufficient to describe the data or whether a slower elimination phase is present which would lead to accumulation of drug if the dosing schedule were once every one or 1.5 half-lives. In addition, further studies on the metabolism of WR 6026 should be performed, addressing two issues in

particular: (1) whether a metabolite of the parent drug--possibly 4-OH WR 6026--is present in significant amounts in plasma, and (2) the identification of the fate of the administered drug, approximately 85% of which was not accounted for in the present investigation. Possibilities which need to be examined include determining whether other as yet unidentified metabolites are excreted in the urine, such as glucuronides of either the 4-OH or desethyl metabolites, and whether there is significant loss of drug in the gastrointestinal tract.

CONTROL CONTRO

REFERENCES

W

Z

8

- Locksley RM and Plorde JJ, Leishmaniasis, in <u>Principles of Internal Medicine</u>, llth edition, Braunwald E, Isselbacher KJ, Petersdorf RG, et al., eds, McGraw-Hill, New York, 1987, pp 785-787.
- 2. Anabwani GM, Dimiti G, Ngira JA, and Bryceson ADM. Comparison of two dosage schedules of sodium stibogluconate in the treatment of visceral leishmaniasis in Kenya. Lancet I:210-213, 1983.
- 3. Kinnamon KE, Steck EA, Loizeaux PS, Hanson WL, Chapman WL, Jr, and Waits VB: The antileishmanial activity of lepidines. Am J Trop Med Hyg 27:751-757, 1978.
- 4. Berman JD and Lee LS: Activity of 8-aminoquinolines against <u>Leishmania</u> tropica within human macrophages in vitro. Am J Trop Med Hyg 32: 753-759, 1983.
- 5. Reno FE, Trutter JA, Alsaker RD, Burdock GA, Voelker RW, and Hepner KE: Twenty-eight day subacute toxicity study in rats WR 6026 2HCl. U.S. Army Medical Research and Development Command Contract No. DAMD 17-81-C-1138, 1982.
- 6. Reno FE, Trutter JA, Hagen WH, Dawkins BG, and Alsaker RD: Twenty-eight day subacute toxicity study in dogs: WR 6026 2HCl in dogs. U.S. Army Medical Research and Development Command Contract No. DAMD 17-81-C-1138, 1982.
- 7. Caldwell RW and Nash CB: Comparison of cardiovascular and pulmonary effects of WR-6026-2HCL and primaquine diphosphate. U.S. Army Medical Research and Development Command Contract No. DAMD 17-82-C-3011, 1984.
- 8. Reba RC, Barry KG, and Altstatt LB: WR 6026 2HCl: Short term dosage safety and tolerance study: single oral dose, rising dose levels. WR 6026 IND Supplement 1.
- 9. Alving AS, Pullman TN, Craige B, Jr, Jones R, Jr, Whorton CM, and Eichelberger L. The clinical trial of eighteen analogues of pamaquin (plasmochin) in <u>vivax</u> malaria (Chesson strain). J Clin Invest 27(Supp): 34-45, May 1948.
- 10. Lin ET, Benet LZ, Upton RA and Gee WR. Single-Dose Absorption and Pharmacokinetics of WR 6026 Hydrochloride in Healthy Subjects. Report No. AY-86-2-D, U.S. Army Medical Research and Development Command Contract No. DAMD 17-86-C-6150, June 24, 1987.
- 11. Theoharides AD, Kim M, Ashmore RW, and Shipley L. Identification and Quantitation of WR 6026 and Metabolites in Human Urine. Biochemical Pharmacology Report 1987-1, Department of Pharmacology, Division of Experimental Therapeutics, WRAIR, August, 1987.

- 12. Metzler CM and Weiner DL. PC NONLIN. Version Ol.A. Statistical Consultants, Inc., Lexington, Kentucky, 1985.
- 13. Holford N. MK MODEL. Elsevier Science Publishers, Amsterdam, The Netherlands, 1986.
- 14. Fox JL and Lamson ML. RSTRIP. Version 3. Micromath, Inc. Salt Lake City, Utah, 1986.
- 15. Gibaldi M and Perrier D. <u>Pharmacokinetics</u>, 2nd ed., Marcel Dekker, Inc., New York, 1982, p. 5-10.
- 16. Wolf PL, Lott JA, Nitti GJ, and Bookstein R. Changes in serum enzymes, lactate, and haptoglobin following acute physical stress in international-class athletes. Clin Biochem 20:73-77, 1987
- 17. Occhi G, Gemma S, Buselli P, and Miserocchi G. Effects of repeated endurance. J Sports Med 27:184-190, 1987.
- 18. Clarkson PM, Apple FS, Byrnes WC, et al. Creatine kinase isoforms following isometric exercise. Muscle Nerve 10:41-44, 1987.
- 19. Stendig-Lindberg G, Shapiro Y, Epstein Y, et al. Changes in serum magnesium concentration after strenuous exercise. J Am Coll Nutr 6:35-40, 1987.
- 20. Nicholson GA, Morgan GJ, Meerkin M, et al. The effect of aerobic exercise on serum creatine kinase. Muscle Nerve 9:820-824, 1986.
- 21. Evans WJ, Meredith CN, Cannon JG, et al. Metabolic changes following eccentric exercise in trained and untrained men. J Appl Physiol 61:1864-1868, 1986.

POSSESSE TESTOCOL TESTOCOLOGICAL TOSSESSE TESTOCOLOGICAL TESTOCOLOGICA DESCRIPTION DE LA COLOGICA DEL COLOGICA DE LA COLOGICA DEL COLOGICA DE LA COLOGICA DEL COLOGICA DE LA COLOGICA DE LA COLOGICA DEL COLOGICA DE LA COLOGICA DEL COLOGICA DE LA COLOGICA DEL COLOGICA DEL COLOGICA DE LA COLOGICA DEL COLOGICA DE LA COLOGICA DE LA COLOGICA DEL COLOGICA DEL

- 22. Stansbie D, Aston JP, Dallimore NS, et al. Effect of exercise in plasma pyruvate kinase and creatine kinase activity. Clin Chim Acta 132:127-32, 1983.
- 23. Newham DJ, Jones DA, and Edwards RHT. Large delayed plasma creatine kinase changes after stepping exercise. Muscle Nerve 6:380-385, 1983.

TABLE 1
Individual Subject Characteristics

| Subject No. | I.D. | Age | Race | HT (cm) | WT (kg) | PPD* | EtOH** |
|-------------|--------|------|------|---------|---------|------|--------|
| 01 | E-W | 22 | В | 176.0 | 73.0 | 0.5 | 3 |
| 02 | RLH | 28 | В | 178.0 | 73.6 | 0.5 | 0 |
| 03 | W-F | 31 | В | 188.0 | 94.0 | 0.5 | 0.5 |
| 04 | LMH | 24 | В | 165.0 | 54.5 | 0.5 | 3.5 |
| 05 | DLA | 27 | В | 195.0 | 105.0 | 0.5 | 0 |
| 06 | M-P | 20 | В | 176.0 | 70.5 | 0.5 | 0 |
| 07 | AID | 24 | В | 179.5 | 57.5 | 1.0 | 49 |
| 08 | CLH | 34 | В | 178.0 | 76.0 | 0.5 | 6 |
| | MEAN | 26.3 | | 179.4 | 75.5 | | |
| | (± SD) | 4.7 | • | 8.9 | 17.0 | | |

^{*} Average packs of cigarettes smoked per day (from subject's history)

^{**}Ounces of alcohol per week (from subject's history)

TABLE 2

Peak Clinical Laboratory Abnormalities Developing

After WR 6026 Administration

(number in parenthesis indicates onset of abnormality in days after drug administration)*

[number in brackets indicates value just before drug administration]

8

| Subject # | AST <u>(units/l)</u> | ALT (units/1) | LDH (units/l) | Trig. (mg/dl) | T wave changes |
|-----------|-------------------------|------------------|------------------|------------------|----------------|
| 1 | | | 217 (4) [150] | | (0) |
| 2 | | | 394 (4) [145] | | |
| 3 | 41 (4) [23] | | 201 (4) [147] | | |
| 4 | | | | | (1) |
| 5 | | 44 (0)** [39] | 203 (2) [138] | | |
| 6 | 53 (4) [18] | 49 (4) [14] | | 255 (2) [86] | |
| 7 | 57 (0)** [37] | 63 (2) [28] | | | |

Normal ranges: AST 0-35 units/1

ALT 0-30 units/l

LDH 0-200 units/1

Triglycerides 20-190 mg/dl

- * The peak abnormality did not always occur on the day of onset of the abnormality.
- ** These two subjects were normal at screening but had slight transferase elevations on the morning blood sample taken just before drug administration.

<u>የወቅያዘመንያመው የናውነው የችን የራቀናቸው የአገድናው የመጀመርው የመጀመርው የ</u>መቀር እስከላይ የመፈርስ የውዲከተው የውዲስ አገር እንዲስፈው የውዲስ የመፈርስ የመፈርስ የመፈርስ

PLASMA CONCENTRATIONS (in ng free base/ml) OF WR6026 IN 8 VOLUNTEERS AFTER A SINGLE 60 MG ORAL DOSE

Table 3

| | | 9 | CHEDULE | D TIME | OF SAME | PLE (HO | URS AFT | ER DOSI | NG) | | | |
|-------|---------|-------|---------|--------|---------|---------|---------|---------|---------------|-------|-------|--------|
| | 0.00 | 0.25 | 0.50 | 0.75 | 1.00 | 1.25 | 1.50 | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 |
| SUBJ | | | | | | | | | | | | |
| 1 | | | | | 8.23 | 29.80 | | | 51.00 | | 57.20 | 49.20 |
| 2 | | | 7.26 | 30.50 | 50.90 | 60.30 | | | 67.00 | | 69.50 | 59.80 |
| 3 | | | | | 7.47 | 13.00 | 14.70 | 28.90 | | | 36.60 | 48.30 |
| 4 | | | 12.90 | 18.00 | 37.80 | 51.20 | 59.60 | | | | 68.30 | 62.80 |
| 5 | | | | | | 10.60 | 19.40 | | | 48.90 | 53.30 | 63.40 |
| 6 | | | 8.58 | 13.70 | 24.00 | | | | 42.90 | | 54.50 | 60.40 |
| 7 | | | 7.17 | 14.40 | 35.90 | | | | 102.00 | | | 102.00 |
| 8 | | | 7.63 | 13.20 | 14.20 | 16.30 | 45.30 | 65.20 | 60.50 | 64.40 | 47.30 | 66.90 |
| | 0 | 0 | 5 | 5 | 7 | a | ٥ | ۵ | 8 | 8 | 8 | 8 |
| HEAN | U | v | 8.71 | 17.96 | 25.50 | 35.48 | | | | 62.91 | 60.20 | |
| SD | | | 2.41 | 7.26 | 16.62 | 23.44 | | | | | 17.60 | 16.69 |
| CV(X) | | | 27.66 | 40.41 | 65.19 | | | | 41.73 | 22.65 | 29.24 | 26.04 |
| MAX | | | 12.90 | 30.50 | 50.90 | | | | 102.00 | | | 102.00 |
| HIN | | | 7.17 | 13.20 | 7.47 | 10.60 | | | 23.20 | | 36.60 | |
| | 5.00 | 6.00 | | | OF SAM | | | | ING) 60.00 | 72 00 | 00 48 | 00.49 |
| SUBJ | 3.00 | 0.00 | 0.00 | 10.00 | 12.00 | 24.00 | 38.00 | 70.00 | 00.00 | 72.00 | 57.00 | 70.00 |
| 1 | 44.40 | 45.90 | 31.40 | 32.30 | 19.70 | 9.80 | | | | | | |
| 2 | 57.90 | 36.10 | 24.80 | 21.50 | 11.40 | 7.00 | | | | | | |
| 3 | 46.60 | 39.20 | 33.60 | 32.50 | 26.90 | 15.00 | 12.00 | | | | | |
| 4 | 75.50 | 52.40 | | 37.20 | 24.10 | | 8.79 | | | | | |
| 5 | 72.00 | 60.10 | 47.20 | 38.30 | 32.70 | 12.00 | | | | | | |
| 6 | 50.50 | 50.20 | 38.50 | 32.60 | 30.20 | 13.40 | | 7.73 | | | | |
| 7 | 91.40 | 89.70 | 88.40 | 64.60 | 66.00 | 31.00 | 17.00 | | | | | |
| 8 | 62.00 | 54.20 | | 27.60 | 19.70 | 8.43 | 6.60 | | | | | |
| | 8 | 8 | 8 | 8 | 8 | 7 | 6 | 2 | . 0 | 0 | 0 |) 0 |
| HEAN | 62.54 | 53.48 | | | | | | | | • | • | |
| SD | 16.21 | 16.61 | | | | | | | | | | |
| CV(Z) | 25.93 | 31.06 | | | | | | | | | | |
| MAX | 91.40 | 89.70 | | | | | | | | | | |
| | - 44.40 | 36.10 | | | | | | | | | | |
| ***** | | | , •• | | | 30 | | , , , , | • | | | |

Absent values indicate the concentration was less than the minimum detectable concentration of 6.44 ng/ml

Table 4

Pharmacokinetic Parameters for Individual Subjects

PLASMA

| SUBJ | VOLUME/F: | K01≈ | K10 ³ | TLAG* | AUC = | PEAK CONCº | PEAK TIME? |
|-------|-----------|-------|------------------|-------|----------|------------|------------|
| | L | /HR | /HR | HR | NG.HR/ML | NG/ML | HR |
| | | | | . 55 | 100.74 | F7 0 | 7 5 |
| 1 | 813.98 | 1.36 | 0.088 | 0.89 | 692.34 | 57.2 | 3.5 |
| 2 | 512.19 | 1.32 | 0.192 | 0.44 | 503.01 | 75.4 | 1.5 |
| 3 | 1013.90 | 0.72 | 0.048 | 0.79 | 1022.39 | 52.5 | 3.0 |
| 4 | 693.06 | 1.15 | 0.067 | 0.36 | 1073.80 | 85.6 | 2.5 |
| 5 | 699.09 | Ů.44 | 0.076 | 0.88 | 930.48 | 72.0 | 5.0 |
| 6 | 875.82 | 0.83 | 0.051 | 0.32 | 1114.84 | 62.3 | 3.0 |
| 7 | 416.34 | 0.63 | 0.055 | 0.42 | 2141.96 | 116.0 | 2.0 |
| 8 | 843.59 | 0.58 | 0.081 | 0.30 | 722.71 | 66.9 | 3.8 |
| MEAN | 733.50 | 0.88 | 0.082 | 0.55 | 1025.19 | 73.5 | 3.0 |
| | | | | | | | |
| SD | 196.35 | 0.35 | 0.047 | 0.26 | 498.79 | 20.1 | 1.1 |
| CV(%) | 26.77 | 39.77 | 56.736 | 46.74 | 48.65 | 27.4 | 36.0 |
| MAX | 1013.90 | 1.36 | 0.192 | 0.89 | 2141.96 | 116.0 | 5.0 |
| MIN | 416.34 | 0.45 | 0.048 | 0.30 | 503.01 | 52.5 | 1.5 |

BLOOD

| SUBJ | VOLUME/F L | | | PEAK CONC NG/ML | |
|------|-------------------|------|------|--------------------|---|
| - | 1239.19 782.04 | | | 39.1 60.9 | * |

¹ Volume of distribution divided by fraction available, F

² Absorption rate constant

³ Elimination rate constant

⁴ Time delay before onset of absorption

⁵ Area under the concentration time curve for the equation fit to the plasma concentrations

⁶ Observed peak plasma concentration, as free base

⁷ Time at which peak concentration occurred

Table 5
Recovery of WR6026 in the Unine as Unchanged Drug and Metabolites.

| | Hydroxymethyl- WR6026 (micromoles) | Desethyl WR6026 (micromoles) | WR6026 | Total Recovered | |
|-----------|--|------------------------------------|--------------|-----------------|------|
| | (mid: c) combides/ | Aut C. Ourot 6.22 | Amici Omores | (m) C) Omb) es | , |
| Subj 1 | 24.5 | 0.25 | 0.82 | 25.6 | 17.7 |
| 2 | 14.0 | 0.22 | 0.57 | 14.8 | 10.3 |
| 3 | 17.2 | 0.28 | 1.56 | 19.0 | 13.3 |
| 1 | 7.9 | 0.18 | 0.87 | 9.0 | 6.2 |
| p. | 9.8 | 0.14 | 0.53 | 10.5 | 7.3 |
| 6 | 16.9 | 0.26 | 1.33 | 18.5 | 12.3 |
| 7 | 17.6 | 0.79 | 4.57 | 23.0 | 15.9 |
| 8 | 41.3 | 0.52 | 1.47 | 43.3 | 30.0 |
| | | | | | |
| MEAN | 18.7 | 0.3 | 1.5 | 20.5 | 14 |
| STD. | DEV. 10.5 | 0.2 | 1.3 | 10.8 | 7.5 |

Pach subject received a 60 mg dose of WR6026 dihydrochloride, which is 144.6 micromoles.

Table 6

Derived Pharmacokinetic Parameters for Individual Subjects

PLASMA

| SUBJ | KO1 HL1 | K10 HL≃ | TMAX≅ | CMAX* |
|-------|----------------------|-------------------|----------------|------------------|
| | HR | HR | HR | NG/ML |
| 1 | 0.51 | 7.89 | 3.04 | 50.33 |
| 2 | 0.53 | 3.61 | 2.15 | 69.55 |
| 3 | 0.96 | 14.52 | 4.82 | 40.27 |
| 4 | 0.60 | 10.42 | 2.99 | 59.94 |
| 5 | 1.56 | 9.11 | 5.67 | 49.18 |
| 5 | 0.84 | 13.67 | 3.91 | 47.11 |
| 7 | 1.09 | 12.49 | 4.63 | 94.10 |
| 8 | 1.20 | 8.54 | 4.25 | 42.57 |
| MEAN | 0.91 | 8.435 | 3 .9 3 | 56.63 |
| SD | 0.37 | | 1.15 | 17.86 |
| CV(%) | 40.36 | 35.44 | 29,26 | 31.53 |
| MAX | 1.56 | 14.52 | 5.67 | 94.10 |
| MIN | 0.51 | 3.61 | 2.15 | 40.27 |
| | | BLOOD | | |
| | | BEOOD | | |
| SUBJ | KOJ HL HR | KIO HL HR | TMAX HR | CMAX NG/ML |
| 1 | 0.37 | 10.48 | 2.72 | 35.38 |
| 2 | 0.58 | 4.08 | 2.20 | 45.80 |
| .6- | ration in the second | 1 · · · · · · · · | and a district | -4 (*) * (*) (*) |

i Absorbison halflife

² Elimination halflife

³ Time at which maximum concentration occurs according to fitted equation

⁴ Maximum concentration from fitted equation

⁵ Geometric mear.

Table 7

Comparison of Area Under the Curve by Trapezoidal Rule and Fitted Equation

| SUBJECT | FLUID | AUC° TRAPEZOIDAL RULE NG.HR/ML | AUC** CURVE FIT NG.HR/ML | DIFFERENCE TRAPEZOIDAL RULE MINUS CURVE FIT | CURVE FIT AS % TRAPEZOIDAL RULE |
|---------|--------|--------------------------------------|--------------------------------|---|---------------------------------------|
| 1 | PLASMA | 710.17 | 692.34 | 17.83 | 97.49 |
| 2 | PLASMA | 511.89 | 503.01 | 8.88 | 98.27 |
| 2 3 | PLASMA | 1058.06 | 1022.39 | 35.67 | 96.63 |
| 4 | PLASMA | 1118.29 | 1073.80 | 44.49 | 96.02 |
| 5 | PLASMA | 985.85 | 930.48 | 55. 37 | 94.38 |
| 6 | PLASMA | 1144.01 | 1114.84 | 29.17 | 97.45 |
| 7 | PLASMA | 2178.82 | 2141.96 | 36.86 | 98.31 |
| 8 | PLASMA | 821.63 | 722.71 | 98.92 | 87.96 |
| | MEAN | 1066.09 | 1025.19 | 40.93 | 95.81 |
| | SD | 499.58 | 498.79 | 27.61 | 3.43 |
| | CV(X) | 46.86 | 48.65 | 67.44 | 3.58 |
| | MAX | 2178.82 | 2141.96 | 98.92 | 98.31 |
| | MIN | 511.89 | 503.01 | 8.88 | 87.96 |
| | | | | | |
| 1 | BLOOD | 625.06 | 603.70 | 21.36 | 96.58 |
| 2 | BLOOD | 394.21 | 372.14 | 22.07 | 94.40 |
| | | | | | |

- Calculated using the linear trapezoidal rule for the measured concentrations and adding the area extrapolated to zero drug concentration (obtained by dividing the last measured concentration by the elimination rate constant K10 from the curve fitting).
- The area from time zero to infinity for the equation fit to the observed data. This is the integral of the equation: C(T') = [D*KO1/(V/F)/(KO1-K10)]*[EXP(-K10*T')-EXP(-KO1*T'] where T'= T-TLAG

ጀመቼያው ያለው የሚያስፈው ያለው የሚያስፈው የሚያስፈው

Table 8

Elimination Rate Constant and Plasma Halflife of WR6026
Calculated from the Uninary Excretion of WR6026

| Subject | Elimination Rate /hr | Halflife hrs |
|------------------|-------------------------|-----------------|
| i | .050 | 13.9 |
| 2 | .112 | 6.2 |
| 3 | .042 | 16.5 |
| Ą | . 042 | 16.5 |
| 5 | .048 | 14.4 |
| $\dot{\epsilon}$ | .043 | 16.1 |
| 7 | .065 | 10.7 |
| 8 | .078 | 8.9 |
| | | |
| MEAN | ,060 | 11.61 |
| STD. DEV. | .025 | |

^{*} Geometric Mean

Table 9

A

33

88

X

88

%

X

K

Pharmacokinetic Data for Individual Subjects

| 1 | 3/34H DIT THOUSE STORY | SET CUT 110 | J. J. | 5 | 2 | ď | FERTIDIAE | CALCULATED ANT IN URINE | RECOVERED AMT 4 N HETNE | CALCULATED | Trapezoidal | X | CHRX | TIME of PERK P | PERK CONC |
|-------------------|------------------------------|---------------|--|--------------|----------------|------|-----------|-------------------------|----------------------------|---|---|------------------------|-------------|-----------------|----------------|
| 2000 | | | ֡֝֝֝֡֝֝֡֝֝֝֡֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֡֓֓֓֓֓֓֡֓֓֡ | ¥ | ¥ | ¥ | | δ'n | 97 | | | Ŧ | FG/HL | | NG/HL |
| | Urine | | | | 0.050 | | 0.00028 | 282 | | | | | | 10° | 57.5 |
| ~ | Plasma | -2 | 814 | 1.36 | 0.088 | | | | | 632 | 710 | 3.0 | 50.3 | ري. دي. | 57.2 |
| | Plasmat Urine | ć, | 1043 | 2.30 | 0.057 | | 0.00032 | 271 | | | 177 | <u>ب</u> در | ტ ∾, | ю. М. | 57.2 |
| | Plasna+Urine | -1 | 946 | 1.95 | 0.057 | 0.91 | 0.00031 | 271 | \$10 610 | 요 (20 4 년 | 10 to | op o ru o | 47. ± | ကျော် ကြောင် | () () () |
| ⊷ | Plasnai Urine | c | 833 | 1.86 | 0.055 | • | 0.00031 | E N | | | زرر | 7. W | T. | 0 7 | 2.30 |
| €¥ | Urine | | | | 0.112 | | 0.00044 | 199 | 197 | | 554 | | | 2.1 | 75.4 |
| ĆΙ | Pl asna | Ņ | 512 | 1.32 | 0.192 | | | | 197 | 503 | 512 | 2.2 | 59.6 | 1.5 | 75.4 |
| CI. | Plasnat Urine | Ų. | 989 | 2.16 | 0.132 | 0.45 | 0.00051 | 190 | | 1. | 533 | ე. ა | 50.2 | 1.5 | 75,4 |
| | Plasmat Urine | -1 | 604 | 1.83 | 0.141 | • | 0.00054 | 189 | | 581 | 533 | 2.0 | 66.2 | 1.0 | ۲. ان |
| | Plasmat Urine | 0 | 290 | 1.84 | 0.135 | • | 0.00053 | 191 | | 622 | 99. 1- | 2.0 | 68.2 | 1.5 | 75.4 |
| ίω | Urine | | | | 0.042 | | 0.00045 | 988 | 537 | | 1094 | | | 0.6 | 52.5 |
| Φ | P] asna | ç | 1014 | 0.72 | 0.048 | | | | 533 | 1022 | 1058 | م. ش | 40.3 | 0. % | 52.5 |
| m | Plasnaturine | ~ | 1106 | 1.04 | 0.046 | 0.85 | 0.00050 | 525 | 537 | 696 | 1069 | ٥. | 38.7 | ©: M | 52.5 |
| , r. , | Plasnat Urine | -1 | 1041 | 1.07 | 0.045 | | 0.00049 | 523 | 537 | 1042 | 1069 | 4.0 | ۰. ۲ | ©. ₩ | લું |
| 07 | PlasmatUrine | 0 | 1008 | 1.21 | 0.045 | | 0.00049 | 525 | E S | 1101 | 1075 | တ က | ლ ლ | ©. | 52.5 |
| á. | Urine | | | | 0.042 | | 0.00026 | 267 | 300 | | 1196 | | | 2.5 | 85.6 |
| ዋ | Plasna | Çi | 643 | 1.15 | 0.067 | 0.36 | | | 300 | 1074 | 1118 | 0.5 | 59.9 | න. ව | 85.6 |
| | Plasmat Urine | ņ | 759 | 1.24 | 0.056 | 0.35 | 0.00033 | 583 | 3000 | 1160 | 1144 | 0 0 | က္ မွာ | io o | છે. જું |
| ŗ | Plasnat Urine | -1 | 6 83 | 1.41 | 0.058 | 0.41 | 0.00034 | 286 | 300 | 1252 | 1137 | က လ | ଜ୍ୟ ଜ୍ୟ | นา กับ | 9.5 0 |
| | PlasmatUrino | 0 | 632 | 1,54 | 0.059 | 9.46 | 0.00035 | 287 | 300 | 1317 | 1136 | r- ci | .39 | ις: -> | æ. 8 |
| s) | Urine | | | | 0.048 | | 0.00018 | 208 | 181 | | 1045 | | | 5.0 | 72.0 |
| u? | P] asha | 2 | 669 | 0.4 | 0.076 | 68.0 | | | 181 | 931 | 986 | ر. د. | 3.64 | ت. ت. | 72.0 |
| | Plasmat Urine | -2 | 984 | 0.19 | 0.047 | | 0.00017 | 179 | 191 | 1070 | 1048 | φ. α. | 31.7 | o.v | 72.0 |
| | Plasma+Urine | -1 | 967 | 0.61 | 0.041 | 5 | 0.00015 | 180 | 191 | 1382 | 1072 | ۲.۵ | 46.3 | 0.0 | 72.0 |
| m | Plasmat Urine | 0 | 748 | 69.0 | 0.046 | 11 | 0.00017 | 179 | 181 | 1435 | 1052 | ۳. ن | 54.6 | 0.0 | 72.0 |
| .20 | Urine | | | | 0.043 | | 0.00040 | 4 1- | 456 | | 1172 | | | 3.0 | 62,3 |
| w | Plasma | 7 | 876 | 0.83 | 0.051 | 0.32 | | | 456 | 1115 | 1144 | 6.6 | 47.1 | 3.0 | 62.3 |
| | Plasmat Urine | -5 | 946 | 0.43 | 0.040 | 3 | 0.00039 | 447 | 456 | 1233 | 1186 | 6.1 | 41.0 | 0.0 | 65.3 |
| | Plasmat Urine | . | 919 | G. 83 | 0.038 | 32 | 0.00037 | 440 | 455 | 1428 | 1196 | 4.1 | 46.3 | 0 0 | 62.3 |
| | PlasmatUrine | 0 | 86.2 | 66.0 | 0.039 | 0.43 | 0.00038 | <u>प</u> ए | 436 | 1490 | 1191 | တ က် | e. 08 | ⊕. M | 62.3 |
| ~ | Urine | | | | 0.065 | | 0.00207 | 1371 | 1565 | | 2147 | | | 2.0 | 116.0 |
| <i>~</i> | Plasha | 7 | 416 | 0.63 | 0.055 | 0.4 | • | 1 | 1565 | 2142 | 2179 | છ. च ैं। | 94.1 | æ. | 116.0 |
| | Plasma+Urine | N . | 785 785 | | 0.050 | ₽; | 0.00191 | וכנו | 1965 | 5 () () () () () () () () () (| 2000 | | n | ⊃ (N (| 116.0 |
| ~ r_ | PlasmatUrine PlasmatUrine | . 0 | 429 | 0.78 1.75 | 0.056 0.054 | 0.70 | 0.00175 | 1560 1560 | 1959 1959 | 2135 2135 | 2165 2183 | . v. | 103.4 | 0.0 | 116.0 |
| | | | | | | | | ! | ļ | | 1 | | | • | • |
| ⊙ α | Urino | ç | 0 4 | c c | 0.078 | (F | 0.00067 | ~ इ. १७ | 503 508 | 507 | かん かな で で | ir. | 42.6 | က က က | 5. 6. 6. 6. |
| | DI semastificione | 1 (| 2.0 | 4 | 080 | | 0.00082 | EU5 | 505 503 | 750 | 1 10 1 00 1 00 | <u>.</u> | 41.1 | : 03 : 10 | 9.4 |
| | Plasmat Urine | , | 754 | 0.63 | 0.075 | | 0.00077 | 303 | 503 | 877 | 853 | 4.1 | o. • • • | ω. | P. 69 |
| | Plasmat Urine | O | 706 | 1.17 | 0.074 | | 0.00078 | 504 | 503 | <u>ن</u> 4 | 5,78 | м ,6 | ୍ୟୁଷ୍ଟ | ø. (*) | 66.9 |
| | | | | | | | | | | | | | | | |

Table 10

Pharmacokinetic Parameters for Individual Subjects
Estimated Using Observed Plasma and Urinary Data¹

| SUBJ | VOLUME/F? | KO13 ZHR | K10⁴ /HR | TLAG¤ HR | F*KURINE* /HR | TMAX? HR | CMAXª NG/ML |
|--------|-----------|-------------|-------------|-------------|------------------|-------------|----------------|
| 1 | 1043 | 2.30 | 0.057 | 0.92 | 0.00032 | 2.6 | 43.2 |
| 2 | 686 | 2.16 | 0.132 | 0.45 | 0.00051 | 1.8 | 60.2 |
| 3 | 1106 | 1.04 | 0.046 | 0.85 | 0.00050 | 4.0 | 38.7 |
| 4 | 759 | 1.24 | 0.056 | 0.35 | 0.00033 | 3.0 | 56.3 |
| 5 | 984 | 0.19 | 0.047 | 0.00 | 0.00017 | 9.8 | 31.7 |
| 6 | 946 | 0.43 | 0.040 | 0.04 | 0.00039 | 6.1 | 41.0 |
| 7 | 392 | 0.50 | 0.050 | 0.40 | 0.00191 | 5.2 | 97.3 |
| 8 | 810 | 0.41 | 0.080 | 0.19 | 0.00082 | 5.1 | 41.1 |
| MEAN | 840 | 1.03 | 0.045 | 0.40 | 0.00062 | 4.7 | 51.2 |
| as | 234 | 0.82 | 0.030 | 0.34 | 0.00056 | 2.5 | 20.8 |
| CV (%) | 28 | 78.97 | 45.939 | 85.13 | 89.77831 | 53.9 | 40.7 |
| MAX | 1106 | 2.30 | 0.132 | 0.92 | 0.00191 | 9.8 | 97.3 |
| MIN | 382 | 0.19 | 0.040 | 0.00 | 0.00017 | 1.8 | 31.7 |

- 1 Observed data weighted to the reciprocal of the square of the number
- 2 Volume of distribution divided by fraction available, F
- 3 Absorption rate constant
- 4 Elimination rate constant
- 5 Time delay before onset of absorption
- 5 Area under the concentration time curve for the equation fit to the plasma concentrations
- 6 Rate constant of elimination of unchanged drug into the urine times the fraction available
- 7 Time at which maximum concentration occurs according to fitted equation
- 8 Maximum concentration from fitted equation

Figure 1

Chemical Structure of WR 6026

$$CH_3^0$$
 CH₃ · 2HC1
 $NH(CH_2)_6N(C_2H_5)_2$

Figure 2







Figure 3


PLASMA [WR6026]: SUBJECTS 4, 5, AND 6



-u-5 -x-

Figure 4

PLASMA [WR6026]; SUBJECTS 7 AND 8



-0- ? -0- ?

Figure 5





Figure 6

BLOOD [WR6026]: SUBJECTS 1 AND 2



-0 1 -0 2


Figure 7

Z.


8

SS SSS 653

No.










Figure 10

K

8

×

X


O.

8

Š.

N.

38 38



hass broom broom brooms more and brooms of brooms brooms brooms brooms brooms brooms

Figure 11

X

8

8

Z

*3*2

8

Š

8

K




Figure 12

8

*

X

8

*

X,

3

Ş

8

ř

8

Ç




Figure 13

% %

33

7.

8

8

Ů,

3

T.

N

Ø

8

75

30333333 115-53-5354

asserted largeries.




Figure 14

Z

3

53

(K)

Í

K


不完


· •

22.2

E


X






X

X





8

77.77

8

8

2527 1825

7

833

N.

X

3


8


ÇĻ.

X

15

Figure 17






EXX

K.

Subject 4

Figure 19



Ä

*

N.

X

X

S

Y.

28

X

X

8


×.


X

Ø.

8

Ç





APPENDIX A Study Flow Chart

E

| | <u>DAY</u> | HOUR | APPROXIMATE TIME | PROCEDURE |
|-----------------|------------|------|------------------|---|
| 333 83 3 | 1 | -46 | 10:00 a.m. | Subject admitted to unit. Sign consent form. Blood tests: CBC with differential SMA 6 SMA 12 LDH, CK Urinalysis |
| \$\$ \$\$ | 2 | - 24 | 8:00 a.m. | Subject begins 12-hour urine collection. |
| ** | | -12 | 8:00 p.m. | Subject concludes 12-hour urine collection, begins another 12-hour urine collection. |
| X | | - 8 | 12:00 a.m. | NPO |
| 3 | 3 | -1 | 7:00 a.m. | Insert heparin lock for blood sampling. |
| 29 | | -0.5 | 7:30 a.m. | Electrocardiogram |
| | | 0 | 8:00 a.m. | Subject concludes 12-hour urine collection, begins another 12-hour urine collection. Blood tests: CBC with differential SMA6 SMA12 LDH,CK Methemoglobin level Elood for WR 6026 level. Subject takes |
| | | | | four 15 mg capsules of WR 6026. |
| | | 0.25 | 8:15 a.m. | Blood for WR 6026 level |
| 33 | | 0.5 | 8;30 a.m. | Blood for WR 6026 level |
| Y. | | 0.75 | 8:45 a.m. | Blood for WR 6026 level |
| | | 1.0 | 9:00 a.m. | Blood for WR 6026 level |
| * | | 1.25 | 9:15 a.m. | Blood for WR 6026 level |
| 数 | | 1.5 | 9:30 a.m. | Blood for WR 6026 level |

ማመመመመው የመድድ የተመሰቀተው የተመሰቀት የ

| X | DAY | <u>HOUR</u> | APPROXIMATE TIME | PROCEDURE |
|------------------|----------|-------------|------------------|--|
| ax — | 3 (cont) | 2.0 | 10:00 a.m. | Blood for WR 6026 level |
| ·/· | | 2.5 | 10:30 a.m. | Blood for WR 6026 level |
| NS. | | 3.0 | 11:00 a.m. | Blood for WR 6026 level |
| X | | 3.5 | 11:30 a.m. | Blood for WR 6026 level |
| % | | 4.0 | 12:00 p.m. | Blood for WR 6026 level. Subject may resume eating. Electrocardiogram |
| *** | | 5.0 | 1:00 p.m. | Blood for WR 6026 level |
| | | 6.0 | 2:00 p.m. | Blood for WR 6026 level |
| \$ \$ | | 8.0 | 4:00 p.m. | Blood for WR 6026 level |
| NA. | | 10.0 | 6:00 p.m. | Blood for WR 6026 level |
| r F | | 12.0 | 8:00 p.m. | Blood for WR 6026 level. Subject concludes 12-hour urine collection, begin another 12- hour urine collection. |
| Ĭ | 4 | 24.0 | 8:00 a.m. | Blood for WR 6026 level. Subject concludes 12-hour urine collection, begins another 12- |
| 25 | | | | hour urine collection. Blood tests: CBC with differential |
| W. | | | | Methemoglobin level SMA 6 SMA 12 |
| 88 | | | | Lipid profile LDH, CK |
| Ä | | | | Electrocardiogram |
| 7 33 | | 36.0 | 8:00 р.ш. | Blood for WR 6026 level. Subject concludes 12-hour urine collection, begins another 12- |
| | | | | hour urine collection. |

| Q | | | | |
|----------------|-----|------|------------------|---|
| 222 | DAY | HOUR | APPROXIMATE TIME | PROCEDURE |
| Ď | 5 | 48.0 | 8:00 a.m. | Blood for WR 6026 level. Subject concludes 12-hour urine collection, begins another 12- hour urine collection. |
| 8 | | | | Blood tests: CBC with differential Methemoglobin level SMA 6 |
| 3 | | | | SMA 12 Lipid profile |
| \$3. \$4. | | | | LDH, CK Electrocardiogram |
| ~ \$8 | | 60.0 | 8:00 p.m. | Blood for WR 6026 level. Subject concludes 12-hour urine collection, begins another 12- hour urine collection. |
| 82 84 8* | 6 | 72.0 | 8:00 a.m. | Blood for WR 6026 level. Subject concludes 12-hour urine collection, begins another 12- hour urine collection. |
| S. S. | | 84.0 | 8:00 p.m. | Blood for WR 6026 level. |
| | | | | Subject concludes 12-hour urine collection, begins another 12-hour urine collection. |
| 23 | 7 | 96.0 | 8:00 a.m. | Blood for WR 6026 level. Subject concludes 12-hour urine collection, begins another 12- |
| | | | | hour urine collection. Blood tests: |
| 3 | | | | CBC with differential Methemoglobin level SMA 6 |
| | | | | SMA 12 Lipid profile |
| | | | | LDH, CK Electrocardiogram |
| | | 98.0 | 10:00 a.m. | Urinalysis Discharge from unit. |

| Form C (Revised | 9/85 |
|-----------------|------|
| J.H.U.M.S. | 1 |

APPENDIX B

CLINICAL INVESTIGATION CONSENT FORM The Johns Hopkins Medical Institutions

Title of Research Project: Single-Dose Absorption and Pharmacokinetics of WR 6026 Hydrochloride in Healthy Subjects

Patient I.D. Plate

Explanation of Research Project to Subject:

You are invited to participate in a study of a drug called WR 6026. WR 6026 is not a licensed drug and therefore is considered investigational. This is a drug used to treat an infection called leishmaniasis which usually occurs in tropical countries and is caused by a parasite. WR 6026 has shown promise as an effective drug to treat this infection. The purpose of this project is to see how healthy volunteers absorb the drug after it is given by mouth, how long the drug stays in the bloodstream, and what byproducts of drug metabolism are eliminated in the urine.

If you agree to join the study you will be hospitalized for six days. You will collect all urine for analysis starting on the second hospital day. On the third hospital day you will take twelve 5 mg capsules of WR 6026 for a total of 60 mg. Following drug administration repeated blood samples will be taken by means of a "heparin lock," a device like an intravenous catheter that stays in your vein and allows blood to be removed without sticking a new needle through the skin each time. A total of 24 blood samples, each less than one tablespoon, will be removed for analysis over the four days after the drug is given. In combination with blood specimens taken before and after the drug is given to monitor its safety, the total amount of blood to be removed for this project is less than one pint. This is less than the amount routinely donated at a blood bank.

We believe that the risks involved with this study are small. When given to animals in large doses for long periods, WR 6026 has caused changes in the kidney, gallbladder, liver, spleen, lungs and heart. In this project however, a much smaller dose for body weight will be given and it is given only once. Healthy human subjects in the 1940's were given WR 6026 for two weeks at a daily dose one half as much as in this project, and they developed only minimal changes in blood tests and EKG but no symptoms. More recently, healthy volunteers have taken single doses of WR 6026, including two who took the same amount you will take in this project, and no symptoms or laboratory abnormalities were seen. The chance of your clinical laboratory tests being abnormal is small. However, if your clinical laboratory tests on Day 7 are abnormal, you will be invited to return for follow-up clinical testing on Day 13 and weekly thereafter until the tests become normal or an alternative explanation is determined. A risk of having a heparin lock for blood withdrawal is the possible discomfort of swelling, soreness and bruising.

Benefits to you for participation in this study are primarily financial, but another potential asset is the comprehensive medical evaluation which accompanies this project, the records of which will be available in the future. If WR 6026 is found to be well absorbed and safe, then it can be studied further to see if it is a better form of therapy for the parasite infection than current drugs.

You are under no obligation to participate in this project. Should you decide not to participate or should you decide to withdraw during the course of the project, your future care at Hopkins will not be affected. You will be paid by check for the proportion of the study which you have completed at the time of withdrawal. Successful completion of the entire study will pay \$250. You will be paid by check when you leave the hospital.

Army inspectors may look at the relevant part of your medical record as part of their job to review this study. You are authorized all medical care for injury or disease which is a proximate result of your participation in this research. The medical treatment provided might include, if necessary, laboratory tests, X-rays and other procedures used in diagnosis and treatment. No other compensation for injury is offered.

Appendix C: Measured Concentrations of WR6026

803

22.2

| atiett. Tal | (TI LITT) | SCHEDULED TIME | ACTUAL TI | ME NGpML |
|-------------|---------------------|---|---------------------|---|
| DUDUB | FLOID | hr post dose | | se base |
| 1 | E-[_ | 0.00 | -0.23 | ± 6 ± 6 ± 6 ± 6 ± 6 ± 6 ± 6 ± 6 ± 6 ± 6 |
| 1. | F.L. | 0.25 | 0.25 | * |
| 1 | PL | 0.50 | 0.50 | · · · · · · · · · · · · · · · · · · · |
| 1 | LD (| 0.75 | 0.75 | * |
| 1 | Fi. | 1.00 | 1.00 | 8.23 |
| 1 | F-L | 1.25 | 1.25 | 29.8 |
| 1 | 1-1-L | 1.50 | 1.50 | 26.8 |
| 1 | PL | 2.00 | 2.00 | 48.7 |
| 1 | P1. | 2.50 | 2.50 | 51.0 |
| 1 | F'L | 3.00 | 3.00 | 53.1 |
| 1 | PL. | 3.50 | 3.50 | 57.2 |
| j | PL | 4.00 | 4.00 | 49.2 |
| 1 | PL | 5.00 | 5.00 | 44.4 |
| 1 | Pi_ | 6.00 | 6.00 | 45.9 |
| 1 | PL | 8.00 | 8.00 | 31.4 |
| 1 | FL. | 10.00 | 10.00 | 32.3 |
| 1 | F'L | 12.00 | 12.00 | 19.7 |
| 1 | PL | 24.00 | 24.00 | · /• / 두• 음 |
| 1 | PL. | 36.00 | 36.00 | / . |
| 1 | F/L | 48.00 | 48.00 | ₩ |
| L. | PL. | 40.00 60.00 | 60.00 | |
| ì | F-L | 72.00 | 72.00 | · · · · · · · · · · · · · · · · · · · |
| 1 | P1_ | 84.00 | 84.00 | * |
| 1 | : 12 PL | 96.00 | 96.00 | |
| , | i 1 | 70400 | A Contract Contract | , |
| | | | | |
| | 6.24 _{2.1} | (\(\frac{1}{2}\) (\(\frac{1}{2}\) (\(\frac{1}{2}\)) | -0.28 | * |
| Ξ' | FL | 0.25 | 0.25 | * |
| 4.5 4.5 | F'L | 0.50 | 0.50 | 7.26 |
| <u> </u> | FI | 0.75 | 0.75 | 30.5 |
| 2) | FL | 1.00 | 1.00 | 50.9 |
| - | 1." | 1.25 | 1.25 | 60.3 |
| 2 | FL | 1.50 | 1.50 | 75.4 |
| 2 | PL. | 2.00 | 2.00 | 56.3 |
| 7 | F·L_ | 2.50 | 2.50 | 67.0 |
| 2 | ۶L. | 3. 00 | 3.00 | 59.4 |
| 2 | PL. | 3.50 | 3.50 | 69.5 |
| - | F'L. | 4.00 | 4.00 | 59.8 |
| 11 11 11 11 | FL | 5.00 | 5.00 | 57.9 |
| 2 | F1 | 6.00 | 6.00 | 36.1 |
| - | PL. | 8.00 | 8.00 | 24.8 |
| Ω | F·L | 10.00 | 10.00 | 21.5 |
| 1 | FL | 12.00 | 12.00 | 11.4 |
| ?? | E-1 | 24.00 | 24.00 | * |
| | 174 | 36.00 | 3 5. 00 | * |
| *** | <u>: 1, 1, 1</u> | 48.00 | 48.00 | • |
| 2 | Fil | 60.00 | 58.22 | * |
| 22 | FfL, | 72.00 | 72.00 | * |

reconstration below the detectability limit. 6.44 mg/ml

Appendix C: Measured Concentrations of WR6026

| 2 2 | PL PL | 84.00 96.00 | 84.00 96.00 | * |
|---------------------------------|----------------------------------|--|--|---|
| аксакананананананана | | 0.00 0.25 0.50 0.75 1.00 1.25 1.50 2.50 3.00 3.50 4.00 5.00 10.00 12.00 24.00 24.00 48.00 48.00 48.00 72.00 | -0.47 0.25 0.50 0.75 1.00 1.25 1.50 2.00 2.50 3.00 3.50 4.00 5.00 8.00 10.02 12.05 24.00 36.03 48.17 61.37 72.05 | * * * * * 7.47 14.7 14.9 46.2.6.3 46.2.6.3 46.2.6.3 12.6.3 |
| 3 3 | PL PL | 84.00 96.00 | 84.12 96.17 | * |
| 4 4 4 4 4 4 4 | PL PL PL PL PL PL | 0.00 0.25 0.50 0.75 1.00 1.25 1.50 2.00 | -0.17 0.25 0.50 0.75 1.00 1.25 1.50 2.00 | * 12.9 18.0 37.8 51.2 59.6 82.6 85.6 |
| 4 4 4 4 4 4 4 | PL PL PL PL PL | 3.00 3.50 4.00 5.00 6.00 8.00 10.00 12.00 | 3.00 3.50 4.00 5.00 6.00 8.05 10.00 12.00 24.00 | 68.6 62.8 75.5 52.4 36.8 37.2 24.1 |
| 4 4 | PL PL | 36.00 48.00 | 36.00 48.00 | 8.79 * |

Concentration below the detectability limit, 6.44 ng/ml

Appendix C: Measured Concentrations of WR6026

| 4 4 4 | PL PL | 50.00 72.00 84.00 | 59.43 72.03 84.08 | * * |
|-------------|--------------|-------------------------|-------------------------|--------------|
| 4 | FL. | 96.00 | 96.00 | * |
| 5 5 | PL PL | 0.00 0.25 | -0.17 | * |
| 5 | PL | 0.50 | 0.25 0.50 | * |
| 5 | FL. | 0.75 | 0.75 | ≅ |
| 5 | FL | 1.00 | 1.00 | * |
| 5 | F.L. | 1.25 | 1.25 | 10.6 |
| 5 | PL. | 1.50 | 1.50 | 19.4 |
| 5 | FL | 2.00 | 2.00 | 24.9 |
| 5 | FL | 2.50 | 2.50 | 23.2 |
| 5 | PL | 3.00 | 3.03 | 48.9 |
| 5 | {F'}. | 3.5 0 | 3.50 | 53.3 |
| 5 | FL. | 4.00 | 4.00 | 63.4 |
| 5 | PL | 5.00 | 5.00 | 72.0 |
| 5 | F'L | 6.00 | 6.00 | 60.1 |
| 5 | FL | 8.00 | 8.05 | 47.2 |
| 5 | F'L | 10.00 | 10.02 | 38.3 |
| 5 | FL | 12.00 | 12.00 | 32.7 |
| 5 | FL | 24.00 | 24.00 | 12.0 |
| 5 | PL. | 36.00 | 34.00 | 7.67 |
| 5 | FL | 48.00 | 48.00 | * |
| 5 5 | PL G | 50.00 | 61.62 | * |
| 5 | Fil. Fil. | 72.00 84. 00 | 72.08 | * |
| 5 | P-[_ | 96.00 | 84.45 96.00 | * |
| J | I L. | 7 D • CIO | 70.00 | * |
| 6 | PL PL | 0.00 | ~0.08 | * |
| 5 6 | FL. | 0.25 0.50 | 0.25 | * |
| 6 | FL | 0.75 | 0.50 0.75 | 8.58 13.7 |
| ద | FL | 1.00 | 1.00 | 24.0 |
| 6 | F.L | 1.25 | 1.25 | 29.4 |
| 6 | PL | 1.50 | 1.50 | 31.6 |
| 6 | FL | 2.00 | 2.00 | 49.0 |
| 5 | ₽L_ | 2.50 | 2.50 | 42.9 |
| 6 | PL | 3.00 | 3.00 | 62.3 |
| က် | PL | 3.50 | 3.50°° | 54.5 |
| 6 | F1 | 4.00 | 4.00 | 60.4 |
| Ó | FL | 5 00 | 5.00 | 50.5 |
| 6 | PL. | 6.00 | 6.00 | 50.2 |
| 6 | PL | 8.00 | 8. 00 | 38.5 |
| 6 | PL | 10.00 | 10.00 | 32.6 |
| 5 | PL | 12.00 | 12.00 | 30.2 |
| 6 | Ft. | 24.00 | 24.08 | 13.4 |

Concentration below the detectability limit, 6.44 ng/ml

| | | | · | | |
|---|----------|-----------|----------------|-----------------|-------------------|
| | App | pendix | C: Measured | Concentrations | us WR 6026 |
| | Ć | FL | 36.00 | 36.00 | 10.4 |
| | 6 | PL | 48.00 | 48.00 | 7.73 |
| | გ გ | PL Pl | 60.00 72.00 | 60.05 72.03 | * * |
| | చ | PL. | 84.00 | 84.00 | * |
| | చ | PL | 96.00 | 96.00 | * |
| | | | | | |
| | 7 | Fi_ | 0.00 | -0.53 | }. |
| | 7 | PL | 0.25 | 0.25 | * |
| | 7 | FL | 0.50 | 0.50 | 7.17 |
| | フ フ | PL. | 0.75 | 0.75 | 14.4 35.9 |
| | 7 | PL PL | 1.00 1.25 | 1.00 1.25 | 33.7 73.2 |
| | 7 | FL. | 1.50 | 1.50 | 85.7 |
| | 7 | F'L | 2.00 | 2.00 | 116.0 |
| | 7 | FL | 2.50 | 2.50 | 102.0 |
| | 7 | FL | 3.00 | 3.00 | 94.1 |
| | 7 | PL | 3.50 | 3.50 | 94.9 |
| | 7 7 | FL | 4.00 5.00 | 4.00 5.00 | 102.0 91.4 |
| | 7 | FL. | 5.00 | 6.00 | 89.7 |
| | 7 | Fi_ | 8.00 | 8.00 | 88.4 |
| | 7 | PL | 10.00 | 10.00 | 64.6 |
| | 7 | PL | 12.00 | 12.00 | 66.0 |
| | 7 | F.F | 24.00 | 24.00 | 31.0 |
| | 7 7 | PL. | 36.00 | 36.00 | 17.0 |
| | 7 | F'L PL | 48.00 60.00 | 48.05 59.30 | 11.3 * |
| | 7 | PI. | 72.00 | 72.08 | * |
| | 7 | FL | 84.00 | 84.08 | * |
| | 7 | ₽D() | 96.00 | 96.05 | * |
| | a | F., | 0.00 | ه مح | v |
| | 8 8 | PL Pl | 0.00 0.25 | -0.25 0.25 | * |
| | ອ 8 | PL | 0.50 | 0.50 | 7.63 |
| | 8 | PL | 0.75 | 0.75 | 13.2 |
| - | 8 | F'L | 1.00 | 1.00 | 14.2 |
| | 8 | FL | 1.25 | 1.25 | 16.3 |
| | 8 | PL Pl | 1.50 | 1.50 | 45.3 45.3 |
| | 8 8 | PL PL | 2.00 2.50 | 2.00 2.50 | 65.2 60.5 |
| | 8 | FL. | 3.00 | 3.00 | 64.4 |
| | ខ | FL. | 3.50 | 3.50 | 47.3 |
| | 8 | PL. | 4.00 | 3.83 | 66.9 |
| | 8 | PL | 5.00 | 5.00 | 62.0 |
| | 8 | FIL. | 6. 00 | 6.00 | 54.2 |
| | 8 | PI. | 8.00 | 8.00 | 30.8 |
| | 8 | FL. | 10.00 | 10.02 | 27.5 |
| - | - Cosa | antrat | io, balow th | e detectability | limit 4 A |
| | s court. | with att | LESSO DELOW CI | | AAMACA ORT |

Concentration below the detectability limit, 6.44 ng/ml

Appendix C: Measured Concentrations of WR6026

| 0 0 0 0 0 0 0 | PL PL PL PL PL PL | 12.00 24.00 36.00 48.00 60.00 72.00 84.00 96.00 | 12.02 24.00 36.00 48.12 59.42 72.00 84.10 96.00 | 19.7 8.43 6.60 * * * * |
|---|--|--|---|--|
| 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | BLDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDD | 0.00 0.25 0.50 0.75 1.00 1.25 1.50 2.00 3.50 4.00 5.00 4.00 10.00 12.00 24.00 36.00 48.00 48.00 48.00 84.00 | -0.28 0.25 0.50 0.75 1.00 1.25 1.50 2.00 2.50 3.50 4.00 5.00 6.00 10.00 12.00 24.00 36.00 48.00 58.22 72.00 84.00 | * 6.69 6.89 6.89 7.7 21.7 234.7 235.1 355.3 355.2 21.0 355.2 |
| | BLD BLD BLD BLD BLD BLD BLD BLD BLD BLD | 0.00 0.25 0.50 0.75 1.00 1.25 1.50 2.00 2.50 3.00 3.50 4.00 5.00 | -0.28 0.25 0.50 0.75 1.00 1.25 1.50 2.00 2.50 3.50 4.00 5.00 6.00 | * 14.8 19.2 38.9 44.7 60.9 57.7 42.2 37.0 35.7 40.1 |

Concentration below the detectability limit, 6.44 ng/ml

Appendix C: Measured Concentrations of WR6026

| 2 | BLD | a.00 | 8.00 | 25.0 |
|--------------|------|-------|-------|--------------|
| \mathbb{Z} | BLD | 10.00 | 10.00 | 12.6 |
| 2 | BLD | 12.00 | 12.00 | 10.2 |
| 2 | BLD | 24.00 | 24.00 | * |
| 2 | BL.D | 36.00 | 36.00 | * |
| 2 | BLD | 48.00 | 48.00 | * |
| 2 | ELD | 60.00 | 58.22 | * |
| 2 | BLD | 72.0Q | 72.00 | * |
| 2 | FLD | 84.00 | 84.00 | * |
| | חום | 94 AA | 94.00 | <u>.</u> |

Appendix D

Table 1*

Precision and Accuracy Data for Analysis of WR 6026,
Desethyl WR 6026 and 4-Hydroxymethyl WR 6026 in Human Urine^a

| Amount Added ^b (ng/ml) | Amount Measured ^b (mean ± SD) (ng/ml) | Coefficient of Variation (%) | N |
|-----------------------------------|--|------------------------------------|-------|
| | WR 6026 | | |
| 17 | 18 ± 4 | 27 | 6 |
| 83 | 84 ± 6 | 7 | 10 |
| | Desethyl WR 6 | 026 | |
| 16 | 15 ± 3 | 20 | 6 |
| 79 | 83 ± 6 | . 7 | 8 |
| | 4-Hydroxymethyl | WR 6026 | |
| 15 | 14 ± 2 | 14 | 8 |
| 75 | 75 ± 8 | 8 | 10 |

aData represents a compilation of N separate experiments with 3 replicates of each sample determined for each experiment. A standard curve of 10-500 ng/ml of each compound bracketed spiked unknowns.

unknowns.

bSamples were spiked 20 or 100 ng/ml of the salts of each compound.

Data shown is corrected for presence of various salt forms of the compounds.

Z

^{*} reproduct 1 from reference #11.

TABLE 2*
4-HYDROXYMETHYL WR6026

| | 4-HYDROXYMETHYL WROOZO | | | | | | | |
|--|------------------------|------------|--------------|-------------|-------------|--------------|-------------|--------------|
| | | N | ANOGR | AMS/ML | URINE | | | |
| SAMPLE | PT 1 | PT 2 | PT 3 | PT 4 | PT 5 | PT 6 | PT 7 | PT 8 |
| | | | | | | | | |
| -12-0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0-12 | 2276 | 1700 | 789 | 1054 | 651 | 958 | 1427 | 2657 |
| 12-24 24-36 | 2316 882 | 549 636 | 585 883 | 1406 967 | 1043 771 | 1189 402 | 1279 542 | 2637 1324 |
| 36-48 | 1096 | 290 | 346 | 335 | 517 | 665 | 712 | 956 |
| 48-60 | 336 | 257 | 298 | 490 | 291 | 521 | 425 | 406 |
| 60-72 | 579 | 110 | 222 | 374 | 260 | 237 | 168 | 304 |
| 72-84 | 505 | 146 | 382 | 193 | 253 | 266 | 311 | 357 |
| 84-96 | 336 | 118 | 108 | 121 | 144 | 127 | 143 | 268 |
| | | | | | | | | |
| | | | | VOLUME | (ML) | | | |
| SAMPLE | PT 1 | PT 2 | PT 3 | PT 4 | PT 5 | PT 6 | PT 7 | PT 8 |
| 12.0 | | | | ===== | | 22222 | | ===== |
| -12-0 0-12 | 680 820 | 590 960 | 1045 | 735 | 815 | 1550 | 1500 | 1920 |
| 12-24 | 1050 | 2020 | 1390 1855 | 510 595 | 485 1050 | 1230 1345 | 960 1300 | 1135 1525 |
| 24-36 | 1835 | 1330 | 1328 | 432 | 802 | 1960 | 1605 | 2590 |
| 36-48 | 830 | 1630 | 3130 | 745 | 1160 | 1480 | 1505 | 1615 |
| 48-60 | 1750 | 1426 | 1695 | 685 | 1000 | 950 | 1095 | 2600 |
| 60-72 | 1315 | 1860 | 1805 | 490 | 1070 | 1635 | 1195 | 1890 |
| 72-84 | 715 | 1410 | 1650 | 1050 | 750 | 1600 | 1565 | 1860 |
| 84-96 | 790 | 1720 | 1720 | 745 | 1025 | 1570 | 1460 | 1970 |
| | • | | | | | | | |
| | | | | | | ICROMO | | |
| SAMPLE | PT 1 | PT 2 | PT 3 | PT 4 | PT 5 | PT 6 | PT 7 | PT 8 |
| -12-0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 0-12 | 5.2 | 4.5 | 3.1 | 1.5 | 0.9 | 3.3 | 3.8 | 8.4 |
| 12-24 | 6.8 | 3.1 | 3.0 | 2.3 | 3.0 | 4.5 | 4.6 | 11.2 |
| 24-36 | 4.5 | 2.4 | 3.3 | 1.2 | 1.7 | 2.2 | 2.4 | 9.6 |
| 36-48 | 2.5 | 1.3 | 3.0 | 0.7 | 1.7 | 2.7 | 3.0 | 4.3 |
| 48-60 | 1.6 | 1.0 | 1.4 | 0.9 | 0.8 | 1.4 | 1.3 | 2.9 |
| 60-72 | 2.1 | 0.6 | 1.1 | 0.5 | 0.8 | 1.1 | 0.6 | 1.6 |
| 72-84 | 1.0 | 0.6 | 1.8 | 0.6 | 0.5 | 1.2 | 1.4 | 1.9 1.5 |
| 84-96 | 0.7 | 0.6 | 0.5 | 0.3 | 0.4 | 0.6 | 0.6 | 1.5 |
| mom a r | 74 5 | 14 0 | 17 2 | 7 9 | 9.8 | 16.9 | 17.6 | 41.3 |
| LOTAL | 27.5 | 14.0 | 21.2 | ••• | J. U | 20.5 | 2 | |
| | | | | | | | | |
| | | RATE | OF EXC | RETION | (NANO | MOLES/ | HOUR) | |
| SAMPLE | PT 1 | PT 2 | PT 3 | PT 4 | PT 5 | PT 6 | PT 7 | PT 8 |
| | | | | | | | | |
| -12-0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | | 067 6 | 252 0 | 104 2 | 254 2 | 271 2 | 206 0 | 700.1 |
| 12-24 | 275 4 | 105 3 | 272.0 | 134.Z | 147 K | 182 0 | 202.2 | 933.5 |
| 21-36 76-40 | 3/3.b | 100 7 | 212.2 | 57.0 | 130.3 | 228.5 | 248.8 | 358.5 |
| , 30-40 48-60 | 136 7 | 85.0 | 117.2 | 77.9 | 67.5 | 114.8 | 107.9 | 245.0 |
| 60-72 | 176.6 | 47.7 | 93.1 | 42.5 | 64.6 | 89.9 | 46.5 | 133.4 |
| 12-24 24-36 , 36-48 48-60 60-72 72-84 | 83.9 | 47.9 | 146.2 | 47.2 | 44.0 | 98.6 | 112.9 | 154.3 |
| 84-96 | 61.7 | 47.3 | 43.0 | 21.0 | 34.4 | 46.4 | 48.6 | 122.4 |

^{*} reproduced from reference #11.

Appendix D

TABLE 3* DESETHYL WR6026

| | | • | | | • | | | |
|---------|--------|---------|------------|--------|--------|---------|-------------|----------------|
| | | ì | NANOGRA | MS/ML | URINE | | | |
| SAMPLE | PT 1 | PT 2 | | PT 4 | PT 5 | PT 6 | PT 7 | PT 8 |
| 2222222 | | | | | | | | |
| -12-0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0-12 | 42 | 48 | 20 | 54 | 17 | 23 | 90 | 48 |
| 12-24 | 25 | 11 | 14 | 12 | 24 | 24 | 90 | 51 |
| 24-36 | 10 | 8 | 16 | 24 | 13 | 7 | 2 | 14 |
| 36-48 | 10 | Ö | 8 | 10 | 4 | 'n | 19 | 11 |
| 48-60 | 0 | Ö | Õ | 8 | 0 | 8 | 17 | 0 |
| 60-72 | 0 | 0 | 0 | 3 | Ö | Ö | 8 | 0 |
| 72-84 | 0 | 0 | 0 | 4 | Ö | o | 12 | 0 |
| 84-96 | ű | 0 | 0 | 0 | Ö | 0 | 2 | 0 |
| 04-30 | U | U | U | U | U | U | 2 | U |
| | | | URINE ' | OT UME | /MT \ | | | |
| 03W017 | Dm 1 | | | PT 4 | PT 5 | PT 6 | OM 7 | PT 8 |
| SAMPLE | PT 1 | PT 2 | | | | | PT 7 | ====== =1 0 |
| | | | | | | | | |
| -12-0 | 680 | 590 | 1045 | 735 | 815 | 1550 | 1500 | 1920 |
| 0-12 | 820 | 960 | 1390 | 510 | 485 | 1230 | 960 | 1135 |
| 12-24 | 1050 | 2020 | 1855 | 595 | 1050 | 1345 | 1300 | 1525 |
| 24-36 | 1835 | 1330 | 1328 | 432 | 802 | 1960 | 1605 | 2590 |
| 36-48 | 830 | 1630 | 3130 | 745 | 1160 | 1480 | 1505 | 1615 |
| 48-60 | 1750 | 1426 | 1695 | 685 | 1000 | 950 | 1095 | 2600 |
| 60-72 | 1315 | 1860 | 1805 | 490 | 1070 | 1635 | 1195 | 1890 |
| 72-84 | 715 | 1410 | 1650 | 1050 | 750 | 1600 | 1565 | 1860 |
| 84-96 | 790 | 1720 | 1720 | 745 | 1025 | 1570 | 1460 | 1970 |
| | | | | | | | | |
| | | TOTAL | EXCRET | ED/SAM | PLE (M | IICROMO | LES) | |
| SAMPLE | PT 1 | PT 2 | PT 3 | PT 4 | PT 5 | PT 6 | PT 7 | PT 8 |
| 2222255 | :===== | ===:=== | ===== | ====== | 22222 | ====== | ===== | *** |
| -12-0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| C-12 | 0.09 | 0.13 | 0.08 | 0.08 | 0.02 | 0.08 | 0.24 | 0.15 |
| 12-24 | 0.07 | 0.06 | 0.07 | 0.02 | 0.07 | 0.09 | 0.33 | 0.22 |
| 24-36 | 0.05 | 0.03 | 0.06 | 0.03 | 0.03 | 0.04 | 0.01 | 0.10 |
| 36-48 | 0.02 | 0.00 | 0.07 | 0.02 | 0.01 | 0.03 | 0.08 | 0.05 |
| 48-60 | 0.00 | 0.00 | 0.00 | 0.02 | 0.00 | 0.02 | 0.05 | 0.00 |
| 60-72 | 0.00 | 0.00 | 0.00 | | 0.00 | 0.00 | 0.03 | 0.00 |
| 72-84 | 0.00 | 0.00 | 0.00 | 0.01 | 0.00 | 0.00 | 0.05 | 0.00 |
| 84-96 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.01 | 0.00 |
| • | | | | | | | | |
| TOTAL | 0.25 | 0.22 | 0.28 | 0.18 | 0.14 | 0.26 | 0.79 | 0.52 |
| | 7.20 | | •••• | *** | | | | |
| | | | | | | | | |
| | | RATE | OF EXCE | RETION | (NANO | MOLES/I | (OUR | |
| SAMPLE | DT 1 | | | | | | | PT 8 |
| 2222222 | | | | | | | | |
| | | | 0.0 | | | | | |
| 0-12 | 7 0 | 10.0 | ر د د | £ 3 | 1 0 | £ £ | 20.0 | 12.6 |
| 12-24 | 1.3 | XU. / | 6.3 E n | 1 6 | E 0 | 7 4 | 27.0 | 18.0 |
| 12-24 | 0.1 | 2.3 | 0.U | 7 4 | 2.2 | 2.3 | 21.3 A E | 8.7 |
| 24-36 | 4.3 | 2.3 | 5.0 | 4.4 | 4.4 | 3.3 | 0.0 | 4 1 |
| 36-48 | 2.0 | 0.0 | 5.5 | 1.8 | 1.2 | 2.5 | 6.6 | 4.7 |
| | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 | 1.7 | 4.4 | 0.0 |
| 60-72 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 2.2 | 0.0 |
| 72-84 | | 0.0 | 0.0 | | | 0.0 | 4.2 | 0.0 |
| | | | | | | | | |

^{*} reproduced from reference #11.

Appendix D TABLE 4* WR6026

| | | | | AMS/ML | | | | |
|----------------|----------------|--------------|--------------|--------------|--------------|-----------------|--------------|--------------|
| SAMPI.E | PT 1 =====: | PT 2 | PT 3 | PT 4 | PT 5 | PT 6 | PT 7 | PT 8 |
| -12-0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0-12 | 157 | 149 | 148 | 254 | 90 | 119 | 706 | 230 |
| 12-24 | 56 | 20 | 71 | 155 | 70 | 106 | 396 | 103 |
| 24-36 | 25 | 10 | 59 | 61 | 33 | 32 | 61 | 20 |
| 36-48 | 26 | 0 | 18 | 18 | 15 | 34 | 128 | 14 |
| 48-60 | 3 | 0 | 13 | 23 | 8 | 17 | 49 | 2 |
| 60-72 | 9 | 0 | 8 | 20 | 4 | 23 | 24 | 3 |
| 72-84 84-96 | 9 0 | 0 | 17 0 | 12 0 | 6 3 | 0 0 | 0 0 | 0 0 |
| 04-36 | Ü | U | 0 | U | 3 | U | U | U |
| | | | URINE | VOLUME | (ML) | | | |
| SAMPLE | PT 1 | PT 2 | PT 3 | PT 4 | PT 5 | PT 6 | PT 7 | PT 8 |
| ========= | | | | ====== | | | ===== | |
| -12-0 | 680 | 590 | 1045 | 735 | 815 | 1550 | 1500 | 1920 |
| 0-12 | 820 | 960 | 1390 | 510 | 485 | 1230 | 960 | 1135 |
| 12-24 | 1050 | 2020 | 1855 | 595 | 1050 | 1345 | 1300 | 1525 |
| 24-36 | 1835 | 1330 | 1328 | 432 | 802 | 1960 | 1605 | 2590 1615 |
| 36-48 48-60 | 830 1750 | 1630 1426 | 3130 1695 | 745 685 | 1160 1000 | 1480 950 | 1505 1095 | 2600 |
| 60-72 | 1315 | 1860 | 1805 | 490 | 1070 | 1635 | 1195 | 1890 |
| 72-84 | 715 | 1410 | 1650 | 1050 | 750 | 1600 | 1565 | 1860 |
| 84-96 | 790 | 1720 | 1720 | 745 | 1025 | 1570 | 1460 | 1970 |
| | | | | | | | | |
| | | TOTAL | | red/sam | | ICROMO | LES) | |
| SAMPLE | PT 1 | PT 2 | PT 3 | PT 4 | PT 5 | PT 6 | PT 7 | PT 8 |
| ======= | | | | | | | | |
| -12-0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 0-12 | 0.38 | 0.42 | 0.60 | 0.38 0.27 | 0.13 | 0.43 | 1.97 | 0.76 0.46 |
| 12-24 24-36 | 0.17 | 0.12 | 0.38 | 0.27 | 0.21 | 0.19 | 1.50 0.29 | 0.46 |
| 36-48 | 0.06 | 0.00 | 0.16 | 0.04 | 0.05 | 0.14 | 0.56 | 0.07 |
| 48-60 | 0.01 | 0.00 | 0.07 | | 0.02 | 0.05 | 0.16 | 0.01 |
| 60-72 | 0.04 | 0.00 | | | 0.01 | 0.11 | 0.08 | 0.02 |
| 72-84 | 0.02 | 0.00 | | | 0.01 | 0.00 | 0.00 | 0.00 |
| 84-96 | 0.00 | 0.00 | 0.00 | 0.00 | 0.01 | 0.00 | 0.00 | 0.00 |
| | | | | | | | | |
| TOTAL | 0.82 | 0.57 | 1.56 | 0.87 | 0.53 | 1.33 | 4.57 | 1.47 |
| | | | | | | | | |
| | | RATE | OF EYC | RETION | (NANO) | (A) [. P.G. / I | inip i | |
| SAMPLE | PT 1 | | PT 3 | | - | - | | PT 8 |
| | | | | | | | | |
| -12-0 | 0.0 | | | | 0.0 | | | |
| 0-12 | | | 49.8 | 31.5 | | | 164.6 | |
| | | | 31.9 | | | 34.5 | 125.1 | |
| | 11.2 | | | 6.4 | | | 23.9 | |
| | | | | 3.3 | | | | 5.6 |
| | 1.2 | | | 3.8 | | 3.9 | | |
| 60-72 | | | | 2.4 | | | | |
| | | | | 3.1 | | | 0.0 | |
| dr-40 | U.U | U. U | , 0.0 | . 0.0 | U.8 | U. U | U.U | U. U |

^{*} reproduced from reference #11.

8

1

8

X

page 1

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | | |
|---------------------|----------------|------|-----------------------------------|--|--|
| LIETMAN | E - W F M L | 01 | PROTOCOL:DAMD 17-85-C-5133- 03 | | |

PROTOCOL

| Study Day | Date ddmmmyy | Procedures |
|--------------|--------------|------------------------|
| | 25Jun86 | Screening laboratory |
| | czJul86 | History, Physical Exam |
| 0 | 05 Jul 86 | Admission |
| 3 | 07JUL36 | Drug Administration |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of $\frac{14}{2}$ pages, for subject $\frac{2}{2}$.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Kut / Filly M.D.

16/Dec/86

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | | |
|---------------------|--------------|------|-----------------------------------|--|--|
| LIETMAN | E W F M L | 01 | PROTOCOL:DAMD 17-85-C-5133- 03 | | |

MEDICAL HISTORY

Date of evaluation 02/Jul/86 dd mmm yy

Examiner

But Fritty N'I

Date of birth 26/Jan/64 dd mmm yy

print nam

Age <u>22</u>yrs

Sex M

30

**

2

Race \mathcal{B}_{-}

| | No | Yes | Comments |
|------------------------------|--------------|----------|--|
| Allergy | ~ | | |
| Tobacco Use | | V | 1/2 ppd |
| Alcohol Use | | / | 6 pack/week |
| Recreational Drug Use | | ✓ | MJ~Turek |
| Medications past 2 weeks | V | | |
| Experimental Drug Exposure | | ✓ | Jни |
| Blood or plasma donor | J | | |
| Prior Surgery | | / | (R Knee Sungery & MUA agak |
| Eye, ear, nose, throat | | V | (R Knee Swigery & MUA ag 6 No flow) "Sniffles" x That plung incu. Conductor. |
| Endocrine(diabetes, thyroid) | ✓ | | 10 |
| C-V (heart murmur, HBP) | ✓ | | |
| Pulmonary (cough, asthma) | \checkmark | | |
| Hepatitis, gastro-intestinal | √′ | | |
| Genito-urinary | | V | RXD & injection Lues 3-1 (Kata age |
| Musculoskeletal | V | | / |
| Neuropsychiatric | / | | |
| Other | | | |
| | | | |
| | <u> </u> | <u> </u> | <u> </u> |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | | OI | PROTOCOL:DAMD 17-85-C-5133- 03 |

B

3

8

2

PHYSICAL EXAMINATION

Date 02/Jul/86 dd mmm yy

| 1 | Temperature | | | | | <u> </u> | | |
|---|----------------|-----------|--------|----|------|----------|-----|------|
| | <u>3 5.4 c</u> | Q 4 2/min | 12/min | 12 | 8158 | 1. 7 | 6.0 | 73.5 |

| GENERAL EXAMINATION: | | | | | | | | | |
|----------------------|--------------|----------|--|--|--|--|--|--|--|
| (check) | Nor. | Abn. | Provide details of abnormalities | | | | | | |
| Head/Neck | / | | | | | | | | |
| EENT | | V | lymphoid exerseences post Atazuny | | | | | | |
| Chest, lungs | (3) | v | hight chet slightly deformed from bour and/or ort tusice (ely muscle) at 200 costochondal articulation, first nuter 1992 | | | | | | |
| Heart | \sqrt | | | | | | | | |
| Abdomen | ✓ | | | | | | | | |
| Genitalia | | | Not done | | | | | | |
| Rectal | | | Not done | | | | | | |
| Extremities | / | | | | | | | | |
| Skin | | ✓ | tattoos @ ant chest + Duppic arm + a ha. | | | | | | |
| Neurologic | V | } | | | | | | | |

CHEST X-RAY
Date p2/Jul/86

| NORMAL | X | ABNORMAL | Describe | abnormalities: |
|--------|---|----------|----------|----------------|
| | | | | |

Examiner

print name

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND:WR6026 |
|---------------------|----------------|------|-----------------------------|
| LIETMAN | E - W F M L | 01 | PROTOCOL:DAMD 17-85-C-5133- |

MEDICATION RECORD

STUDY: WR6026

| | | | End drug admin- istration (0-2400) | | Bottle I.D. # |
|---|--------|------|---------------------------------------|----|------------------|
| 3 | 07JU86 | 0802 | NA. | PO | |

* Below Assay Sensitivity

DOSAGE (total) 60 mg

PLASMA CONCENTRATIONS

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo- globin % |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|----------|-------------------------|
| | 0 | c 7Jul 86 | 0748 | 0748 | * | c /· |
| | 0.25 | c + Jul 86 | 0817 | 0817 | * | , |
| | 0.50 | c7)ul 86 | c 832 | C & 32 | * | |
| | 0.75 | 079ul86 | (847 | 1847 | * | |
| | 1.0 | 0+, 4186 | 1.902 | 0902 | 8.23 | |
| | 1.25 | 079-186 | 6917 | 1917. | 29.8 | |
| | 1.50 | 079486 | 0932 | 0932 | X.8 | |
| | 2.0 | 077486 | 1002 | 1002 | 48.7 | |
| | 2.5 | 07 Jul 86 | 1032 | 1632 | 51.0 | |
| | 3.0 | 077 ul Sk | 1102 | 1102 | 53.1 | |
| | 3.5 | 07 Jul 86 | 1132 | 1132 | 57.2 | |
| | | c7 Jul 86 | 1202 | 1202 | 49,2 | |
| · | E ^ | 079ul 86 | /302 | 131.2 | 44.4 | |
| | _ | 07Gul 86 | 1402 | 1412 | 45.9 | 14% |
| | | 079486 | 1612 | 1602 | 31.4 | |
| | امما | 079-186 | 1812 | 1862 | 32.3 | |
| | 14 | 07 Jul 8 is | 3662 | 3002 | 197 | |

8

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | E-W FML | 01 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICATION RECORD

STUDY: WR6026

| Study Day | Date ddmmmyy | Start drug admin- istration (0-2400) | End drug admin- istration (0-2400) | | Bottle I.D. # |
|--------------|--------------|---|---------------------------------------|----|------------------|
| 3 | 079ul86 | 0802 | N.A. | PO | |

* Below Assay Sensitivity

PLASMA CONCENTRATIONS

page 2

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo- globin % |
|---------------------------------------|-----------------------------|-------------------------------------|-------------------------------------|----------------------------------|----------|-------------------------|
| | 24.0 | 05Jul86 | CSC2 | 0802 | 9.80 | 0.5 |
| | 36.0 | ce.Jul 86 | 2002 | 2002 | * | |
| | 48.0 | 09 Tu (86 | 0862 | 0802 | * | c.7 |
| | 60.0 | 09gul 86 | 2002 | 2002 | * | |
| | 72.0 | 10 Jul 86 | 0802 | 080a | * | |
| | 84.0 | 10 Jul 86 10 Jul 86 11 Jul 86 | 2002 | 2002 | * | |
| | 96.0 | 11 Jul 86 | 0802 | 0802 | * | 10 |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| · · · · · · · · · · · · · · · · · · · | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | 87 | | | |

%

8

*

8

Ä

8

₹

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | 1 N L | 01 | PROTOCOL:DAMD 17-85-C-5133- 03 |

URINE CONCENTRATIONS

STUDY:WR6026

| Study | Date | Start drug admin- | End drug admin- | Route | Bottle |
|-------|----------|--------------------|--------------------|-------|--------|
| Day | ddmmmyy | istration (0-2400) | istration (0-2400) | | I.D. # |
| 3 | 07.Jul86 | 0802 | NA | PO | |

DOSAGE (total) 60 mg

URINE CONCENTRATIONS

| Sample | Scheduled | Start Co | ollection | End Col | lection | Total Volume | [WR6026] | |
|--------|----------------------------|------------|-----------|-----------|---------|-----------------|-----------|--|
| No. | Collection Time (hours) | ddmmmyy | 0-2400 | ddmmmyy | 0-2400 | (ml) | UN) | |
| U01 | -24 TO -12 | Ck Jul St | 0800 | 06 Jul 86 | 2000 | 1315 | | |
| nos | -12 TO O | 06 Jul Blo | 2000 | 07 Jul 86 | 0800 | 680 | 0 | |
| П03 | 0 TD 12 | 07 jul 8/2 | 0800 | 0794186 | 2000 | 826 | 0.16742 | |
| U04 | 12 TO 24 | 0794186 | 2000 | 0892186 | 0800 | 105C | 0.04335 | |
| U05 | 24 TO 36 | 08 Jul 86 | 0800 | 08 Jul 86 | 2000 | 1835 | 0.03853 | |
| N0E | 36 TO 48 | 08 Jul 86 | 2000 | 099ul 86 | 0800 | 83c | 0.01826 | |
| U07 | 48 TO 60 | 09 Jul 86 | 0800 | 09 Jul 86 | 2000 | 1750 | V00 2 052 | |
| 80U | 60 TD 72 | 09 Jul 86 | 2000 | 10 Jul 86 | 0800 | 1315 | 0.010388 | |
| U09 | 72 TO 84 | 10 Jul 86 | 0800 | 10 Jul 86 | 2000 | 715 | D.005648 | |
| U10 | 84 TO 96 | 10 Jul 86 | 2000 | Digue 86 | 0800 | 790 | 0 | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| * | | | 88 | | | | ~ | |

| INVESTIGATOR'S NAME | PΤ | INI. | TIAL | PΤ | # | COMPOUND: WR6026 |
|---------------------|-----|------|------|----|---|-----------------------------------|
| LIETMAN | E F | M | 718 | 0 | / | PROTOCOL:DAMD 17-85-C-5133- 03 |

8

85

K W

X

**

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| | Screen - | Predrug 1 | Predrug 23 | Post 4 | tdrug 5 | Day |
|----------------------|--------------------|--------------------|---------------|--------------------|------------|------|
| TEST:NORMAL | 35Jun86 ddmmmyy | OSJU186 ddmmmyy | ddinmmyy | CSTul&6 ddmmmyy | | Date |
| NA:135-148 MEQ/L | 142 | 138 | <i>i</i> 42 | 139 | 142 | |
| K:3.5-5.0 MEQ/L | 3.9 | 3.8 | 4.6 | 4.3 | 4.1 | |
| CL:96-109 MEQ/L | 110 | 107 | 104 | 105 | 101 | |
| CO2:24-30 MEQ/L | 24 | 22 | 23 | 24 | 25 | |
| SUN:12-25 MG/DL | 16 | 26 | 12 | 13 | 14 | |
| CREAT:0.4-1.5 MG/DL | 1.0 | 1.1 | 1.0 | 1.0 | 1.0 | |
| GLU:70-115 MG/DL | 96 | £ 5 | 82 | 88 | 91 | |
| T.BILI:0.3-1.2MG/DL | 1.4 | 1.3 | 1.6 | 1.1 | 1.3 | |
| D. BILI:0.1-0.4MG/DL | c.1 | 0.2 | 0.2 | 0.1 | 0.1 | |
| CA:9.0-11.0 MG/DL | 9.5 | 9.1 | 9.3 | 9.4 | 10,1 | |
| PO4:3.0-4.5 MG/DL | 4.4 | 4.1 | 4,4 | 4.6 | 5.2 | |
| URIC A:4.2-8.8MG/DL | 5.5 | 5.8 | 4.1 | 4.1 | 4.3 | |
| T. PROT:6.0-8.5G/DL | 6.7 | 6.7 | 6.4 | 6.3 | 7.5 | |
| ALB.:3.2-5.3 G/DL | | 4.6 | 4,5 | 4,5 | ND | |
| AST:0-35 IU/L | 19 | 36 | 16 | 25 | 31 | |
| ALT:0-30 IU/L | 10 | 15 | 15 | 14 | 21 | |
| ALK PHOS:0-95 IU/L | 44 | 52 | 45 | 42 | 47 | |
| CHOL:151-268 MG/DL | 203 | 201 | 228 | 205 1390 | 137 | |
| LDH:0-200 IU/L | | 183 | 150 | /39 | 137 | |
| CPK:0-160 U/L(male) | _ | 712 | 434 | 557 | 247 | |
| TG:20-190 MG/DL | 54 | _ | 66 | 58 | 69 | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | E - W F M L | c/ | PROTOCOL:DAMD 17-85-C-5133- 03 |

Ä

30

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Postdruj 7 | g | | | | Day |
|---------------------|--------------------|---------|---------|---------|---------|-----|
| TEST:NORMAL | 11Ju/86 ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy | Dat |
| NA:135-148 MEQ/L | 140 | | | | | |
| K:3.5-5.0 MEQ/L | 4.3 | | | | | |
| CL:96-109 MEQ/L | 99 | | | | | |
| CO2:24-30 MEQ/L | 29 | | | | | |
| SUN:12-25 MG/DL | 18 | | | | | |
| CREAT:0.4-1.5 MG/DL | 1.0 | | | | |] |
| GLU:70-115 MG/DL | 80 | | | | | |
| T.BILI:0.3-1.2MG/DL | 1.3 | | | | | |
| D.BILI:0.1-0.4MG/DL | 0,0 | | | | | |
| CA:9.0-11.0 MG/DL | 14.5 | | | | | |
| PO4:3.0-4.5 MG/DL | 4.9 | | | | | } |
| URIC A:4.2-8.8MG/DL | 4.8 | | | | | |
| T. PROT:6.0-8.5G/DL | 7.6 | | | | | |
| ALB.:3.2-5.3 G/DL | 5.1 | | | | | |
| AST:0-35 IU/L | 27 | | | | |] |
| ALT:0-30 IU/L | 13 | | | | | |
| ALK PHOS:0-95 IU/L | 42 | | | | |] |
| CHOL:151-268 MG/DL | 279 | | | | | |
| LDH:0-200 IU/L | 217 | | | | | |
| CPK:0-160 U/L(male) | AN | | | | | |
| TG:20-190 MG/DL | ND. | | | | | |

page 9

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | E U | c.I | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| | • | Screen - | Predrug 1 | Predrug 3 | Postdrug 4 5 | | Day |
|----------|------------|-------------|--------------|--------------|-----------------|--------------------|------|
| TEST | NORMAL | 2.5Jun86 | | | 08JUL 86 | 09Jul86 ddmmmyy | Date |
| MBC | 4500-11000 | 5600 | 6200 | 580c | 55cc | 5000 | |
| RBC | 4.50-5.90 | 4.80 | 4.65 | 5.10 | 4.90 | 5.it | |
| Hgb | 13.9-16.3 | 14.4 | 13.7 | 15.0 | 14.7 | 15.4 | |
| PCV | 41.0-53.0 | 42.4 | 41.0 | 45.2 | 43.8 | 45.4 | |
| Plt | 150-350 | 304 | 281 | 271 | 264 | 285 | |
| Bands | 2-6× | 0 | 5 | | 2. | | |
| Polys | 31-76% | 54 | 45 | 5¢ | 42 | 54 | |
| Eos | 1-4% | á | 1 | C | 5 | 4 | |
| Bas | | 0 | 1 | C | 2 | C | |
| Lymphs | 24-44% | 4C | 36 | 36 | 35 | 38 | |
| Atyp Lym | | 2 | 1 | 1 | 1 | t | |
| Monos | 2-11% | 2 | 11 | 13 | 13 | 4 | |
| Other | | 0 | 0 | O | 0 | C | |
| | | | | | | | |
| Methemi | | 0.3% | | 0.0 | 0.3 | 0,7 | |
| G-6-PD | 7.4-9.4 | 6.9 | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |

page 10

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | E - W F M L | CI | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug 7

Day

| | | | | , | | | 1 |
|----------|-------------|--|--------------|--------------|---------|---------|-----|
| TEST | NORMAL | 11JUSC ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy | Dat |
| MBC | 4500-11000 | 7000 | | | | | |
| RBC | 4.50-5.90 | 5.07 | | | | | |
| Ндь | 13.9-16.3 | 15.2 | | | | | |
| PCV | 41.0-53.0 | 45.9 | | | | | |
| Plt | 150-350 | 302 | | | | | |
| Bands | 2-6% | 4 | | | | | |
| Polys | 31-76% | 45 | | | | | |
| Eos | 1-4% | 2 | | | | | |
| Bas | | Ĉ: | | | | | |
| Lymphs | 24-44% | 37 | | | | | |
| Atyp Lym | | to | | | | | |
| Monos | 2-11% | 12 | | | | | |
| Other | | 0 | | | | | |
| | | | | | | | |
| methem | | 1.0 | | | | | |
| G-6-PD | 7.4-9.4 | | | | | | |
| | | | | | | | |
| | | | | | | | |
| } | | | | | | | |
| | | | | | | | 1 |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WREOZE |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | E - W F M L | 01 | PROTOCOL:DAMD 17-85-C-5133- 03 |

URINALYSIS VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Screen Predrug - 1 | | Postdrug 4 5 | | 7 | |
|----------|-----------------------|--------------------|--------------------|--------------------|----------|---------------------------|
| | | 02Jul86 ddmmmyy | 05Jun86 ddmmmyy | CEJu(86 ddmmmyy | | <u>//Ju/86</u> ddmmmyy |
| 5p.Gr. | | 1.034 | 1.03元 | 1016 | 1.017 | 1,629 |
| рн | | 6.0 | 6.5 | 7.0 | 7.0 | 7.0 |
| Protein | V | 4 France | Neg | NEW TO | New (sp) | Ney Trace 47) |
| Glucose | | Nea | Neg | Neg | Neg | Neg |
| Ketones | | Neg | Neg | Neg | Nea | Neg |
| Bili. | | Neg | Neg | Neg | Neg | Neg |
| ūcc.Bld. | | Neg | Neg | Neg | Neg | Nog |
| Cast/lpf | | c / | C | 0 | 0 | 0 |
| WBC/hpf | | C | C | 0 | 0 | \mathcal{C} |
| RBC/hpf | | D. | C | O | O | C |
| Epi./hpf | | C | C. | 0 | C | 0-1 |
| Crys/hpf | | O | t c | 0 | D | O |
| Bact/hpf | | C. | 0 | 0 | 0 | 0 |

ELECTROCARDIOGRAM

| Date ddmmmyy | Time 0-2400 | | ABNORMAL check | Describe abnormalities |
|-----------------|----------------|-----|-------------------|---|
| 02July 86 | 11.3c | | / | non specific Tunue changes, przeibly a roban placin |
| 07July86 | C725 | | | son-specific Twave changes |
| 07Jul 86 | 1157 | | V | non specific Twee A, 2° ban placement |
| 08JW 86 | 0750 | V | | l' l' |
| 109 Til 86 | 0750 | V . | | |

Scrun

93

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | <u>ال</u> | C I | PROTOCOL:DAMD 17-85-C-5133- 03 |

ADVERSE EXPERIENCES

check if none occurred



| | OVERSE EVENT | ONSET (Date/Time) | CESSATION Date/Time | | REL'N TO TEST DRUG | ACTION (Check) |
|---|-----------------|----------------------|------------------------|------------|-----------------------|-------------------|
| * | | 11 JJ B | dd mmm yy | Mild | DEF. | None |
| 1 | Increase in | <u>0890</u> | (0-2400) | Mod | PROB. | |
| | | | | □ Sev | DEF. NOT | Stop test |
| * | | dd mmm yy | dd mmm yy | Mild | DEF. | drug |
| | | (0-2400) | (0-2400) | ☐ Mod | PROB. POSS. | Treatment |
| | | | | Sev | DEF. NOT | Stop test drug |

COMMENTS (Indicate # and event)

page 13

| | | | | · · | |
|---------------------------------------|-----------------------------------|--|-------------------------|---------------|---------------------|
| INVESTIG | 026 | | | | |
| LIETM | IAN , | F M L | 01 | PROTOCOL:DAMD | 17-85-C-5133- 03 |
| | (То | | UTCOME ed for | all subjects) | |
| Ø | Protocol co | ompleted | | | 11Juler ddmmmyy |
| | Premature t | ermination | of pro | otocol | ddmmmyy |
| | (Cr Adverse Exp Died During | g Study Return For operate iolation ation nt Illness | riate (| category) | |
| If terminated early, explain briefly: | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |

| PT INITIAL | PT # | СОМРОИ | ND: WR602 | 6 | |
|-------------------------|------------------------------|---|--|--|--|
| E - N F M L | 01 | PROTOC | OL:DAMD 1 | 7-85- C-5133 03 | |
| CONCOMITANT MEDICATIONS | | | | | |
| | | | DATE STARTED | DATE STOPPED | |
| heparin flush | | wuts | c7ful 86 | 10JU178 | |
| | | · · · · · · · · · · · · · · · · · · · | | | |
| | | | | | |
| | | comme | nts) | | |
| | | | | | |
| | E - N F M L CONCOMITAN | E - N F M L CI CONCOMITANT MEDIC DAILY 344 - 1444 | E - N CI PROTOC CONCOMITANT MEDICATIONS TOTAL DAILY DOSE 300 - 1000 mults | PT INITIAL PT # COMPOUND: WREOZ E - N PROTOCOL:DAMD 1 CONCOMITANT MEDICATIONS TOTAL DATE DAILY DOSE STARTED 300 - 1000 mults C7 ful 86 | |

X

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | |
|---------------------|------------|------|-----------------------------------|--|
| LIETMAN | RLH | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

PROTOCOL

| Study Day | Date ddmmmyy | Procedures |
|--------------|---------------|----------------------|
| | 09Jul86 | Screening laboratory |
| | 10 Jul 86 | |
| 0 38 | 12 05]a186 | Admission |
| 3 | # Jul 86 | Drug Administration |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

I certify that:

Ž.

- 1) I have carefully examined all entries on the data collection forms, consisting of 14 pages, for subject 4 02.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Brent Mathy M.D.

signature

161 Dec 1 12 dd/mmm/yy

ይሄውት መቀመቀው የመቀመቀው የመቀመቀ የመቀመቀው የመቀ

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | R L H | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICAL HISTORY

Date of evaluation $\frac{10}{\text{Jul}} \frac{86}{\text{dd}}$

Examiner

Kinte Mitty My)

Date of birth

13 / A us/ 57 dd mmm yy

orint name

Age

X S

8

%

3

28 yrs

Sex

<u>M</u>

Race

В

| | No | Yes | Comments |
|------------------------------|----|----------|-----------------------------|
| Allergy | V | | |
| Tobacco Use | | √ | 1/2 PPD |
| Alcohol Use | / | | Quit 6 menths age |
| Recreational Drug Use | / | | Quit 6 months ago |
| Medications past 2 weeks | ✓ | | |
| Experimental Drug Exposure | | ✓ | Johns Hopkins Hospital |
| Blood or plasma donor | | / | August 1985 |
| Prior Surgery | | / | Appy 1978, 7+A 1975 |
| Eye, ear, nose, throat | ~ | | ,,,, |
| Endocrine(diabetes, thyroid) | 1 | | |
| C-V (heart murmur, HBP) | / | | |
| Pulmonary (cough, asthma) | ✓ | | |
| Hepatitis, gastro-intestinal | / | | |
| Genito-urinary | | / | GC-1978, Lues 1978, Rxd PCN |
| Musculoskeletal | / | | |
| Neuropsychiatric | ~ | | |
| Other | / | | |
| | | | |

page 3

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | R L H F M L | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 |

PHYSICAL EXAMINATION

Date 10/Jul/86 dd mmm yy

| Temperature | | | | | f | 1 1 |
|---------------|-----------|--------|-----|------|-------|------|
| <u>35.9</u> c | _ 6 d/min | 14/min | 1_0 | 8,72 | 178.0 | 73.6 |

| | GENERAL EXAMINATION: | | | | | | | |
|--------------|----------------------|----------|---|--|--|--|--|--|
| (check) | Nor. | Abn. | Provide details of abnormalities | | | | | |
| Head/Neck | ✓ | | Neck-Shetty nodes | | | | | |
| EENT | √ | | throat numeral hypophoid tissue in (k to is in poss | | | | | |
| Chest, lungs | V | | | | | | | |
| Heart | 1 | | | | | | | |
| Abdomen | V | | | | | | | |
| Genitalia | | · | Not dong | | | | | |
| Rectal | | | not done | | | | | |
| Extremities | ~ | | | | | | | |
| Skin | | ✓ | Appy scar, antehulital scar, foreheard scar | | | | | |
| Neurologic | / | | | | | | | |

CHEST X-RAY
Date 09/Jen/86

IPSS

| NORMAL | χ | ABNORMAL | Describe abnormalities: |
|--------|---|----------|-------------------------|
| | | | • |

Examiner

Dr. B.G. Pety

print name

5

汉

XX XX

**

8

8

No.

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND:WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | RLH FML | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICATION RECORD

STUDY: WR6026

| Study | Date | Start drug admin- | End drug admin- | Route | Bottle |
|-------|---------|--------------------|--------------------|-------|--------|
| Day | ddmmmyy | istration (0-2400) | istration (0-2400) | | I.D. # |
| 3 | # Jan86 | 0802 | . A, N | PO | |

DOSAGE (total) 60 mg

* Below Assay Sensitivity

PLASMA CONCENTRATIONS

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Metheme= globin -*- |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|------------------|---------------------------|
| | 0 | 14Jul 86 | 0745 | 0745 | * | |
| | 0.25 | 14Jul86 | 08/7 | 0817 | * | |
| | 0.50 | 14 Jul 86 | 0832 | 0832 | 7.26 | |
| | 0.75 | 14 Jul 86 | 0847 | C847 | 30.5 | |
| | 1.0 | 14 Jul 86 | 0902 | 0902 | 50.9 | |
| | 1.25 | 14 Jul 86 | C917 | 0917 | 60.3 | |
| | 1.50 | 14 Jul 86 | 0932 | 0932 | 75.4 | |
| | 2.0 | 14 Jul 86 | 1002 | 1002 | 56.3 | |
| | 2.5 | 14 Jul 86 | 1032 | 1032 | G7.0 | |
| | 3.0 | 14 Jul 86 | 1102 | 1102 | 59.4 | |
| | 3.5 | 14 Jul 86 | 1132 | 1132 | 69.5 | |
| | 4.0 | 14 Jul 86 | /aca | 1202 | 59.8 | |
| | 5.0 | 14 Jul 86 | 13C2 | 1302 | 51.4 | <u> </u> |
| | 6.0 | 14 Jul 86 | 1402 | 1402 | 36.1 | |
| | 8.0 | 14 Jul 86 | 1602 | 1602 | 8.PG | <u></u> |
| | 10.0 | 14 Jul 86 | 1812 | 1862 | 3 1.5 | |
| | 12.0 | 14 J W86 | | 200,2 | J 1 | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | | |
|---------------------|----------------|------|-----------------------------------|--|--|
| LIETMAN | R L H F M L | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 | | |

MEDICATION RECORD

STUDY: WR6026

 \ddot{x}

N.

8

% %

8

8

8

85

| Study Day | Date ddmmmyy | Start drug admin- istration (0-2400) | End drug a istration | admin- (0-2400) | Route | Bottle I.D. 4 | |
|--------------|--------------|---|----------------------|--------------------|-------|------------------|--|
| 3 | 14JW 86 | 0802 | NA. | | PO | | |

* Bebw Assay Sensitivity DOSAGE (total) _____ 60 mg

ASMA CONCENTRATIONS

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo- globin |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|----------|--------------------|
| | 24.0 | 15JW86 | C802 | 0802 | * | |
| | 36.0 | 15 Jul 86 | 2002 | 2002 | * | |
| | 48.0 | 16Jul86 | 0802 | 0802 | * | |
| | 60.0 | 16Ju/86 | 2002 | 2015 | * | |
| | 72.0 | 17Jul86 | 0802 | 1302 | * | |
| | 84. C | 17JW86 | 3003 | 3702 | * | |
| | 96.0 | 18Jul 86 | 6363 | 0803 | * | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | 101 | | | |

page 6

| INVESTIGATOR'S NAME | PT INITIAL P | T # | COMPOUND: WR6026 | |
|---------------------|----------------|-----|-----------------------------------|--|
| LIETMAN | R L H F M L | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

URINE CONCENTRATIONS

STUDY: WR6026

| | | Start drug admin- istration (0-2400) | | | Bottle I.D. # |
|---|---------|---|------|----|------------------|
| 3 | 14Jul86 | 0802 | N.A. | PO | |

DDSAGE (total) 60 mg

URINE CONCENTRATIONS

| Sample No. | · | | lection | Total Volume | [WR6026] | | | | |
|---------------|-------|--------|------------|-----------------|-----------|-----------|--------|-------|-------------------|
| 140. | | nours) | - 4 iii ee | ddmmmyy | 0-2400 | ddmmmyy | 0-2400 | (ml) | ma |
| U01 | -24 1 | ro -12 | | 13Jul 86 | 0800 | 13 Jul 86 | 2000 | 1355 | (|
| บงอ | -12 7 | ro o | | 13Jul 86 | → | 13Jul86 | 0800 | S | O HEADER |
| U03 | 0 1 | ro 12 | | 14JU186 | 0800 | 14Ju/86 | 2000 | 960 | 0.11904 |
| U04 | 12 7 | TO 24 | | 14 Jul 86 | 2000 | 15JU186 | 0800 | 3030 | 0.03434 G-0104 |
| U05 | 24 1 | TO 36 | | 15Jul 86 | 0800 | 15Jul 86 | 20083 | 1330 | 0.0064 |
| noe | 36 1 | TO 48 | 9 | 15 HUJU186 | 20082 | 16Jul86 | 08082 | 1636 | 0 |
| U07 | 48 1 | TO 60 | | 16Jul 86 | 0800295 | 16 Jul 86 | | 1436 | 0 |
| UOB | 60 1 | ro 72 | | 16Jul86 | | 17 Jul 86 | 0800 | 1860 | 0 |
| U09 | 72 1 | TO 84 | | 17Jul86 | | 17 Jul 86 | 2000 | 1410 | 0 |
| U10 | 84 1 | ro 96 | | 17 Jul 86 | 2000 | 18 Jul 86 | 0806 | 17.3C | 0 |
| | | | | | | | | | |
| | | ·· | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | * Smt | urine | Jus | auss ot. | failed to | accusat | ely sa | r ur | nco |
| | , , | | | 186 at 20 | 1 | | , | | ٧ |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | R L H F M L | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 |

2

8

X

3

2

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Screen - | Predrug 1 | Predrug 2 | Post 4 | tdrug 5 | Day |
|------------------------|-------------|--------------|--------------------|---------------------|----------------------|------|
| TEST:NORMAL | 69Ju/86 | | 1454/86 ddmmmyy | 15Jul 16 ddmmmyy | 16 Jul 20 ddmmmyy | Date |
| NA:135-148 MEQ/L | 144 | 144 | 139 | 145 | 140 | |
| K:3.5-5.0 MEQ/L | 4./ | 3.9 | 4.3 | 4.1 | 4.5 | |
| CL:96-109 MEQ/L | 111 | 113 | 106 | 104 | 136 | |
| CO2:24-30 MEQ/L | 25 | 19 | 22 | 25 | 22 | |
| SUN:12-25 MG/DL | 14 | 13 | 10 | 12 | 15 | |
| CREAT:0.4-1.5 MG/DL | 1.3 | 1.4 | 1.1 | 1.2 | 1.2- | |
| GLU:70-115 MG/DL | 60 | 84 | 70 | רר | 76 | |
| T.BILI:0.3-1.2MG/DL | 0.7 | 0-3 | 0.6 | 0.5 | 0.4 | |
| D. BILI:O. 1-O. 4MG/DL | 0.0 | 0.0 | 0.1 | 0.1 | 0.2 | |
| CA:9.0-11.0 MG/DL | 9.4 | 8.7 | 9.4 | 10-1 | 9.3 | |
| PO4:3.0-4.5 MG/DL | 4.1 | 3.1 | 3.7 | 3.5 | 4.0 | |
| URIC A:4.2-8.8MG/DL | 7.5 | 5.4 | 5.0 | 5.1 | 5.7 | |
| T. PROT:6.0-8.5G/DL | 6.3 | 5.8 | 6.4 | 6.7 | 6.5 | |
| ALB.:3.2-5.3 G/DL | ND | 3. 8 | 3- 9 | 4.4 | 4.2 | |
| AST:0-35 IU/L | 34 | 33 | 24 | 24 | J3 | |
| ALT:0-30 IU/L | 16 | 17 | 19 | 2.2 | 20 | |
| ALK PHOS:0-95 IU/L | 43 | 52- | 51 | 46 | 44 | |
| CHQL:151-268 MG/DL | 140 | 125 | 167 | 174 | 164 | |
| LDH:0-200 IU/L | 158 | 144 | 145 | 122 | 131 | |
| CPK:0-160 U/L(male) | 570 | 635 | 296 | 283 | 245 | |
| TG:20-190 MG/DL | 69 | 12 For | 62 | 68 | 45 | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WREOZE |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | R L H F M L | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 |

CHEMISTRY VALUES

H

X

Laboratory: JOHNS HOPKINS HOSPITAL

| 7 | • | | | Day |
|-----------|---------|---------|---------|------|
| i [Jul Ro | ddmmmyy | ddmmmyy | ddmmmyy | Date |

| TEST: NORMAL | 18Jul Ro | ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy | Dat |
|----------------------|----------|---------|---------|---------|---------|-----|
| NA:135-148 MEQ/L | 141 | | | | | |
| K:3.5-5.0 MEQ/L | 4.3 | | | | | |
| CL:96-109 MEQ/L | /03 | | | | | |
| CO2:24-30 MEQ/L | 27 | | | | | |
| SUN:12-25 MG/DL | 14 | | | | | |
| CREAT:0.4-1.5 MG/DL | 1.3 | | | | | |
| GLU:70-115 MG/DL | 71 | | | | | |
| T. BILI:0.3-1.2MG/DL | 0.6 | | | | | |
| D. BILI:0.1-0.4MG/DL | 0.1 | | | | | |
| CA:9.0-11.0 MG/DL | 10,2 | | | | | |
| PO4:3.0-4.5 MG/DL | 4.2 | | | | | |
| URIC A:4.2-8.8MG/DL | 5.5 | | | | | |
| T. PROT:6.0-8.5G/DL | 7.3 | | | | | |
| ALB.:3.2-5.3 G/DL | 4.5 | | | | | |
| AST:0-35 IU/L | 29 | | | | | |
| ALT:0-30 IU/L | 27 | | | | | |
| ALK PHOS:0-95 IU/L | 47 | | | | | |
| CHOL:151-268 MG/DL | 183 | | | | |] |
| LDH:0-200 IU/L | 394 | | | | | |
| CPK:0-160 U/L(male) | 272 | | | | | |
| TG:20-190 MG/DL | 62 | | | | | 1 |

page 9

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | RLH FML | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUESLaboratory: JOHNS HOPKINS HOSPITAL

| | | Screen | Predrug 1 | Predrug 3 | Post 4 | drug 5 | Day |
|----------|------------|--------------------|--------------|---------------------|-----------|--------------------|------|
| TEST | NORMAL | 09Jul86 ddmmmyy | | 14Jul 26 ddmmmyy | | 161x136 ddmmmyy | Date |
| WBC | 4500-11000 | 8800 | 7800 | 7800 | 7700 | 7700 | |
| RBC | 4.50-5.90 | 4.65 | 4.31 | 4.93 | 5.06 | 4.99 | |
| Ндь | 13.9-16.3 | 14.5 | 13.4 | 15.6 | 15.5 | 15.6 | |
| PCV | 41.0-53.0 | 43.2 | 39.9 | 45.4 | 48.3 | 47.1. | |
| Plt | 150-350 | 253 | 238 | 272 | 309 | 274 | |
| Bands | 2-6% | 1 | 2 | 6 | 4 | 1 | |
| Polys | 31-76% | 45 | 45 | 49 | 38 | 49 |] |
| Eos | 1-4% | 3 | 1 | 1 | 2. | 4 |] |
| Bas | | 0 | 0 | 1 | 0 | 0 | |
| Lymphs | 24-44% | 41 | 46 | 32 | 45 | 40 | |
| Atyp Lym | | 0 | 0 | 0 | 0 | 0 | |
| Monos | 2-11% | 10 | 6 | 11 | 11 | 6 | |
| Other | | 0 | 0 | 0 | 0 | 0 | |
| | | 10 Jul 86 | | | | | |
| Methem: | , | 0.1 | | 0.0/0.1 | 0.0 | 0,0 | |
| G-6-PD | 7. 4-9. 4 | 7,8 | | 7 | | | |
| | | | | | | | |
| | | | | | | | } |
| | | | | | | | |
| | | | | | | |] |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | RLH FML | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOBY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug 7

*

Day

| TEST | NORMAL | 18Jul 16 ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy | qqwwmyy | Date |
|----------|------------|---------------------|---------|---------|---------|---------|------|
| WBC | 4500-11000 | 0003 | | | | | |
| RBC | 4.50-5.90 | 5,04 | | | | | |
| НдЬ | 13.9-16.3 | 15.6 | | | | | |
| PCV | 41.0-53.0 | 46.9 | | | | | |
| Plt | 150-350 | 288 | | | | | |
| Bands | 2-6% | 3 | | | | | |
| Polys | 31-76% | 48 | | | | | |
| Eos | 1-4% | 0 | | | | | |
| Bas | | / | | | | | |
| Lymphs | 24-44% | 40 | | | | | |
| Atyp Lym | | 0 | | | | | |
| Monos | 2-11% | 8 | | | | | |
| Other | | 0 | | | | | |
| | | | | | | | |
| Methem. | | 0.5 | | | | | |
| G-6-PD | 7.4-9.4 | | | | | | |
| | | | | | | |] |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WREOZE |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | RLH FML | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 |

URINALYSIS VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| | Screen Predrug | | | Postdrug | | |
|----------|--------------------|--------------------|----------|----------|--------------------|--|
| | | 1 | 4 | 5 | 7 | |
| | 1014186 ddmmmyy | 12Jul86 ddmmmyy | 15Jul 86 | ddmmmyy | qquuuuyy 187486 | |
| Sp.Gr. | 1.023 | 1.023 | 1.011 | i 1 | 1.017 | |
| рН - | 6.0 | 5.0 | 6.0 | | 6.5 | |
| Protein | Neg | Neg | Neg | | heg | |
| Glucose | Neg | Neg | Ncg | | hea | |
| Ketones | Neg | Neg | Neg | | heg | |
| Bili. | Neg | Neg | Neg | | hei | |
| Occ.Bld. | Neg | Neg | Neg | | heg | |
| Cast/lpf | 0 | ó | d | | 0' | |
| WBC/hpf | C | 0 | v | | 0-1 | |
| RBC/hpf | c | 0 | O | | 0 | |
| Epi./hpf | C | 0 | 0 | | 0 | |
| Crys/hpf | o | 0 | 0 | | 0 | |
| Bact/hpf | 0 | 0 | o | | 0 | |

K

ELECTROCARDIOGRAM

| Date ddmmmxy 10 141 75 | Time 0-2400 | NORMAL check | ABNORMAL check | Describe abnormalities Sinus biodycardia with occasional jumitinda lavabral dust |
|------------------------------|----------------|-----------------|-------------------|---|
| 14JW 86 | 0725 | ~ | | |
| 19 Jul 86 | 1213 | Derro, | V | nodal low atrial shythm |
| 15 Jul 86 | 0828 | 250 | | model o low atrial nhythm |
| 16Jul 86 | 0823 | V | | |
| 18Jul 86 | 0828 | eng. | / | node o low atrial shuthing |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | R L H | 02 | PROTOCOL:DAMD 17-85-C-5133- 03 |

ADVERSE EXPERIENCES



| , | OVERSE EVENT | ONSET (Date/Time) | CESSATION Date/Time | | REL'N TO TEST DRUG | ACTION (Check) |
|---|-----------------|-------------------------|------------------------|-----------|------------------------|-------------------|
| * | | 18 Jul 76 dd mmm yy | dd mmm yy | Mild | DEF. | None |
|) | Incurred LDH | <u>0 No</u> (0-2400) | William (0-2400) | Мод | PROB. | Treatment |
| | | | | Sev | DEF. NOT | Stop test drug |
| * | | dd mmm yy | dd mmm yy | ☐ Mild | DEF. | None |
| | | (0-2400) | (0-2400) | Mod | PROB. POSS. DEF. NOT | Treatment |
| | | | | Sev | UNKNOWN | Stop test drug |

COMMENTS (Indicate # and event)

| Prema Adver Died Failu Did N Proto Entry | (To | be completermi | completed nation | DUTCO | PROTOC | ibjects) | 17-85-C-5133- 03 18 46 <u>Julib</u> ddmmmyy |
|---|-----------------------------------|---|--------------------------|-------------|----------------------------------|----------|---|
| Proto Prema Adver Died Failu Did N Proto Entry Inter | ture f | be completermi | comple eted nation | n of | ME or all so protocol E TERMINA | ibjects) | 03 18 46 <u>Julib</u> ddmmmyy |
| Prema Adver Died Failu Did N Proto Entry Inter | ture f | omple termi ON FO | eted nation | n of MATURI | protocol | ATION | ddmmmyy |
| Prema Adver Died Failu Did N Proto Entry Inter | REAS(| termi DN FO | nation OR PREI | MATURI | E TERMINA | | ddmmmyy |
| Adver Died Failu Did N Proto Entry Inter | REAS(| DN FO | R PREI | MATURI | E TERMINA | | ddmmmyy |
| Died Failu Did N Proto Entry Inter | (Cf | heck | appro | | | | |
| If terminated | ot Cod col V Viola curre | Retuopera iolat ation nt Il tive/ | idy irn For ite ion i | | low-up | | |

page 14

| | | | | • | |
|---------------------|--|--------------------|------------|----------------|--------------------|
| INVESTIGATOR'S NAME | PT INITIA | - PT # | COMPOU | ND: WRE | 026 |
| LIETMAN | RLH | 02 | PROTOC | DL:DAMD | 17-85-C-5133 03 |
| | CONCOMITA | NT MEDIC | CATIONS | | |
| DRUG NAME | | TO' DAILY | AL DOSE | DATE STARTE | |
| hipasin flush | | 100-100 | Cunits | 14 Jul 8 | 6 17.71LPG |
| | | | | | |
| | | | | | |
| (please | CO: date and : | MMENTS sign all | commer | nts) | |
| | | | | | |
| <u> </u> | ······································ | | | ···· | |

25

X

page 1

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | W F | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 |

PROTOCOL

| Study Day | Date' ddmmmyy | Procedures |
|--------------|------------------|------------------------|
| | 08Jul 86 | Screening laboratory |
| | 10Jul 86 | History, Physical Exam |
| 0 | 38 WTP1 | Admission |
| 3 | 21Jul 86 | Drug Administration |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 4 pages, for subject # 03
- 2) All information entered onto the data collection forms for subject, to the best of my knowledge, is correct.

16, Dec, 86

dd/mmm/yy

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | W = F | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICAL HISTORY

Date of evaluation 10/Jul/86 dd mmm yy

Examiner

Dr.B.G. Petty

Date of birth

30/ALR/55 ad mmm yy

Age _3

31 yrs

Sex

M

Race

<u>B__</u>

| | No | Yes | Comments |
|------------------------------|---------|-----|----------------|
| Allergy | / | | Astura es inti |
| Tobacco Use | | ✓ | Yappo |
| Alcohol Use | | 1 | 1 beer/week |
| Recreational Drug Use | / | | |
| Medications past 2 weeks | <i></i> | | |
| Experimental Drug Exposure | V | | |
| Blood or plasma donor | / | | |
| Prior Surgery | / | | |
| Eye, ear, nose, throat | ~ | | |
| Endocrine(diabetes, thyroid) | ~ | | |
| C-V (heart murmur, HBP) | ~ | | |
| Pulmonary (cough, asthma) | ~ | | |
| Hepatitis, gastro-intestinal | / | | |
| Genito-urinary | r | | |
| Musculoskeletal | ~ | | |
| Neuropsychiatric | ~ | | |
| Other | • | | |
| | | | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | W - F F M L | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 |

Ä

X

27.6

 $\ddot{\aleph}$

3

深

É

3

رب برج برج

Ŝ,

PHYSICAL EXAMINATION Date 10/Jul/86 dd mmm yy

| Temperature | | 1 | l . | | _ | | _ |
|-------------|-----------|--------|-------|------|-----|---|------|
| 18.44 | _ 2 Q/min | 14/min | 1_ 1_ | 0152 | 188 | 0 | 94.0 |

| GENERAL EXAMINATION: | | | | | | | |
|----------------------|------|------|--|--|--|--|--|
| (check) | Nor. | Abn. | Provide details of abnormalities | | | | |
| Head/Neck | ~ | | | | | | |
| EENT | | / | mouth - a small 1-2 mm mucular red patchis on hard patches | | | | |
| Chest, lungs | | V | from 2nd remnestin + language clears to cough | | | | |
| Heart | r | | , | | | | |
| Abdomen | V | | | | | | |
| Genitalia | | | Not Done | | | | |
| Rectal | | | Not Done | | | | |
| Extremities | V | | patch over @ 3rd MCP point from nemote trauma | | | | |
| Skin | | / | patch over @ 3rd MCP joint from nemote houma orge scar Wiewer thorax from remote trauma while we would be the cafe au last spot 4000 LUO | | | | |
| Neurologic | V | | | | | | |

CHEST X-RAY Date 15/Jul/86

| NORMAL | Х | ABNORMAL | Describe abnormalities: | |
|--------|---|----------|-------------------------|--|
| | | | | |

Examiner

Dr. E.G. Felly

print name

20

10

Ņ

8

8

8

33

XX

33

X

page 4

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | |
|---------------------|------------|------|-----------------------------------|--|
| LIETMAN | W-E | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

MEDICATION RECORD

STUDY: WR6026

| Study | Date | Start drug admin- | End drug admin- | Route | Bottle |
|-------|---------|--------------------|--------------------|-------|--------|
| Day | ddmmmyy | istration (0-2400) | istration (0-2400) | | I.D. # |
| 3 | 2170186 | 0825 | N.A | PO | |

* Below Assay Sensitivity

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methems- globin _¥ |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|----------|--------------------------|
| | 0 | aJul86 | 0757 | 0757 | * | |
| | 0.25 | 41Jul 86 | 0840 | 0840 | * | |
| | 0.50 | 21 Jul 86 | 0855 | 0855 | * | |
| | 0.75 | al Jul 86 | 0910 | 0910 | * | |
| | 1.0 | 21Jul 86 | 0955 | 0925 | 7.47 | |
| | 1.25 | 21Ju186 | 0940 | 0940 | 13.0 | |
| | 1.50 | 2Nul 86 | 0955 | 0955 | 14.7 | |
| | 2.0 | 21Jul 86 | 1025 | 1025 | P.8G | |
| | 2.5 | 21Ju/86 | 1055 | 1055 | 46.8 | |
| | 3.0 | 21Ju/86 | 1155 | 1125 | 52.5 | _ |
| | 3.5 | 21 Jul 86 | 1155 | 1155 | 36.6 | |
| | 4.0 | 21 Jul 86 | 1225 | 1225 | 48.3 | |
| | 5.0 | 21 Jul 86 | 1325 | 1325 | 466 | |
| | 6.0 | 21Ju/86 | 14×5 | 1425 | 39.2 | |
| | 8.0 | 21Ju/86 | 1625 | 1625 | 33.6 | |
| | 10.0 | 21Jul 86 | 1825 | 1326 | 36.2 | |
| | 12.0 | 21 Jul 86 | 2025 114 | 500E8 | 26.9 | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WRED26 | |
|---------------------|------------|------|-----------------------------------|--|
| LIETMAN | WIF | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

MEDICATION RECORD

STUDY: WR6026

8

X

8

| Study Day | Date ddmmmyy | Start drug admin- istration (0-2400) | End drug | admin- (0-2400) | Route | Bottle I.D. # |
|--------------|--------------|---|----------|--------------------|-------|------------------|
| 3 | 21Jul 86 | 0825 | N.A. | | PO | |

* Below Assay Sensitivity

DOSAGE (total) 60 mg.

PLASMA CONCENTRATIONS

page 2

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo g lobin % |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|-------------|------------------------------------|
| | 24.0 | 22Jul 86 | 0825 | 0825 | 15.0 | |
| | 36.0 | 22Jul86 | 2025 | 2027 | 12.0 | |
| | 48.0 | 23Jul 86 | 0825 | 0835 | * | |
| | 60.0 | 23 Jul 86 | 2025 | 2147 | - *k | |
| | 72.0 | 24Jul 86 | 0825 | 0828 | * | |
| | 84.0 | 24Jul86 | 2025 | 2032 | * | |
| | 96.0 | 25Jul86 | 0825 | 0835 | * | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | 115 | | | |

page 6

| INVESTIGATOR'S NAME | PT INITIAL PT # | COMPOUND: WR6026 |
|---------------------|-------------------|-----------------------------------|
| LIETMAN | W - F F M L 03 | PROTOCOL:DAMD 17-85-C-5133- 03 |

URINE CONCENTRATIONS

STUDY: WR6026

| Study | Date | Start drug admin- | End drug admin- | Route | Bottle |
|-------|----------|--------------------|--------------------|-------|--------|
| Day | ddmmmyy | istration (0-2400) | istration (0-2400) | | I.D. # |
| 3 | 21Jul 86 | 0825 | NA | PO | |

DOSAGE (total) 60 mg

URINE CONCENTRATIONS

| Sample | Sched | | Start Co | ollection | End Col | lection | Total Volume | FURCO263 |
|-------------|------------------|-----|-----------|-----------|-----------|---------|-----------------|-----------|
| No. | Collecti (hou | | ddmmmyy | 0-2400 | ddmmmyy | 0-2400 | (ml) | [WR6026] |
| U01 | -24 TO | -12 | 20 Tul 86 | 0800 | 20 Jul 86 | 2005 | 1270 | |
| nos | -12 TO | 0 | 201186 | 2005 | 21 Jul 86 | 0800 | 1045 | 0 |
| U03 | о то | 12 | 2771186 | 0800 | 21 Jul86 | 2000 | 1390 | 0.17097 |
| U04 | 12 TO | 24 | 21Jul 86 | 2000 | 22 Jul 86 | 0800 | 1855 | 0.10944 |
| U0 5 | 24 TO | 36 | 22 Jul 86 | 0800 | 22 Jul 86 | 2000 | 1328 | 0.065073 |
| noe | 36 TO | 48 | 22 Jul 86 | 2000 | 23 Jul 86 | 0805 | 3/30 | O.0418FG |
| U07 | 48 TO | 60 | 23 Jul 86 | 0805 | 23 Jul 86 | 2100 | 1695 | D. 018645 |
| uos | 60 TO | 72 | 23 Jul 86 | 2100 | 24 JUl 86 | 0835 | 1805 | 0.013635 |
| 009 | 72 TO | 84 | 24Jul 86 | 0835 | 29 Jul 86 | 2000 | 1650 | 0.0331 |
| U10 | 84 TO | 96 | 24 Jul 86 | 2000 | 25 Jul 86 | 0800 | 1720 | 0 |
| | | | | | | | | |
| | | | , | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |

NI SONO CONTRACTO CONTRACTOR CONTRACTO CONTRACTOR C

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | W - F | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 |

1

8

X

1

Š

88

8

X

888

7

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| | Screen - | Predrug 1 | Predrug 2 | Postdrug 4 5 | | Day |
|------------------------|--------------|--------------|--------------|---------------------|-----------------------------|------|
| TEST:NORMAL | | | | 22Jul 86 ddmmmyy | 23 <u>Ju</u> 186 ddmmmyy | Date |
| NA:135-148 MEQ/L | 142 | 140 | 142 | 143 | 142 | |
| K:3.5-5.0 MEQ/L | 4.9 | 4.6 | 3.7 | 4.3 | 4.6 | |
| CL:96-109 MEQ/L | 108 | 109 | 108 | 11.1 | 108 | |
| CO2:24-30 MEQ/L | 26 | 15 | 20 | 19 | 19 | |
| SUN:12-25 MG/DL | 20 | 15 | 15 | 14 | 15 | |
| CREAT:0.4-1.5 MG/DL | 1.2 | 0.7 | 1.0 | 0.9 | 0.9 | |
| GLU:70-115 MG/DL | 98 | 70 | 69 | 90 | 86 | |
| T.BILI:0.3-1.2MG/DL | 0.5 | 0.2 | 0.6 | 0.6 | 0.5 | |
| D. BILI:0. 1-0. 4MG/DL | 0.1 | 0.1 | 0.1 | 0,1 | 0.0 | |
| CA:9.0-11.0 MG/DL | 10.0 | 9.6 | 9.5 | 9.7 | 9.9 | |
| PO4:3.0-4.5 MG/DL | 3.7 | 3.6 | 2.8 | 3.5 | 3.5 | |
| URIC A:4.2-8.8MG/DL | 9.4 | 7.4 | 6.5 | 7.5 | 7.4 | |
| T. PROT:6.0-8.5G/DL | 7.8 | 7.1 | 6.7 | 7.2 | 7.5 | |
| ALB.:3.2-5.3 G/DL | 4.8 | 4.4 | 4.2 | N.D. | 4.7 | |
| AST:0-35 IU/L | 20 | 14 | 23 | 3 | 27 | |
| ALT:0-30 IU/L | 7 | 7 | 10 | D | 16 | |
| ALK PHOS:0-95 IU/L | <i>5</i> 3 | 45 | 48 | 46 | 49 | } |
| CHOL:151-268 MG/DL | 190 | 177 | 184 | 183 | 197 | |
| LDH:0-200 IU/L | 15 / | 161 | 147 | 190 | 170 | |
| CPK:0-160 U/L(male) | a 8 5 | 338 | <i>á13</i> | 200 | 187 | |
| TG:20-190 MG/DL | 164 | - 60cgs | 62 | 62 | 87 | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | V-F FML | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 |

CONTROL OF THE CONTRO

H

3

8

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Postdrug 7 follow=w) | | | | | |
|----------------------|------------------------|------------|---------|---------|---------|------|
| TEST: NORMAL | 25Jull6 | OS ALLA SA | ddmmmyy | ddmmmyy | ddmmmyy | Date |
| NA:135-148 MEQ/L | 141 | ND | | | | |
| K:3.5-5.0 MEQ/L | 3.8 | ND | | | | |
| CL:96-109 MEQ/L | 106 | ND | | | | |
| CO2:24-30 MEQ/L | 22 | ND | | | | |
| SUN:12-25 MG/DL | 15 | 10 | | | | |
| CREAT: 0.4-1.5 MG/DL | 1.0 | 1.1 | | | | |
| GLU:70-115 MG/DL | 76 | 70 | | | | |
| T.BILI:0.3-1.2MG/DL | 0.7 | 0.3 | | | | |
| D. BILI:0.1-0.4MG/DL | ND- | 0.1 | | | | |
| CA:9.0-11.0 MG/DL | 9.8 | 9.7 | | | | |
| PD4:3.0-4.5 MG/DL | 3.1 | 2,2 | | | | |
| URIC A:4.2-8.8MG/DL | 6.8 | 8.6 | | | | |
| T. PROT:6.0-8.5G/DL | 7.7 | ND | | | | |
| ALB.:3.2-5.3 G/DL | 4.8 | 4.3 | | | | |
| AST:0-35 IU/L | 41 | 22 | | | | |
| ALT:0-30 IU/L | 21 | 11 | | | | |
| ALK PHOS:0-95 IU/L | 46 | 47 | | | | |
| CHOL:151-268 MG/DL | 203 | 166 | | | | |
| LDH:0-200 IU/L | 201 | ND | | | | |
| CPK:0-160 U/L(male) | 193 | ND | | | | |
| TG:20-190 MG/DL | 60 | NP | | | | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | W-F FML | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUESLaboratory: JOHNS HOPKINS HOSPITAL

| | • | Screen - | Predrug 1 | Predrug 3 | Post | tdrug 5 | Day |
|----------|------------|--------------------|--------------|---------------------|------------|------------|------|
| TEST | NORMAL | 08Jul86 Www.mpb | | 21Jul 86 ddmmmyy | | | Date |
| MBC | 4500-11000 | 6200 | 6406 | 5700 | 5000 | 4400 | |
| RBC | 4.50-5.90 | 4.74 | 4.21 | 4.28 | 4.37 | 4.58 | |
| Ндъ | 13.9-16.3 | 15,3 | 13.5 | 14.4 | 14.6 | 13.8 | |
| PCV | 41.0-53.0 | 44.9 | 40.1 | 41.8 | 42.6 | 44.2 | |
| Plt | 150-350 | 331 | 323 | 298 | 330 | 309 | |
| Bands | 2-6% | 3 | 7 | 3 | 2 | 1 | |
| Polys | 31-76% | 38 | 38 | 35 | 48 | 3 3 | |
| Eos | 1-4% | 5 | 3 | 3 | 1 | 2 | |
| Bas | | 0 | 0 | 2 | 1 | 0 | |
| Lymphs | 24-44% | 45 | 46 | 47 | 4 3 | 53 | } |
| Atyp Lym | | 2 | 0 | 1 | 0 | 0 | |
| Monos | 2-11% | 7 | 6 | 9 | 5 | 11 | |
| Other | | 0 | 0 | 0 | 6 | 6 | |
| | | | | | | | |
| Methem: | | 1:0 | | 0.1/0.0 | 0./ | 0.4 | |
| G-6-PD | 7.4-9.4 | 7.0 | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | |] |
| | | | | | | | |

Ľ

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | |
|---------------------|----------------|------|-----------------------------------|--|
| LIETMAN | W - F F M L | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

HEMATOLOBY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug

7

8

が

8

Day

Date

| TEST | NORMAL | 25 Jul 86 ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy |]1 |
|----------|------------|-------------------|---------|---------|---------|---------|----|
| MBC | 4500-11000 | 7300 | | | | | |
| RBC | 4.50-5.90 | 4.57 | | | | | |
| Ндь | 13.9-16.3 | 14.9 | | | | | |
| PCV | 41.0-53.0 | 43.3 | | | | | |
| Plt | 150-350 | 304 | | | | | |
| Bands | 2-6% | 4 | | | | | |
| Polys | 31-76% | 50 | | | | | |
| Eos | 1-4% | 2 | | | | | |
| Bas | | 1 | | | | | |
| Lymphs | 24-44% | 33 | | | | | |
| Atyp Lym | | 1 | | | | | |
| Monos | 2-11% | 9 | | | | | |
| Other | | 0 | | | | | |
| | | | | 1 | | | |
| methem. | 0.0 - 1.5 | 0.3 | | | | | |
| G-6-PD | 7.4-9.4 | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | W - F F M L | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 |

URINALYSIS VALUES

8

8

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Screen Predrug - 1 | | 4- | Postdrug 4 5 7 | | |
|-----------|-----------------------|-----------------------------|---------|-------------------|--------------------|--|
| | LSJul86 ddmmmyy | 19 <u>Jul</u> %6 ddmmmyy | qqwwwyy | ddmmmyy | 25Ju/86 ddmmmyy | |
| 5p.6r. | 1.022 | 1.020 | | | 1.018 | |
| pH · | 6.0 | 6.0 | | | 6.0 | |
| Protein | Neg | Neg | | | Nea | |
| Glucose | Neg | Neg | | | Nea | |
| Ketones | Neg | Neg | | | Nea | |
| Bili. | Neg | Neg | | | Neg | |
| Occ. Bld. | Neg | Neg | | | Neg | |
| Cast/lpf | 0 | L' [] | | | 0 | |
| WBC/hpf | 0 | | | | 0 | |
| RBC/hpf | 0 | | | | 0 | |
| Epi./hpf | 0-1 | | | | 0 | |
| Crys/hpf | 0 | , | | | 0 | |
| Bact/hpf | 0 | | | | b | |

ELECTROCARDIOGRAM

| Date ddmmmyy | Time 0-2400 | NORMAL check | ABNORMAL check | Describe abnormalities |
|-----------------------|----------------|-----------------|-------------------|------------------------|
| 21 Jul86 | 0705 | / | | |
| 21 Jul86 | 1145 | V | | |
| 22 Jul 86 | 09/2 | / | | |
| 237486 | 0954 | | | |
| 25Jul Pb 15 Jul 86 | | / | | |
| 15 Jul 86 | 1401 | | | 121 |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | |
|---------------------|----------------|------|-----------------------------------|--|
| LIETMAN | W - F F M L | 03 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

| rational and the state of the state of the | 18.456/8319/856/856/856/856/856/856/856/85 | ŎĸŶĸŎijŶĸŎĸĸŶĸĬĠĊŶĸŎĬĸŎĸŖŶĊĸŎŨĸĠĸŶŎĊĠ | `\$``\$\\$\`\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\\$\ | <u>₹</u> \$}\$\\$\$\\$\$\\$\$\\$\$\\$\$\\$\$\\$\$\\$\$\\$\$\\$\$\\$\$\\$\$\ | en e | ndy anadronomonomonomo |
|--|--|---------------------------------------|---|---|--|--|
| | | | | | page 12 | 2 |
| 22 | INVESTIGATOR'S | NAME PT INI | TIAL PT # | COMPOUND | :WR6026 | |
| 83 ∑ | LIETMAN | <u> </u> | E 03 | PROTOCOL | :DAMD 17-85 | 5-C-5133- 03 |
| Ti. | | ADVER | SE EXPERIE | NCES | | |
| & | | check if non | e occurred | | | |
| g | ADVERSE EVENT | ONSET (Date/Time) | CESSATION Date/Time | | REL'N TO TEST DRUG | ACTION (Check) |
| \$ \$ | * Malel elevation | 25 Jul 82 dd mmm yy | 08 Aug PG dd mmm yy | Mild | DEF. | None . |
| % | | (0-2400) | (0-2400) | Mod Mod | PROB. | Treatment |
| | | | | □ Sev | DEF. NOT | Stop test drug |
| ** | * | dd mmm yy | dd mmm yy | ☐ Mild | DEF. | None |
| \$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\ | | (0-2400) | (0-2400) | Mod | PROB. | Treatment |
| ************************************** | | | | Sev | DEF. NOT | Stop test drug |
| * | COMMENTS (Indic | ate # and eve | ent) | | | ······································ |
| K | | | | | | |
| ** | | | | | | |
| <u>F</u> | | | 122 | | | |
| | | KOKO40K0K0K0K0K0 | NESCHOLIGIES SE | 2 5 250262626 | 20202020202020 | \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\ |

| | | | | page 13 | | |
|--|-----------------------|-------------------------|----------------|----------------------------|--|--|
| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: | NR6026 | | |
| LIETMAN , | <u>W - F</u> F M L | 03 | PROTOCOL : DAI | MD 17-85-C-5133- 03 | | |
| (То | - | UTCOME ed for | all subject: | 5) | | |
| Protocol c | ompleted | | | <u>157</u> u/86 ddmmmyy | | |
| Premature | termination | of pr | otocol | ddmmmyy | | |
| REASON FOR PREMATURE TERMINATION (Check appropriate category) Adverse Experience Died During Study Failure To Return For Follow-up Did Not Cooperate Protocol Violation Entry Violation Intercurrent Illness Administrative/Other If terminated early, explain briefly: | | | | | | |
| | | | | | | |

A CONTRACTOR OF THE PROPERTY O

80

V.

8

N

**

X

30

8

page 14

| INVESTIGATOR'S NAME PT INITIAL PT # COMPOUND: WR6026 W - F PROTOCOL: DAMD 17-85-C-51 | | | | | | | | | |
|---|----|------------------|-------------|-------|----|-----------|--|--|--|
| LIETMAN | 03 | -85-C-5133 03 | | | | | | | |
| CONCOMITANT MEDICATIONS | | | | | | | | | |
| DRUG NAME TOTAL DATE DATE DRUG NAME DAILY DOSE STARTED STOPP | | | | | | | | | |
| heparin flush | | 100-10 | 00 unit | artul | 86 | 25 1. 1 K | | | |
| | | | | | | | | | |
| | | ···· | | L | | | | | |
| COMMENTS (please date and sign all comments) | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |

B32 538

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | L M H F M L | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 |

PROTOCOL

| Study Day | Date ddmmmyy | Procedures |
|--------------|--------------|------------------------|
| | 14Jul 86 | Screening laboratory |
| | 15Jul86 | History, Physical Exam |
| 0 | 38/JUEP1 | Admission |
| 3 | 21Ju186 | Drug Administration |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 14 pages, for subject # 04.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

freut Willy M.D.

nvestigator's signature

dd/mmm/y)

| INVESTIGATOR'S NAME | PT INITIAL | # TG | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | F M L | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICAL HISTORY

Date of evaluation \$5/Ju| 86

Examiner

Breat Willy 14

Date of birth

17/MAY/62 dd mmm yy print name

Age

32

Š

Š

\$32

24 yrs

Sex

M

Race

B

| Allergy Tobacco Use Alcohol Use Recreational Drug Use Medications past 2 weeks Experimental Drug Exposure Blood or plasma donor Prior Surgery Eye, ear, nose, throat Endocrine(diabetes, thyroid) C-V (heart murmur, HBP) Pulmonary (cough, asthma) Hepatitis, gastro-intestinal | |
|---|-------|
| Alcohol Use Recreational Drug Use Medications past 2 weeks Experimental Drug Exposure Blood or plasma donor Prior Surgery Eye, ear, nose, throat Endocrine(diabetes, thyroid) C-V (heart murmur, HBP) Pulmonary (cough, asthma) | |
| Alcohol Use Recreational Drug Use Medications past 2 weeks Experimental Drug Exposure Blood or plasma donor Prior Surgery Eye, ear, nose, throat Endocrine(diabetes, thyroid) C-V (heart murmur, HBP) Pulmonary (cough, asthma) | |
| Recreational Drug Use Medications past 2 weeks ? Darvon ~ 2Dago | |
| Medications past 2 weeks Experimental Drug Exposure Blood or plasma donor Prior Surgery Eye, ear, nose, throat Endocrine(diabetes, thyroid) C-V (heart murmur, HBP) Pulmonary (cough, asthma) | 70 |
| Experimental Drug Exposure Blood or plasma donor Prior Surgery Eye, ear, nose, throat Endocrine(diabetes, thyroid) C-V (heart murmur, HBP) Pulmonary (cough, asthma) | |
| Blood or plasma donor Prior Surgery Eye, ear, nose, throat Endocrine(diabetes, thyroid) C-V (heart murmur, HBP) Pulmonary (cough, asthma) | |
| Eye, ear, nose, throat Endocrine(diabetes, thyroid) C-V (heart murmur, HBP) Pulmonary (cough, asthma) | |
| Eye, ear, nose, throat Endocrine(diabetes, thyroid) C-V (heart murmur, HBP) Pulmonary (cough, asthma) | LR ID |
| C-V (heart murmur, HBP) Pulmonary (cough, asthma) | 1 |
| Pulmonary (cough, asthma) | |
| | |
| Hepatitis, gastro-intestinal 🗸 | |
| | |
| Benito-urinary GC:n 1981, resolved of | PCN |
| Musculoskeletal | |
| Neuropsychiatric ✓ | |
| Other | |
| 126 | |

| INVESTIGATOR'S NAM | E PT INITIAL | PT # | COMPOUND: WR6026 | | |
|--------------------|--------------|------|-----------------------------------|--|--|
| LIETMAN | L M H | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 | | |

| | | i | Blood Pressure | L | L J |
|--------|----------|--------|----------------|--------|------|
| 36.4 c | _ 68/min | 12/min | 118162 | 16.5.0 | 54.5 |

| 8 | | | _ | | | | page 3 | | |
|--------------|--------------|--------|--------|----------------------------------|---------------------------|---------------------|-----------------|----------------|--|
| Cui | INVESTIGATOR | 'S NAM | 1E PI | INITIAL | PT # | COMPOUR | ND:WR6026 | | |
| | LIETMAN | |) F | = M H | 04 | PROTOCO | DL:DAMD 17-85- | -C-5133- 03 | |
| | | | | | N. EXAM 15/Ju dd mm | | | | |
| ý | Temperature | Puls | i e | Respir | Blood | Pressure | Height (cm) | Wt. (kg | |
| 3 | 36.4 c | _68, | min | 12/min | 1 1 2 | 3_1_6 2 | 16.5.0 | 54.5 | |
| C | | | | | | | | | |
| Ž. | | | | GENEI | RAL EXA | MINATIO | V: | | |
| 3 | (check) | Nor. | Abn. | Provide details of abnormalities | | | | | |
| - | Head/Neck | | 7 | Shottu | nodes | both ar | t. cervical are | nS | |
| Ž. | EENT | | > | · 1 | | TH'S | | | |
| | Chest, lungs | V | | | | | | | |
| Ĭ | Heart | ~ | | | | | | | |
| Š | Abdomen | / | | | | | | | |
| : | Genitalia | | | Not. | Done | | | | |
| 8 | Rectal | | | Not | | | | | |
| | Extremities | / | | | | | | | |
| 80 82 | Skin | | / | Scar Du | stiet fro | m sunger on back | 9 1785 178 | | |
| F M | Neurologic | V | | 17.20 | | | | | |

CHEST X-RAY
Date 15/JU/86

Ÿ.

7

8

| NORMAL | Y | ABNORMAL | Describe | abnormalities: |
|--------|---|----------|----------|----------------|
| | | | | |

Examiner

Dr. B.G. Petty Dry)

Dr. B.G. Petty

print name

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|--------------|------|-----------------------------|
| LIETMAN | <u>- M +</u> | 04 | PROTOCOL:DAMD 17-85-C-5133- |

MEDICATION RECORD

STUDY: WR6026

| Study | Date | Start drug admin- | End drug | admin- | Route | Bottle |
|-------|---------|--------------------|-----------|----------|-------|--------|
| Day | ddmmmyy | istration (0-2400) | istration | (0-2400) | | I.D. # |
| 3 | 21Jul86 | 0810 | NA | | PO | |

DOSAGE (total) 60 mg

* Below Assay Sensitivity

PLASMA CONCENTRATIONS

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo- globin |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|----------|--------------------|
| | 0 | 21Jul 86 | 0800 | 0800 | * | |
| | 0.25 | 2Nu186 | 0825 | 0825 | * | |
| | 0.50 | 21 Jul 86 | 0840 | 0840 | P.C1 | |
| | 0.75 | 21 Jul86 | 0855 | 0855 | 18.0 | |
| | 1.0 | 21 Jul 86 | 0910 | 0910 | 37.8 | |
| | 1.25 | 21 Jul 86 | 0925 | 0925 | 51.2 | |
| | 1.50 | 21 Jul 86 | 0940 | 0940 | 59.6 | |
| | 2.0 | 21 Jul 86 | 1010 | 1010 | D.C8 | |
| | 2.5 | 2171 86 | 1046 | 1040 | 85.6 | |
| | 3.0 | 2154186 | 1110 | 1110 | 68.6 | |
| | 3.5 | 2154186 | 1140 | 1140 | G8.3 | |
| | 4.0 | 21Jul 86 | 1210 | 1210 | 62.8 | |
| | 5.0 | 2154186 | 1310 | 1310 | 75.5 | |
| | 6.0 | 215486 | 1410 | 1410 | 52.4 | |
| | 8.0 | 21 Jul 86 | 1610 | 1613 | 36.8 | |
| | 10.0 | 2/Jul 86 | 1810 | 1810 | 37.2 | |
| | 12.0 | 21JW 86 | 20 10 128 | 2010 | 24.1 | |

555 557

2000 S

35

(A)

松

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | | |
|---------------------|------------|------|-----------------------------------|--|--|
| LIETMAN | F M H | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 | | |

MEDICATION RECORD

STUDY: WR6026

8

8

3

8

32

X

33

8

Ž.

| Study Day | Date ddmmmyy | Start drug admin- istration (0-2400) | End drug admin- istration (0-2400) | Route | Bottle I.D. # |
|--------------|--------------|---|---------------------------------------|-------|------------------|
| 3 | 21Jul 86 | 0810 | N.A. | PO | |

* Below Assay Sensitivity

DOSAGE (total) 60 mg

PLASMA CONCENTRATIONS

page 2

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemos globin |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|----------|--------------------|
| ! | 24.0 | 22 Jul 86 | 0810 | 0810 | 8.79 | |
| | 36.0 | 22Jul 86 | 2010 | 2010 | * | |
| | 48.0 | 23Jul 86 | 0810 | 0816 | * | |
| | 60.0 | 23Jul 86 | 2010 | 2/37 | * | |
| | 72.0 | 24Jul 86 | 0810 | 08/2 | * | |
| | 84.0 | 24Jul 86 | 2010 | 2015 | * | |
| ! | 96.0 | 25 Jul 86 | 0810 | 0810 | * | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | 129 | | | |

page 6

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | | | |
|---------------------|------------|------|-----------------------------------|--|--|--|
| LIETMAN | <u> </u> | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 | | | |

URINE CONCENTRATIONS

STUDY:WR6026

R

X

8

8

| Study | Date | Start drug admin- | End drug admin- | Route | Bottle |
|-------|----------|--------------------|--------------------|-------|--------|
| Day | ddmmmyy | istration (0-2400) | istration (0-2400) | | I.D. # |
| 3 | 21Jul 86 | 0810 | N.A. | PO | |

DOSAGE (total) _60mg___

URINE CONCENTRATIONS

| Sample Scheduled No. Collection Time | | Start Co | ollection | End Coli | lection | Total Volume | [WR6026] | | |
|--------------------------------------|--------------------|----------|-----------|-----------|---------|-----------------|----------|------|----------|
| | (hours) | | ddmmmyy | 0-2400 | ddmmmyy | 0-2400 | (ml) | my. | |
| U01 | -24 | TO | -12 | 20 Jul 86 | 0800 | 20 Jul 86 | 2000 | 755 | |
| nos | -12 | το | 0 | 20 Jul 86 | 2000 | 21 Jul 86 | 0800 | 735 | 0 |
| no3 | 0 | TO | 12 | 21Jul 86 | 0800 | 21 Jul 86 | 2000 | 510 | 0-101920 |
| U04 | 12 | TO | 24 | 21 Jul 86 | 2000 | 22 Jul 86 | 0800 | 595 | 0077053 |
| U05 | 24 | TO | 36 | 22 Jul 86 | 0800 | 22Jul 86 | 2000 | 432 | 0.021987 |
| U06 | 36 | TO | 48 | 22 Jul 86 | 2000 | 23 Jul 86 | 0800 | 745 | C.011287 |
| U07 | 48 | TO | 60 | 23 Jul 86 | 0800 | 23 Jul 86 | 2000 | 685 | 0.013104 |
| UOB | 60 | TO | 72 | 23 JU86 | 2000 | 29 Jul 86 | 0810 | 490 | 0.008100 |
| U09 | 72 | TO | 84 | 24 Jul 86 | 0810 | 24 Jul 86 | 2000 | 1050 | 0.01&92 |
| U10 | 84 | TO | 96 | 24 Jul 86 | 2000 | 25 Jul 86 | 0800 | 745 | 0 |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | -, -, - | | | | 130 | | | | , |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | F M H | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 |

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| , | Screen - | Predrug 1 | Predrug 2 | Postdrug 4 5 | | Day |
|------------------------|-------------|--------------|----------------------------|-----------------|--------------|------|
| TEST:NORMAL | | | 21 <u>Jul86</u> ddmmmyy | | | Date |
| NA:135-148 MEQ/L | 140 | 144 | 142 | 138 | 139 | |
| K:3.5-5.0 MEQ/L | 45 | 4.3 | 4.4 | 5.0 | 4.2 | |
| CL:96-109 MEQ/L | 106 | 104 | 106 | 107 | 108 | |
| CD2:24-30 MEQ/L | 28 | 24 | 28 | 20 | 20 | |
| SUN:12-25 MG/DL | 12 | 12 | 9 | 10 | 13 | |
| CREAT:0.4-1.5 MG/DL | 0.7 | 1.0 | 1.0 | 1.0 | 0.9 | |
| GLU:70-115 MG/DL | 76 | 72 | 70 | 84 | 76 | |
| T.BILI:0.3-1.2MG/DL | 0.6 | 6.5 | 0.6 | 0.9 | 0.7 | |
| D. BILI:0. 1-0. 4MG/DL | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | |
| CA:9.0-11.0 MG/DL | 10.3 | 9.8 | 9.9 | 10.2 | 9.5 | |
| PD4:3.0-4.5 MG/DL | 4.7 | 5.2 | 3.7 | 3.5 | <i>3</i> . 8 | |
| URIC A:4.2-8.8MG/DL | 7.0 | 1.3 | 5.8 | 6.5 | 6.9 | |
| T. PROT:6.0-8.5G/DL | 7.4 | 4.6 | 7.4 | 6.9 | 6.7 | |
| ALB.:3.2-5.3 G/DL | 4.6 | 4.6 | 4.3 | Da | 4.3 | |
| AST:0-35 IU/L | 25 | 27 | 21 | 21 | 13 | |
| ALT:0-30 IU/L | 12 | 13 | 15 | 15 | 7 | |
| ALK PHOS:0-95 IU/L | 66 | 61 | 57 | 62 | 57 | |
| CHOL:151-268 MG/DL | 207 | 220 | 214 | 173 | 170 | |
| LDH:0-200 IU/L | 189 | ND | 168 | ND | 166 | |
| CPK:0-160 U/L(male) | 379 | ND | 278 | 169 | 169 | |
| TG:20-190 MG/DL | 50 | 9243 | 92 | 79 | 82 | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | | |
|---------------------|------------|------|-----------------------------------|--|--|
| LIETMAN | L M H | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 | | |

8

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| TEST:NORMAL | 25Jul 96 | ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy | Date |
|-------------------------|--------------|------------|-------------|---------|---------|------|
| | | admining y | - Comming y | dummiyy | | |
| NA:135-148 MEQ/L | 143 | | | | | |
| K:3.5-5.0 MEQ/L | 4.2 | | | | ·_ · | |
| CL:96-109 MEQ/L | 105 | | | | | |
| CO2:24-30 MEQ/L | 25 | | | | | |
| SUN:12-25 MG/DL | 10 | | | | | |
| CREAT: 0.4-1.5 MG/DL | 1.0 | | | | | |
| GLU:70-115 MG/DL | 72 | | | | | |
| T.BILI:0.3-1.2MG/DL | 0.8 | | | | | |
| D. BILI: 0. 1-0. 4MG/DL | 0.1 | | | | | |
| CA:9.0-11.0 MG/DL | 9.5 | | | | | |
| PD4:3.0-4.5 MG/DL | 3.8 | | | | | |
| URIC A:4.2-8.8MG/DL | 7,1 | | | | | |
| T. PROT:6.0-8.5G/DL | 7.0 | | | | | |
| ALB.:3.2-5.3 G/DL | N.D. | | | | | |
| AST:0-35 IU/L | 18 | | | | | |
| ALT:0-30 IU/L | 14 | | | | | |
| ALK PHOS:0-95 IU/L | 60 | | | | | |
| CHOL:151-268 MG/DL | 181 | | | | · | |
| LDH:0-200 IU/L | 146 | | | | | |
| CPK:0-160 U/L(male) | 18E | | | | | |
| TG:20-190 MG/DL | 89 | | | | | |
| | | 132 | | | | • |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|---------------|------|-----------------------------------|
| LIETMAN | LM H F M L | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUESLaboratory: JOHNS HOPKINS HOSPITAL

| | • | Screen - | Predrug 1 | Predrug 3 | Post 4 | Postdrug 4 5 | |
|----------|------------|--------------------|--------------|---------------------|-----------|-----------------|------|
| TEST | NORMAL | 14Jul86 ddmmmyy | | 21Jul 86 ddmmmyy | | | Date |
| MBC | 4500-11000 | 7200 | 6700 | 6400 | 9500 | 8400 | |
| RBC | 4.50-5.90 | 4.73 | 4.87 | 4.70 | 4.49 | 4.61 |] |
| Hgb | 13.9-16.3 | 14.6 | 14.6 | 14.8 | 14.5 | 14.0 | } |
| PCV | 41.0-53.0 | 43.6 | 44.8 | 44.2 | 42.1 | 43.1 | |
| Plt | 150~350 | 308 | 341 | 295 | 271 | 260 | |
| Bands | 2-6% | 4 | | 0 | 3 | 5 | |
| Polys | 31-76% | 52 | 54 | 64 | 69 | 64 | |
| Eos | 1-4% | 3 | | 0 | 1 | O |] |
| Bas | | , | 0 | 0 | 0 | O | |
| Lymphs | 24-44% | 27 | 43 | 32 | á l | á4 | |
| Atyp Lym | | 0 | D | 0 | 0 | Ö | |
| Monos | 2-11% | /3 | 1 | 4 | 6 | 7 | |
| Other | | 0 | 0 | 0 | 0 | 0 | |
| | | | | | | | |
| Methem | 0.0-1.5 | 0.0 | | 0.0/0.1 | 0.0 | 0.3 | |
| G-6-PD | 7.4-9.4 | 8.3 | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | } |
| | | | | | | | } |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|--------------|------|-----------------------------------|
| LIETMAN | LM H FM L | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug

7

Day

25Jul 86 Date TEST NORMAL ddmmmyy ddmmmyy ddmmmyy ddmmmyy ddmmmyy 4500-11000 WBC 6800 RBC 4.50-5.90 4.66 13.9-16.3 14.1 Hgb PCV 41.0-53.0 43.2 Plt 150-350 279 2-6% Bands 2 31-76% Polys 51 Eos 1-4% Bas 24-44% Lymphs 39 Atyp Lym 0 2-11% 6 Monos Other 0 nethem. 0.0 - 1-5 0.0 G-6-PD 7.4~9.4

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WREOSE |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | LMH FML | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 |

URINALYSIS VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| | Screen Predrug | | Postdrug | | |
|----------|----------------|----------|----------|----------|-----------|
| | - | 1 | 4 | 5 | 7 |
| | 15Ju186 | 19 Ju/86 | | | 25 Jul 86 |
| | qqwwwyy | ddmmmyy | aquuumyy | ddmmmyy | ddmmmyy |
| Sp.Gr. | 1.029 | 1.017 | 1 | 1 | 1.025 |
| pH · | 6.0 | 6.5 | | | 6.5 |
| Protein | Neg | Neg | | | Neg |
| Glucose | Neg | Weg | | | Neg |
| Ketones | Neg | Neg | | 1 | Nea |
| Bili. | Neg | Neg | | | Neg |
| ūcc.Bld. | Nes | Neg | | | Nea |
| Cast/lpf | 0) | ND | | | 0 |
| WBC/hpf | U | | | | 0 |
| REC/hpf | c | | | | 0 |
| Epi./hpf | C | | | | 0 |
| Crys/hpf | 0 | | | | 0 |
| Bact/hpf | 0 | | | <u> </u> | 0 |

ELECTROCARDIOGRAM

| Date ddmmreyy | Time 0-2400 | NORMAL check | ABNORMAL check | Describe abnormalities Short Printernal Uware |
|------------------|----------------|-----------------|-------------------|--|
| 21 Jul 86 | 0650 | · | V | Scot PR interval 4 waves |
| 21 Jul 86 | 1226 | | 1 | No significant change |
| 22JU86 | , , , , | | V | Aleno Twair diange in TI sheet PR |
| 23Jul 86 | l | | V | In significant diange |
| 25Jul 86 | 0826 | | | Twom both to riginal oppose were in III, dent P. |
| 15 Tel 86 | | — | | 135 |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|-------------|------|-----------------------------------|
| LIETMAN | LMH_ FML | 04 | PROTOCOL:DAMD 17-85-C-5133- 03 |

ADVERSE EXPERIENCES

check if none occurred

| abla | \neg |
|------------|---------------|
| 1 > | 71 |
| (/ | \mathcal{N} |
| ᢞ | ~ |

| | | | <u>/ </u> | | |
|------------------|----------------------|------------------------|--|-----------------------|-------------------|
| ADVERSE EVENT | ONSET (Date/Time) | CESSATION Date/Time | | REL'N TO TEST DRUG | ACTION (Check) |
| * | dd mmm yy | dd mmm yy | ☐ Mild | DEF. | None |
| | (0-2400) | (0-2400) | ☐ Mod | PROB. | Treatment |
| | | | □ Sev | DEF. NOT | Stop test drug |
| * | dd mmm yy | dd mmm yy | ☐ Mild | DEF. | None |
| | (0-2400) | (0-2400) | Mod | PROB. POSS. | Treatment |
| | | | Sev | DEF. NOT | Stop test drug |

COMMENTS (Indicate # and event)

teres " Receivery Vicationist Despitation Described

| | | | | | pa | ge 13 | |
|---------------------------------------|--|------------|--------|-----------|------|---------------------------------|--|
| INVESTI | GATOR'S NAME | PT INITIAL | PT # | COMPOUND: | WR60 | 26 | |
| LIET | LIETMAN . $\frac{L}{F} \frac{M}{M} \frac{H}{L} = 04$ PROTOCOL: DAMD | | | | AMD | 17-85-C-5133- 03 | |
| | OUTCOME (To be completed for all subjects) | | | | | | |
| \boxtimes | Protocol co | ompleted | | | | 25J ₁ 186 ddmmmyy | |
| | Premature t | ermination | of pro | otocol | | ddmmmyy | |
| | Died During Study Failure To Return For Follow-up Did Not Cooperate Protocol Violation Entry Violation Intercurrent Illness | | | | | | |
| If terminated early, explain briefly: | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |

enter that have been parties that the property of the property

X

| INVESTIGATOR'S NAME | PT INITIA | _1 | ND: WR6026 | | | |
|--|-------------|---------------------|-----------------|--|--|--|
| LIETMAN | L M H | 04 PROTOC | OL:DAMD 17 | -85-C-5133 03 | | |
| CONCOMITANT MEDICATIONS | | | | | | |
| DRUG NAME | | TOTAL DAILY DOSE | DATE STARTED | DATE STOPPED | | |
| heparin | | 100-1000 units | 21 Jul 86 | 24Jul 86 | | |
| | | | | | | |
| | | | | | | |
| COMMENTS (please date and sign all comments) | | | | | | |
| | | | | | | |
| L | | | - | ا استان میں میں استان | | |

10

8

X

X

X

122

8

page 1

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | D L A | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 |

PROTOCOL

| Study Day | Date ddmmmyy | Procedures |
|--------------|--------------|------------------------|
| | 14 Jul 86 | Screening laboratory |
| | 15Jul 86 | History, Physical Exam |
| 0 | 19Ju186 | Admission |
| 3 | 2474/86 | Drug Administration |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 14 pages, for subject 40.5
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Men States M.D.

nvestigator's signature

16/ Dec/ 80 dd/mmm/yy

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | DLA | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICAL HISTORY

Date of evaluation 15/Jul/86 dd mmm yy

Examiner

Date of birth

29, Sen, 58
dd mmm yy

Age

1

X

777

27 yrs

Sex

<u>M</u>_B

Race

| | No | Yes | Comments |
|-------------------------------|----------|----------|-------------------------------|
| Allergy | V | | |
| Tobacco Use | | ~ | 1/2 PPD |
| Alcohol Use | √ | | Quit 1/86 |
| Recreational Drug Use | | ./ | lary : 136 (MJ), 140 drup sys |
| Medications past 2 weeks | V | | |
| Experimental Drug Exposure | | √ | ourds cettriaxore |
| Blood or plasma donor | ~ | | Chalilical herne repair |
| Prior Sur ge ry | | V | Umbilical herne recair |
| Eye, ear, nose, throat | V | | |
| Endocrine (diabetes, thyroid) | ~ | | |
| C-V (heart murmur, HBP) | V | | |
| Pulmonary (cough, asthma) | ~ | | |
| Hepatitis, gastro-intestinal | V | | |
| Genito-urinary | / | | |
| Musculoskeletal | V | | |
| Neuropsychiatric | V | | |
| Other | • | | |
| | | | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | D L H F M L | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 |

PHYSICAL EXAMINATION Date 15/15/15/ dd mmm yy

| ı | Temperature | | | | | _ | 1 |
|---|---------------|---------|--------|----|------|-------|-------|
| | <u>36.7</u> t | _72/min | /Q/min | 10 | 8,64 | 195.0 | 105.0 |

CONTRACT CONTRACTOR NOTICE AND SOUND ASSESSMENT ASSESSM

E

Ž,

 Σ

₹ \$

7

33

,

8

W

50000

BSS > 2 20 MED ACCOUNT FOR SECTION

| | GENERAL EXAMINATION: | | | | | | | |
|--------------|----------------------|------|---|--|--|--|--|--|
| (check) | Nor. | Abn. | Provide details of abnormalities | | | | | |
| Head/Neck | 1/ | | 1x0.5cm node Preck unchanged x your occiden | | | | | |
| EENT | | L' | ears cureminouse librat | | | | | |
| Chest, lungs | ~ | | | | | | | |
| Heart | ٧. | | | | | | | |
| Abdomen | L-1 | | | | | | | |
| Genitalia | | | \mathcal{N}^{\prime} | | | | | |
| Rectal | | | N | | | | | |
| Extremities | U | | | | | | | |
| Skin | | ·/ | Scar O heels, appealock, un bilirux | | | | | |
| Neurologic | V | | | | | | | |

CHEST X-RAY Date 15/Je1/86

| NORMAL | X | ABNORMAL | Describe | abnormalities: |
|--------|---|----------|----------|----------------|
| | _ | | | |

Dr. B.G. Petty

print name

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | DLA | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICATION RECORD

STUDY: WR6026

| Study | Date | Start drug admin- | End drug admin- | Route | Bottle |
|-------|---------|--------------------|--------------------|-------|--------|
| Day | ddmmmyy | istration (0-2400) | istration (0-2400) | | I.D. # |
| 3 | 215,186 | 0805 | N.A. | PO | |

* Below Assay Sensitivity

DI AGMA CONCENTRATIONS

DOSAGE (total) 60 mg

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo- globin- * |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|--------------|--------------------------|
| | 0 | ENJW 86 | 0755 | 075 | * | |
| | 0.25 | 21 Jul 86 | 0820 | 0820 | * | |
| | 0.50 | 21/1/86 | 0835 | 0836 | * | |
| | 0.75 | 215486 | 0850 | 0850 | * | |
| | 1.0 | 215/186 | 0905 | 0905 | * | |
| | 1.25 | 21 Jul 86 | 0920 | 0920 | <u>0</u> | |
| | 1.50 | 21 Jul 86 | 0925 | 0935 | 19.4 | |
| | 2.0 | 21 Jul 86 | 1005 | 1005 | 94.9 | |
| | 2.5 | 215.186 | 1035 | 1035 | <i>33.</i> 5 | |
| | 3.0 | 21 Jul 86 | 1105 | 1107 | 48.9 | |
| | 3.5 | 21 Jul 86 | 1135 | 1135 | 53.3 | |
| | 4.0 | 2154186 | 120E | 1205 | G3.4 | |
| | 5.0 | 21 Jul 86 | 1305 | 1305 | 73.0 | |
| | 6.0 | 4Jul 86 | 1400 | 1405 | 30.1 | |
| | 8.0 | 21 Jul 86 | 1605 | 1608 | 47.2 | |
| | 10.0 | 21 Jul 86 | 1805 | 1706 | 38.3 | |
| | 12.0 | 21 Jul 86 | 2005 142 | 200E | 32.7 | |

*

Ä

*

23

XX

88

ŽŽ

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | DLA | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICATION RECORD

STUDY: WR6026

| Study Day | Date ddmmmyy | Start drug admin- istration (0-2400) | End drug admin- istration (0-2400) | Route | Bottle I.D. # |
|--------------|--------------|---|---------------------------------------|-------|------------------|
| 3 | 21Jul 86 | 0805 | N.A. | PO | |

* Below Assay Sensitivity

60 mg DOSAGE (total)

PLASMA CONCENTRATIONS

| page 2 | | | | | | |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|----------|--------------------------|
| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methems- globin -% |
| | 24.0 | 22Jul 86 | 0805 | 0805 | 13.0 | |
| | 36.0 | 22 Jul 86 | 2005 | 2005 | 7.67 | |
| | 48.0 | 23JW86 | 0805 | 0805 | * | |
| 1 | 60.0 | 23 Jul 86 | 080 2005 | 2142 | * | |
| | 72.0 | 24 JU86 | 0805 | 08/0 | * | |
| | 84.0 | 24 JW86 | 2005 | 2032 | * | |
| | 96.0 | 25 Jul 86 | 0805 | 0805 | * | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| - | | | | | | |
| | | | | | | |
| • | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | 143 | | | |

page 6

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | |
|---------------------|------------|------|-----------------------------------|--|
| LIETMAN | D L A | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

URINE CONCENTRATIONS

STUDY: WR6026

A

X

| | | Start drug admin- istration (0-2400) | | | Bottle I.D. # |
|---|--------|---|------|----|------------------|
| 3 | 215486 | 0805 | 0805 | PO | |

DOSAGE (total) 60 mg

| Sample | Scheduled Start Collection | | End Col | lection | Total Volume | FURCAGES | |
|--------|----------------------------|-----------|---------|-----------|-----------------|----------|----------|
| No. | Collection Time (hours) | ddmmmyy | 0-2400 | ddmmmyy | 0-2400 | (ml) | (WR6026) |
| U01 | -24 TO -12 | 20 Jul 86 | 0800 | 20 Jul 86 | 2000 | 1790 | |
| nos | -12 TO 0 | 20 Jul 86 | 2000 | 2/Jul 86 | 0800 | 815 | 0 |
| N03 | 0 TD 12 | 21 Jul86 | 0800 | 21 Jul 86 | 2000 | 485 | 0.63837 |
| U04 | 12 TO 24 | 21 Jul 86 | 2000 | 22 Jul 86 | 0800 | 1050 | 0.061018 |
| U05 | 24 TO 36 | 22 Jul 86 | 0800 | 22 Jul 86 | 2000 | 802 | 0.02177 |
| noe | 36 TO 48 | 22 Jul86 | 2000 | 23 Jul 86 | 0800 | 1160 | O.014318 |
| U07 | 48 TO 60 | 23 Jul 86 | 0800 | 23Jul 86 | 2000 | 1000 | 0.006443 |
| Bou | 60 TO 72 | 23 Jul 86 | 2000 | 29 Jul 86 | 0800 | 1070 | 0.003892 |
| U09 | 72 TO 84 | 29 Jul 86 | 0800 | 29 Jul 86 | 2000 | 750 | 0.003619 |
| U10 | 84 TO 96 | 4 Jul 86 | 2000 | 25 Jul 86 | 0800 | 1025 | 0.00057 |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | <u> </u> | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | D L A F M L | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 |

X

X) X

8

B

88

3

**

X

7

8

5

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Screen - | Predrug 1 | Predrug 2 | Post 4 | tdrug 5 | Day |
|------------------------|--------------------|-------------------|--------------|---------------------|------------|------|
| TEST:NORMAL | 14Ju/86 ddmmmyy | 19JW86 ddmmmyy | , | 22Tul 86 ddmmmyy | , | Date |
| NA:135-148 MEQ/L | 143 | 139 | 138 | 137 | 138 | |
| K:3.5-5.0 MEQ/L | 4.6 | 4.6 | 4.0 | 3.9 | 4.4 | |
| CL:96-109 MEQ/L | 105 | 103 | 101 | 104 | 106 | |
| CO2:24-30 MEQ/L | 24 | 20 | 24 | 21 | <i>á</i> / | |
| SUN:12-25 MG/DL | 13 | // | 10 | 13 | 15 | |
| CREAT:0.4-1.5 MG/DL | 1.1 | 1.1 | 0.9 | 1.1 | 1.0 | |
| GLU:70-115 MG/DL | 103 | 81 | 86 | 92 | 89 | |
| T.BILI:0.3-1.2MG/DL | 0.3 | 0.2 | 0.5 | 0.4 | 0.4 | |
| D. BILI:0. 1-0. 4MG/DL | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | |
| CA:9.0-11.0 MG/DL | mot done | 9.7 | 9.2 | 9.3 | 9. + | |
| PO4:3.0-4.5 MG/DL | 3.7 | 3.4 | 3.6 | 3.3 | 4.1 | |
| URIC A:4.2-8.8MG/DL | 4,9 | 4.9 | 5.4 | 5.7 | 6.2 | |
| T. PROT:6.0-8.5G/DL | 7.6 | 7.4 | 6.8 | 6.9 | 7./ | |
| ALB.:3.2-5.3 G/DL | 4.6 | 4.4 | 4.1 | 4.2 | 4.3 | |
| AST:0-35 IU/L | 28 | 28 | 30 | 27 | N.D. | |
| ALT:0-30 IU/L | 25 | 33 | 39 | 36 | 40 | |
| ALK PHOS:0-95 IU/L | 56 | 55 | 49 | 59 | 51 | |
| CHOL:151-268 MG/DL | 179 | 193 | 172 | 162 | 174 | |
| LDH:0-200 IU/L | 153 | 154 | /33 | 143 | 203 | |
| CPK:0-160 U/L(male) | 210 | 271 | 159 | 142 | 140 | |
| TG:20-190 MG/DL | 224 | ND | 157 | 139 | 147 | |

145

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | |
|---------------------|------------|------|-----------------------------------|--|
| LIETMAN | DA | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

X

N.

8

8

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Postdrug 7 | follow u | 0 | | | Day |
|----------------------|---------------------|------------------|---|---------|---------|------|
| TEST: NORMAL | 25Jul 06 ddmmmyy | 05 Aug 86 | | ddmmmyy | ddmmmyy | Date |
| NA:135-148 MEQ/L | 137 | ND | | | | |
| K:3.5-5.0 MEQ/L | 4.1 | ND | | , | | |
| CL:96-109 MEQ/L | 104 | ND | | | | |
| CO2:24-30 MEQ/L | 21 | ND | | | | |
| SUN:12-25 MG/DL | 13 | 10 | | | | |
| CREAT: 0.4-1.5 MG/DL | 1.1 | 1, 1 | | | | |
| GLU:70-115 MG/DL | 87 | 108 | | | | |
| T.BILI:0.3-1.2MG/DL | 0.4 | 0.5 | | | | |
| D. BILI:0.1-0.4MG/DL | 0.1 | 0,1 | | | | |
| CA:9.0-11.0 MG/DL | 10.2 | 10.0 | | | | |
| P04:3.0-4.5 MG/DL | 4.0 | 3.2 | | | | |
| URIC A:4.2-8.8MG/DL | 6.6 | channel mall. | | | | |
| T. PROT:6.0-8.5G/DL | 7.8 | 7.8 | | | | |
| ALB.:3.2-5.3 G/DL | 4.4 | 4.6 | | | | |
| AST:0-35 IU/L | 29 | 16 | | | | |
| ALT:0-30 IU/L | 44 | 19 | | | | |
| ALK PHOS:0-95 IU/L | 54 | 41 | | | | |
| CHOL:151-268 MG/DL | 191 | 175 | | | | |
| LDH:0-200 IU/L | 166 | 141 | | | | |
| CPK:0-160 U/L(male) | 169 | ND | | | | |
| TG:20-190 MG/DL | 160 | ND | | | | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | D L A F M L | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUESLaboratory: JOHNS HOPKINS HOSPITAL

| | • | Screen - | Predrug 1 | Predrug 3 | Post | drug 5 | Day |
|----------|------------|-------------|---------------|--------------|--------------------|---------------------|------|
| TEST | NORMAL | 14 Jul 86 | (| | 22Jul86 ddmmmyy | 23Jul 86 ddmmmyy | Date |
| WBC | 4500-11000 | 4700 | 6100 | 6300 | 7500 | 8600 | |
| RBC | 4.50-5.90 | 5.11 | 5.00 | 4.73 | 4.77 | 4.96 | |
| НдЬ | 13.9-16.3 | /4.8 | 13.8 | 13.9 | 14.1 | 13.9 | |
| PCV | 41.0-53.0 | 43.8 | 42.4 | 40.9 | 41.4 | 42.6 | |
| Plt | 150-350 | 207 | 232 | 233 | 283 | 267 | |
| Bands | 2-6% | 2 | 2998 | 2 | 4 | 4 | |
| Polys | 31-76% | 29 | 40 | 52 | 46 | 45 | |
| Eos | 1-4% | 29-1 | Į | l l | 2 | 1 | |
| Bas | | ABO | 0 | 0 | 0 | 0 | |
| Lymphs | 24-44% | 55 | 50 | 37 | 40 | 44 | |
| Atyp Lym | | 0 | 0 | l | D | 1 | |
| Monos | 2-11% | 13 | 7 | 7 | 8 | 4 | |
| Other | | | | | | 1 met | |
| | | | | | | | |
| Methemi | , | 1.4 | | 0.0 | 0.0 | 0.2 | |
| G-6-PD | 7.4-9.4 | 7.8 | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |

page 10

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | DAA | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug

7

25 Jul 86

STATES OF THE PROPERTY OF THE

Day

Date

TEST NORMAL ddmmmyy ddmmmyy ddmmmyy ddmmmyy ddmmmyy WBC 4500-11000 7800 4.50-5.90 RBC 4.97 Hgb 13.9-16.3 14.2 PCV 41.0-53.0 42.5 286 Plt 150-350 2-6% Bands 31-76% Polys 36 1-4% Eos Bas 0 24-44% Lymphs 54 Atyp Lym Monos 2-11% Other D methem 0.0 - 1.5 F.0 G-6-PD 7.4-9.4

| INVESTIGATOR'S NAME | PT INITIAL | # 19 | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | D L A F M L | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 |

URINALYSIS VALUES

JOHNS HOPKINS HOSPITAL Laboratory:

| · | Screen - | Predrug 1 | 4 | Postdrug 5 | 7 |
|----------|-------------------|--------------|---------|---------------|--------------------|
| | 15JUS6 ddmmmyy | , | ddmmmyy | ddmmmyy | 25J.186 ddmmmyy |
| Sp.Gr. | 1.026 | 1.024 | | | 1.027 |
| рН | 6.5 | 6.0 | | | 6.0 |
| Protein | Neg | Neg | | | heg |
| Glucose | Neg | Neg | | | hea |
| Ketones | Neg | Neg | | | Nea |
| Bili. | Neg | Neg | | | hea |
| Occ.Bld. | Neg | Neg | | | hea |
| Cast/lpf | o' | 12 | | | 0 |
| WBC/hpf | 0 | | | | 0-1/hpf |
| RBC/hpf | 10 | | | | 0 |
| Epi./hpf | 0 | | | | 0 |
| Crys/hpf | 0 | | | | 0 |
| Bact/hpf | 0 | | | | 0 |

ELECTROCARDIOGRAM

| 157 W 8 | | | incomplete right hundle branch & |
|-------------|----------|------|----------------------------------|
| 21 Jul 8 | | V | IRBBB |
| 22 Jul | 96 0905 | / | TRBBB |
| 23 Jul 8 | 6 1000 | V | IRBBB |
| 25 1 18 | 16 0817 | ~ | IRBBA |
| | <u>-</u> | | 149 |

N.

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | DLA | 05 | PROTOCOL:DAMD 17-85-C-5133- 03 |

ADVERSE EXPERIENCES

check if none occurred

| • | OVERSE EVENT | ONSET (Date/Time) | | B | REL'N TO TEST DRUG | ACTION (Check) |
|---|--------------------|----------------------|------------------------|--------------|------------------------|-------------------|
| # | Mildler | 19 Jul Rodd mmm yy | 05 Au Po | X Mild | DEF. | None |
| | Milder diestret | (0-2400) | (0-2400) | ☐ Mod | PROB. D POSS. | Treatment |
| | | | | □ Sev | DEP. NOT | Stop test drug |
| * | | 23 Jul Redd mmm yy | 25 Jul 86 dd mmm yy | Ď Mild | DEF. | None |
| / | Manualle, excepted | (0-2400) | (0-2400) | Mod | PROB. POSS. DEF. NOT | Treatment |
| | | | | Sev | UNKNOWN | Stop test drug |

COMMENTS (Indicate # and event)

| | | | | - | age 13 | | | | |
|--|---|----------------|------|---------------|---------------------|--|--|--|--|
| INVESTIGAT | OR'S NAME | PT INITIAL | PT # | COMPOUND: WR6 | 026 | | | | |
| LIETMAN | l , | D L A F M L | 05 | PROTOCOL:DAMD | 17-85-C-5133- 03 | | | | |
| OUTCOME (To be completed for all subjects) | | | | | | | | | |
| Protocol completed 25 Jul 80 ddmmmyy | | | | | | | | | |
| | Premature termination of protocol ddmmmyy | | | | | | | | |
| REASON FOR PREMATURE TERMINATION (Check appropriate category) Adverse Experience Died During Study Failure To Return For Follow-up Did Not Cooperate Protocol Violation Entry Violation Intercurrent Illness Administrative/Other If terminated early, explain briefly: | | | | | | | | | |

ř

8

| INVESTIGATOR'S NAME | PT INITIAL | PT # COMPOL | JND: WR6026 | | | | |
|-------------------------|----------------|--------------------------|-----------------|------------------|--|--|--|
| LIETMAN | D L A F M L | 05 PROTOC | COL:DAMD 17 | -85-C-5133 03 | | | |
| CONCOMITANT MEDICATIONS | | | | | | | |
| DRUG NAME | | TOTAL DAILY DOSE | DATE STARTED | DATE STOPPED | | | |
| heparin solution | 100-1000 unt | \$ \$21 Jul 86 | 245486 | | | | |
| , v | | | | | | | |
| | | | | | | | |
| | | <u> </u> | <u></u> | | | | |
| | | | | | | | |
| | | | | | | | |
| (please | | MMENTS sign all comme | ents) | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |

X.

X

*2*0

); };

288

R

ń

page 1

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | P L | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 |

PROTOCOL

| Study Day | Date ddmmmyy | Procedures |
|--------------|-----------------|------------------------|
| | 22 Jul 86 | Screening laboratory |
| _ | 13 Aug 86 | History, Physical Exam |
| 0 | 16 Aug 86 | Admission |
| 3 | 18 Aug 86 | Drug Administration |
| | J | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 14 pages, for subject # 06.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

。

page 2

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | M - P F M L | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICAL HISTORY

Date of evaluation 13/Aug/86 dd mmm yy

Date of birth

16/Mar/66 dd mmm yy

Age

20 yrs

Sex

M _B__

Race

50

| | No | Yes | Comments |
|-------------------------------|----------|----------|-----------------------------|
| Allergy | V | | |
| Tobacco Use | | 1 | 1/2 PPD |
| Alcohol Use | V | | |
| Recreational Drug Use | | V | MJ Q weekend |
| Medications past 2 weeks | \ | | |
| Experimental Drug Exposure | | ✓ | ceftriaxone, pyridostigmine |
| Blood or plasma donor | V | | |
| Prior Surgery | V | | |
| Eye, ear, nose, throat | ~ | | |
| Endocrine (diabetes, thyroid) | / | | |
| C-V (heart murmur, HBP) | ~ | | |
| Pulmonary (cough, asthma) | V | | |
| Hepatitis, gastro-intestinal | V | | |
| Genito-urinary | V | | |
| Musculoskeletal | V | | |
| Neuropsychiatric | V | | |
| Other | V | | |
| | | | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | M - P | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 |

器

S

Ś

PHYSICAL EXAMINATION Date 131 Aug 1 80 dd minn yy

| Temperature | Pulse | Respir | Blood | Pressure | Height | (cm) | Wt. (kg) |
|-------------|---------|--------|-------|----------|--------|------|----------|
| 36.7c | 070/min | 16/min | 1 1 | 0190 | 1.7. | eO_ | 70.5 |

| | - | | GENERAL EXAMINATION: |
|--------------|---------------|--------------|---|
| (check) | Nor. | Abn. | Provide details of abnormalities |
| Head/Neck | \checkmark | 493 | 1 |
| BENT | | / | minimal asteriolar marrowing Slight wide + doe optic to cups |
| Chest, lungs | / | | Mild rhinchus Sorlado & experation |
| Heart | i / | | |
| Abdomen | / | | |
| Genitalia | not | Rine | |
| Rectal | not | done | |
| Extremities | | | |
| Skin | | \checkmark | Own scar @ upper asm, bound scarle thumb |
| Neurologic | / | | |

CHEST X-RAY
Date 02/feb/86

| NORMAL | X | ABNORMAL | Describe | abnormalities: |
|--------|---|----------|----------|----------------|
| | | | • | |

Examiner

Dr. B.G. Pitty
print name

| INVESTIGATOR'S NAME | PT IN | VITIAL | PT # | COMPOUND:WR6026 |
|---------------------|-------|--------|------|-----------------------------------|
| LIETMAN | M - | - 12 | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICATION RECORD

STUDY: WR6026

| Study Day | Date ddmmmyy | Start drug admin- istration (0-2400) | End drug administration (0-24 | - Route | Bottle I.D. # |
|--------------|-----------------|---|-------------------------------|---------|------------------|
| 3 | 18 Aug 86 | 0800 | N.A. | PO | |

* Below Assay Sensitivity

DOSAGE (total) 60 mg

PLASMA CONCENTRATIONS

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [8503RW] | Methemo- globin * |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|----------|-------------------------|
| | 0 | 18 Aug Sc | 0755 | 0755 | * | |
| | 0.25 | 18 Ary 80 | 0815 | 0815 | * | |
| | 0.50 | 18141286 | 0830 | 0830 | 8.58 | |
| | 0.75 | 18 Prug So | 0845 | 084.5 | 13.7 | |
| | 1.0 | 1812486 | 0900 | 1900 | 24.0 | |
| | 1.25 | 18 My 80 | 0915 | 0915 | 29.4 | |
| | 1.50 | 18 AUG &C | 0930 | 0930 | 31.6 | |
| | 2.0 | 18 Artale | 1000 | 1000 | 49.0 | |
| | 2.5 | 18 Aug So | 1030 | 1030 | 42.9 | |
| | 3.0 | 18PU486 | 1100 | 1100 | CD 3 | |
| | 3.5 | 18AUG86 | 1130 | 1130 | 54.5 | |
| | 4.0 | 18 AUG | 1200 | 1200 | 60.4 | |
| | 5.0 | BAUGE | 1300 | 1300 | 50.5 | |
| | 6.0 | 18 Aug 80 | 1400 | 1400 | 50.2 | |
| | 8.0 | 18 AL 186 | 1600 | 1600 | 38.5 | |
| | 10.0 | 18Aug Sto | 1800 | 1800 | 32.6 | |
| | 12.0 | 1894480 | 1 . | 2000 | 30.2 | |

page 5

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | | |
|---------------------|------------|------|-----------------------------------|--|--|
| LIETMAN | M-P FML | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 | | |

MEDICATION RECORD

STUDY: WR6026

| Study | Date ' | Start drug admin- | End drug admin- | Route | Bottle |
|-------|----------|--------------------|--------------------|-------|--------|
| Day | ddmmmyy | istration (0-2400) | istration (0-2400) | | I.D. # |
| 3 | 18 Aug86 | 0800 | N.A. | PO | |

* Below Assay Sensitivity

DOSAGE (total) 60 mg

PLASMA CONCENTRATIONS

page 2

| page 2 | | | | | | |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|----------|---------------------------------------|
| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo- g lobi n _≯ |
| | 24.0 | 19 Au 86 | 0800 | 0805 | 13.4 | |
| | 36.0 | KI Ariy Slo | 2000 | 2000 | 10.4 | |
| | 48. Ü | 20 AU 86 | l . | 0807) | 7.73 | |
| | 60.0 | 20 Aug 86 | | 20083 | * | |
| | 72.0 | 21 Alg86 | | 0802 2 | ** | |
| | 84.0 | 21 Aug Bo | 1 - | 2000 | * | |
| | 96.0 | 22 Male | 0800 | 0800 | * | i |
| | | J | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | 157 | | | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | M - P F M L | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 |

URINE CONCENTRATIONS

STUDY: WR6026

Ж

7

*

N.

8

Ý.

8

\$3 \$3

STATESTER BASES

| Study | Date | Start drug admin- | End drug admin- | Route | Bottle |
|-------|-----------|--------------------|--------------------|-------|--------|
| Day | ddmmmyy | istration (0-2400) | istration (0-2400) | | I.D. # |
| 3 | 18 Aug 36 | 0810 | N.A. | PO | |

DOSAGE (total) 60mg

| Sample No. | Scheduled Collection Time | Start Co | ollection | End Cal | lection | Total Volume | [WR6026] |
|---------------|------------------------------|------------|-----------|------------|---------|-----------------|----------|
| No. | (hours) | ddmmmyy | 0-2400 | ddmmmyy | 0-2400 | (ml) | mg. |
| U01 | -24 TO -12 | 17 Aug 80 | 0800 | 17 Aug 86 | 2000 | 2000 | |
| nos | -12 TD 0 | 7 Augst | 2000 | 18 Aug 86 | 9758 59 | 1550 | 0 |
| U03 | O TO 12 | 18 Aug 86 | 8350 55 | 18 Arig & | 2000 | 1230 | 0.12177 |
| U04 | 12 TO 24 | 18 Aug 86 | 2000 | 19 Aug & | 0800 | 1345 | 0.11836 |
| U05 | 24 TO 3 6 | 19 Aug Sto | 0800 | 19 Aug80 | 2000 | 1960 | 0.05292 |
| UOE | 36 TO 48 | 19 mg 80 | 2000 | 20 Aug 86 | 0800 | 1480 | 0.04144 |
| U07 | 48 TO 60 | 20 Augsto | 0800 | 20 Aug Sto | 2000 | 950 | 0.0133 |
| HOB | 60 TO 72 | 20 Aug 86 | 2000 | 21 Aug810 | 8336 4 | 1635 | 0.031065 |
| noa | 72 TO 84 | 21 Ary 86 | BEE 39 | 21 Mig86 | 2000 | 1600 | 0 |
| U10 | 84 TO 96 | 21 Aug86 | 2000 | 22 Aug 80 | 0800 | 1570 | O |
| | | U. | · | U | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | , | | | | | | ~ |

<u></u>

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | M - P F M L | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 |

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Screen - | Predrug 1 | Predrug #3 | Post 4 | tdrug 5 | Day |
|----------------------|--------------------|--------------|---------------|-----------|-------------------|------|
| TEST:NORMAL | 22Jµ/86 ddmmmyy | | 18 Aug 86 | | 20 Aug 86 ddmmwyy | Date |
| NA:135-148 MEQ/L | 140 | 142 | 141 | 140 | 140 | |
| K:3.5-5.0 MEQ/L | 4.3 | 4.1 | 4.4 | 4.1 | 4.2 | |
| CL:96-109 MEQ/L | 109 | 107 | 108 | 106 | 105 | |
| CD2:24-30 MEQ/L | 20 | 17 | 18 | 23 | 25 | |
| SUN:12-25 MG/DL | 13 | 8 | 7 | 7 | 13 | |
| CREAT: 0.4-1.5 MG/DL | 1.3 | 1.0 | 1.1 | 1.1 | 1.0 | |
| GLU:70-115 MG/DL | 129 | 60 | 80 | 79 | 75 | |
| T. BILI:0.3-1.2MG/DL | 1.1 | 0.3 | 0.5 | 0.4 | 0.4 | |
| D. BILI:0.1-0.4MG/DL | 0./ | 0.1 | 0.0 | 0.0 | 0.0 | |
| CA:9.0-11.0 MG/DL | 10.6 | 9.2 | 9.3 | 9.5 | 9.7 | |
| PO4:3.0-4.5 MG/DL | 3.8 | 4.1 | ND | 4.7 | 5.2 | |
| URIC A:4.2-8.8MG/DL | 4.9 | 4.5 | 4.2 | NΔ | 4.0 | |
| T. PROT:6.0-8.5G/DL | 7.6 | 6.1 | 6.2 | 6.1 | 6.4 | |
| ALB.:3.2-5.3 G/DL | 4.7 | 4.0 | 4.0 | 4.1 | 4.3 | |
| AST:0-35 IU/L | 27 | 21 | 18 | 22 | 22 | |
| ALT:0-30 IU/L | 18 | 15 | 14 | 19 | 21 | |
| ALK PHOS:0-95 IU/L | 43 | 42 | 43 | 43 | 48 | |
| CHOL:151-268 MG/DL | 149 | 147 | 164 | 154 | 165 | |
| LDH:0-200 IU/L | 142 | 134 | 146 | 123 | 115 | |
| CPK:0-160 U/L(male) | 348 | 163 | 111 | 102 | 116 | |
| TG:20-190 MG/DL | 44 | 45 | 86 | 92 | 255 | |

225

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | M 1 L | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 |

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug 7

2

8

88

W.

8

Day

Date

| TEST: NORMAL | 22Ax36 ddmmmyy | 33Sept 36 | ddmmmyy | ddmmmyy | ddmmmyy | ı |
|----------------------|-------------------|-----------|---------|---------|---------|---|
| NA:135-148 MEQ/L | 139 | ND | | | | |
| K:3.5-5.0 MEQ/L | 4.1 | | | | | |
| CL:96-109 MEQ/L | 103 | | | | | |
| CO2:24-30 MEQ/L | 27 | | | | | |
| SUN:12-25 MG/DL | 13 | | | | | |
| CREAT: 0.4-1.5 MG/DL | 1.0 | | | | | |
| GLU:70-115 MG/DL | 69 | | | | | |
| T. BILI:0.3-1.2MG/DL | 0.4 | | | | | |
| D. BILI:0.1-0.4MG/DL | 0.0 | | | | | |
| CA:9.0-11.0 MG/DL | 9.4 | | | | | |
| PO4:3.0-4.5 MG/DL | Channel Matt. | | | | | |
| URIC A:4.2-8.8MG/DL | 4.0 | | | | | |
| T. PROT:6.0-8.5G/DL | 7.0 | | | | | |
| ALB.:3.2-5.3 G/DL | 4.7 | | | | | |
| AST:0-35 IU/L | 53 | 34 | | | | |
| ALT:0-30 IU/L | 49 | P 33 10 | | | | |
| ALK PHOS:0-95 IU/L | 40, 0 | 必益 | | | | |
| CHOL:151-268 MG/DL | 182 | ND | | | | |
| LDH:0-500 IN/F | 148 | | | | | |
| CPK:0-160 U/L(male) | 140 | | | | | |
| TG:20-190 MG/DL | 218 | | | | | |

page 9

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | |
|---------------------|------------|------|-----------------------------------|--|
| LIETMAN | M = P | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

HEMATOLOGY VALUESLaboratory: JOHNS HOPKINS HOSPITAL

| | • | Screen - | Predrug 1 | Predrug 3 | Postdrug 4 5 | | Day |
|----------|------------|---------------------|--------------------|--------------|-----------------|----------------------|------|
| TEST | NORMAL | 22Jul 86 ddmmmyy | 16Augsb ddmmmyy | 18 Auf 86 | 19Aug86 | 20 Aug 86 ddmmmyy | Date |
| WBC | 4500-11000 | 6900 | 10300 | 7300 | 7000 | 8100 | } |
| RBC | 4.50-5.90 | 5.31 | 4.71 | 5.14 | 5.01 | 5.10 | |
| Hgb | 13.9-16.3 | 15.2 | 13.6 | 15.4 | 14.7 | 15.4 |] |
| PCV | 41.0-53.0 | 45.8 | 40.8 | 45.2 | 4329 | 44.0 | |
| Plt | 150-350 | 268 | 266 | 265 | 286 | 272 | |
| Bands | 2-6% | 1 | 2 | 0 | 0 | 0 | } |
| Polys | 31-76% | 51 | 61 | 58 | 61 | 56 | |
| Eos | 1-4% | 1 | - 1 | 0 | 4 | 2 | |
| Bas | | 0 | 0 | ſ | 0 | 0 | |
| Lymphs | 24-44% | 40 | .31 | 37 | 33 | 32 | } |
| Atyp Lym | | / | 0 | 0 | 0 | / | } |
| Monos | 2-11% | 6 | 5 | 4 | 2 | 9 |] |
| Other | | 0 | 0 | 0 | 0 | 6 | |
| | | | | | | | } |
| Methem: | | 0.5 | | 0.2/03 | 0.3 | 0.7 | |
| G-6-PD | 7.4-9.4 | 7.8 | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | } |
| | | | <u> </u> | | | | 1 |

EXX.

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | |
|---------------------|------------|------|-----------------------------------|--|
| LIETMAN | M - P | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

THE PROPERTY OF THE PROPERTY O

3

8

Y

HEMATOLOGY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug

7

Day

Date $\mathfrak{M}_{\mathcal{A}}$ ddmmmyy TEST NORMAL ddmmmyy ddmmmyy ddmmmyy ddmmmyy 4500-11000 WBC 9(00) RBC 4.50-5.90 5.21 13.9-16.3 Hgb 16.1 PCV 41.0-53.0 45.7 Plt 150~350 PTC Bands 2-6% Polys 31-76% Eos 1-4% Bas 24-44% Lymphs 29 Atyp Lym 2-11% Monos Other 0 methe m 0.7 G-6-PD 7.4-9.4

page 11

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | M - P F M L | 06 | PROTUCOL:DAMD 17-85-C-5133- 03 |

URINALYSIS VALUES
Laboratory: JOHNS HOPKINS HOSPITAL

| • | Scree - | Screen Predrug - 1 | | Postdrug 9 4 5 | |
|----------|------------|-----------------------|-----|-------------------|----------------------|
| | L3AL ddmm | 16 Augs | 6 8 | ddmmmyy | 22 Aug Su ddmmwyy |
| Sp.Gr. | 1.0 | 17 1.023 | | | 1,016 |
| pri · | 7. | i | | | 6.0 |
| Protein | Nei | 7 hea | | | nea |
| Glucose | ne | , , , | |] | nto |
| Ketones | Ned | neg | | | neg |
| Bili. | $ \ln e$ | 9 hea | | | nca |
| Occ.Bld. | Ne | 9 nea | | | neg |
| Cast/lpf | 10 | 6 | | | Ø |
| wBC/hpf | nane | nare | | | Ø |
| RBC/hpf | rar | e 0 | | | Ø |
| Epi./hpf | 0 | D | | | 2-4 |
| Crys/hpf | 0 | 0 | | | Ø |
| Bact/hpf | 0 | 0 | | | 8 |

ELECTROCARDIOGRAM

| Date ddmmmyy | Time 0-2400 | NORMAL check | ABNORMAL check | Describe abnormalities |
|------------------------|----------------|-----------------|-------------------|-----------------------------|
| 13 Aug 86 18 Aug 86 | 0745 | 7 | | |
| 18 Aug 86 | 1147 | | / | relator low atrial shouther |
| 19 Aug 86 | 0830 | / | | |
| 20 Aug 86 | 0825 | V | | |
| 22 Aug 86 | 0810 | V | | |

X

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | M - P F M L | 06 | PROTOCOL:DAMD 17-85-C-5133- 03 |

ADVERSE EXPERIENCES

check if none occurred

 \aleph

838

| 1 | OVERSE EVENT | ONSET (Date/Time) | CESSATION Date/Time | | REL'N TO TEST DRUG | ACTION (Check) |
|---|----------------------------|--------------------------------|---------------------------|-------------|------------------------------|----------------------------|
| 1 | Incicase in Anglepundes | 20 frug Red dd mmm yy (0-2400) | dd mmm yy White (0-2400) | Mild Mod | DEF. PROB. POSS. DEF. NOT | None Treatment Stop test |
| * | | 22 Aug Be | 23 Sep 12. | | DEF | drug |
| 2 | Jeune James Transferance | · · | <u>/030</u> (0-2400) | Mod | PROB. POSS. DEF. NOT | Treatment |
| | | | | Sev | ONKNOMN | Stop test drug |

COMMENTS (Indicate # and event)

page 13

| INVESTIG | ATOR'S NAME | COMPOUND: WR | 5026 | | | | |
|---------------------------------------|---|--|---------|---------------|-----------------------|--|--|
| LIETMAN , | | M - L | D6 | PROTOCOL:DAMD | | | |
| | OUTCOME (To be completed for all subjects) | | | | | | |
| X | Protocol co | ompleted | | | 22 Aug 86 admindry | | |
| | Premature termination of protocol | | | | | | |
| | | Return For operate dolation at Illness | riate d | category) | | | |
| If terminated early, explain briefly: | | | | | | | |
| | | | | | | | |
| | | | | | | | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOU | ND: WREOZE | |
|---|----------------|----------------|--------------------------|-----------------|-----------------|
| LIETMAN | M - P F M L | 06 | PROTOCOL:DAMD 17-85-C-51 | | 1 |
| CONCOMITANT MEDICATIONS | | | | | |
| DRUG NAME | | TO DAILY | TAL DOSE | DATE STARTED | DATE STOPPED |
| hepasin flush | | 100-1000 units | | \$ 17 Aug 86 | 19 Aug 86 |
| | | | | | |
| COMMENTS (please date and sign all comments) | | | | | |
| | | | | | |
| | | | | | |

X

page 1

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | A I D F M L | 07 | PROTOCOL:DAMD 17-85-C-5133- 03 |

PROTOCOL

| Study Day | Date ddmmmyy | Procedures |
|--------------|--------------|------------------------|
| | 19Jul 86 | Screening laboratory |
| | 165 ul 86 | History, Physical Exam |
| 0 | 19Jul 86 | Admission |
| 3 | 21Jul 86 | Drug Administration |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

I certify that:

82

- 1) I have carefully examined all entries on the data collection forms, consisting of 14 pages, for subject # 07.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

10, Dec, 10 dd/mmm/yy

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | A J D F M L | 07 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICAL HISTORY

Date of evaluation 16/Jul/86 dd mmm yy

Examiner

Dr. B.G. Petly

Date of birth

20/Jun/62 dd mmm yy

print name

Age

A

3

Š

¥.

24 yrs

Sex

<u>M</u>_

Race

<u>B</u>

| | No | Yes | Comments |
|------------------------------|----------|----------|--------------------------|
| Allergy | / | | |
| Tobacco Use | | √ | T PPD |
| Alcohol Use | | ✓ | 1/2 pint liquor +6 mck/d |
| Recreational Drug Use | | Y | 1/d (marineral) |
| Medications past 2 weeks | | 1 | multivits |
| Experimental Drug Exposure | | V | Pharmakinetins |
| Blood or plasma donor | ~ | | |
| Prior Surgery | ~ | | |
| Eye, ear, nose, throat | 1 | | |
| Endocrine(diabetes, thyroid) | / | | |
| C-V (heart murmur, HBP) | ١ | | |
| Pulmonary (cough, asthma) | \ | | |
| Hepatitis, gastro-intestinal | / | | |
| Genito-urinary | / | | |
| Musculoskeletal | V | | |
| Neuropsychiatric | / | | |
| Other | / | | |
| | | | |

page 3

| INVESTIGATOR'S NAME | PT INITIAL | PT # | | | |
|---------------------|------------|------|-----------------------------------|--|--|
| LIETMAN | A I D | 07 | PROTOCOL:DAMD 17-85-C-5133- 03 | | |

PHYSICAL EXAMINATION Date 16/Jul/86 dd mmm yy

| ų | Temperature | | | | | 3 | , , |
|---|-------------|---------|--------|----|------|------|---------|
| | 37.0c | _68/min | 14/min | 12 | 2178 | 179. | 5 5 7.5 |

| | | | GENERAL EXAMINATION: |
|--------------|------|------|---|
| (check) | Nor. | Abn. | Provide details of abnormalities |
| Head/Neck | / | | Mouth-mild pyorrhea Ears-Bilat Ceruminous Pose-Septum St 76 |
| EENT | | / | Ears-Bilat ceruminous Pose-septum 51-70 funduscopil-Perinacular pigmentation & Throat-lymphal oc |
| Chest, lungs | V | | |
| Heart | V | | |
| Abdomen | / | | Liver edge at RCM |
| Genitalia | | | Not Done |
| Rectal | | | not done |
| Extremities | | ~ | hyperpigmented birth mart, scratch mark, and scan (& shoulder scar () upper arm, () crown of head, () lat. Thigh & chest, |
| Skin | | ~ | sour Oupper arm (Ucrown of nead, (L) lat. thigh & chest, both hands |
| Neurologic | 1 | | |

Comments: mild Lumbar Scoliosis "Straight back" thoracic spine

CHEST X-RAY
Date 16/Tul/86

| NORMAL | χ | ABNORMAL | Describe abnormalities: |
|--------|---|----------|-------------------------|
| | · | | |

Examiner Both Mittel

Dr.B.G. Retty

33

X

| INVESTIGATOR'S NAME | PT INITIAL | PT # | PROTOCOL:DAMD 17-85-C-5133- | |
|---------------------|------------|------|-----------------------------------|--|
| LIETMAN | A I D | 07 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

MEDICATION RECORD

STUDY: WR6026

| Study | Date | Start drug admin- | End drug admin- | Route | Bottle |
|-------|----------|--------------------|--------------------|-------|--------|
| Day | ddmmmyy | istration (0-2400) | istration (0-2400) | | I.D. # |
| 3 | 21 Jul86 | 0817 | N.A | PO | |

* Below Assay Sensitivity

DOSAGE (total) 60 mg

PLASMA CONCENTRATIONS

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo globin |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|----------|-------------------|
| | 0 | 21 Jul 86 | 0750 | 0750 | * | |
| | 0.25 | 21 Jul 86 | 0832 | 0832 | * | |
| | 0.50 | 21 Jul 86 | 0847 | 0847 | 7.17 | |
| | 0.75 | 21 Jul 86 | 0902 | 0902 | 14.4 | |
| | 1.0 | 21 Jul 86 | 0917 | 0917 | 35.9 | |
| ı | 1.25 | 21 Jul 86 | 0932 | 0932 | 73.2 | |
| | 1.50 | 21 Jul 86 | 0947 | 0947 | 85.7 | |
| | 2.0 | 21 Jul 86 | 1017 | 1017 | 116. | |
| | 2.5 | 21 Jul 86 | 1047 | 1047 | 103. | |
| | 3.0 | 21 Jul 86 | 1117 | 1117 | 94.1 | |
| | 3.5 | 21 Jul 86 | 1147 | 1147 | 94.9 | |
| | 4.0 | 21 TW 86 | 1217 | 1217 | 102. | |
| | 5.0 | 21 Jul 86 | /3 <i>/7</i> | 1317 | 91.4 | |
| | 6.0 | 21 Jul 86 | 1417 | 1417 | 89.7 | |
| | 8.0 | 21 Jul 86 | 1617 | 1647 | 88.4 | |
| | 10.0 | 21 J W 86 | 1817 | 1817 | G4.6 | |
| | 12.0 | 21 Jul 86 | 2017 170 | | (6.0 | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | A 1 D F M L | 07 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICATION RECORD

STUDY: WR6026

| Study Day | Date ddmmmyy | Start drug admin- istration (0-2400) | End drug admin- istration (0-2400) | Route | Bottle I.D. # |
|--------------|--------------|---|---------------------------------------|-------|------------------|
| 3 | 215486 | 0817 | N.A. | PO | |

* Below Assay Sensitivity

PLASMA CONCENTRATIONS

page 2

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo- -globin |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|------------|---------------------|
| | 24.0 | 22Jul 86 | 0817 | 0817 | 31.0 | |
| | 36.0 | 22 Jul 86 | 2017 | 2017 | 17.0 | |
| | 48.0 | 23 Jul 86 | 0817 | 0820 | 11.3 | |
| | 60.0 | 23 Jul 86 | 2017 | 2135 | * | |
| | 72.0 | 24 Jul 86 | 0817 | 0822 | * | |
| | 84.0 | 29 Jul 86 | 2017 | 2022 | a ⊬ | |
| | 96.0 | 25 Jul 86 | 0817 | 0820 | * | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |

171

| INVESTIGATOR'S NAME | PT INITIAL PT # | COMPOUND: WR6026 |
|---------------------|-----------------|-----------------------------------|
| LIETMAN | A 1 D 07 | PROTOCOL:DAMD 17-85-C-5133- 03 |

URINE CONCENTRATIONS

STUDY:WR6026

| | | Start drug admin- istration (0-2400) | | | Bottle I.D. # |
|---|-----------|---|------|----|------------------|
| 3 | 21 Jul 86 | 0817 | N.A. | PO | |

DDSAGE (total) 60 mg

URINE CONCENTRATIONS

| Sample No. | Scheduled Collection Time | Start Co | ollection | End Coli | lection | Total Volume | [WR6026] |
|---------------|------------------------------|-----------|-----------|-----------|---------|-----------------|-----------|
| NO. | (hours) | ddmmmyy | 0-2400 | ddmmmyy | 0-2400 | (ml) | ma |
| U01 | -24 TO -12 | 20 Jul 86 | 0800 | 20 Jul 86 | 2000 | 1410 | <u> </u> |
| nos | -12 TO O | 20 Jul 86 | 2000 | 21 Jul 86 | 0805 | 1500 | 5 |
| U03 | 0 TO 12 | 21Jul86 | 0805 | 21 Jul 86 | 2000 | 960 | 0.56448 |
| U04 | 12 TO 24 | 21Jul86 | 2000 | 22 Jul86 | 0800 | 1300 | 0.424 |
| U05 | 24 TD 36 | 22Jul86 | 0800 | 22 Jul 81 | 2000 | 1605 | 0.081955 |
| U0E | 36 TO 48 | 22 Jul 86 | 2000 | 23 Jul 86 | 0800 | 1505 | 0.161.035 |
| U07 | 48 TO 60 | 23Jul 86 | 0800 | 23 Jul 86 | 2000 | 1095 | 0.0449 |
| UOB | 60 TO 72 | 23Ju186 | 2000 | 24 Jul 86 | 0805 | 1195 | 0.0239 |
| U09 | 72 TO 84 | 24Jul86 | 0805 | 24 Jul86 | 2000 | 1565 | \circ |
| U10 | 84 TO 96 | 24Ju186 | 2000 | 25 Jul86 | 0820 | 1460 | 0 |
| | ; | | | | | | |
| | | | | | | | |
| | | | · | | | | |
| | | | | | | | |
| - · | | | | | | | |
| | | | | | | | |
| | | | | | | | |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | A 1 D F M L | 07 | PROTOCOL:DAMD 17-85-C-5133- 03 |

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Screen - | Predrug 1 | Predrug 2 | Post 4 | tdrug 5 | Day |
|------------------------|--------------------|--------------|----------------------------|-----------|---------------------|------|
| TEST:NORMAL | 14Jul86 ddmmmyy | | 2 <i>17u/86</i> ddmmmyy | | 23 J./86 ddmmmyy | Date |
| NA:135-148 MEQ/L | 145 | 145 | 139 | 140 | 139 | j |
| K:3.5-5.0 MEQ/L | 4.5 | 4.7 | 4.4 | 4.4 | 4.4 | |
| CL:96-109 MEQ/L | 109 | 105 | 110 | 106 | 106 | |
| CD2:24-30 MEQ/L | 24 | 20 | 24 | 19 | 18 | |
| SUN:12-25 MG/DL | 7 | 6 | 9 | 9 | 1/ | |
| CREAT: 0.4-1.5 MG/DL | 1.0 | 1.1 | 0.9 | 1.C | 1.1 | |
| GLU:70-115 MG/DL | 85 | 66 | 86 | 87 | 80 | |
| T.BILI:0.3-1.2MG/DL | 0.3 | 0.2 | C.2 | C·3 | C.3 | |
| D. BILI:0. 1-0. 4MG/DL | 0.1 | 0.1 | c.c | 0.1 | 0.1 | |
| CA:9.0-11.0 MG/DL | | 9.7 | 10.0 | 9.2 | 9.5 | |
| PD4:3.0-4.5 MG/DL | 3.5 | 4.6 | 4.6 | 4.9 | 5. c | |
| URIC A:4.2-8.8MG/DL | <i>5</i> .0 | 7.0 | 4.2 | 4.9 | 4.5 | |
| T. PROT:6.0-8.5G/DL | 6.8 | 7.2 | 6.5 | 6.4 | 7.0 | } |
| ALB.:3.2-5.3 G/DL | 4.6 | 4.7 | 4.2 | 4.4 | 4.4 |] |
| AST:0-35 IU/L | 30 | 24 | 37 | 36 | 57 | |
| ALT:0-30 IU/L | 26 | 19 | 28 | 26 | 36 | } |
| ALK PHOS:0-95 IU/L | 51 | 56 | 48 | 54 | 60 | |
| CHOL:151-268 MG/DL | 141 | 141 | 160 | 148 | 176 | |
| LDH:0-200 IU/L | 126 | 186 | 131 | 190 | 177 | } |
| CPK:0-160 U/L(male) | 136 | 186 | 83 | 73 | 64 | |
| TG:20-190 MG/DL | 64 | ND | 119 | 114 | 107 | |

| INVESTIGATOR'S NAME | PT | INITIE | IL I | PT # | COMPOUND: WR6026 |
|---------------------|----|--------|------|------|-----------------------------------|
| LIETMAN | Ā | I D | | 07 | PROTOCOL:DAMD 17~85-C-5133- 03 |

ACCEPTATION OF THE PROPERTY OF

N

2

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug

Day

08 Aug R. 21 Aug 86 25 Ju/86 Date TEST: NORMAL ddmmmyy ddmmmyy ddmmmyy ddmmmyy ddmmmyy NA:135-148 MEQ/L 12 140 K:3.5-5.0 MEQ/L CL:96-109 MEQ/L 102 CO2:24-30 MEQ/L 28 SUN:12-25 MG/DL 9 4 6 CREAT: 0.4-1.5 MG/DL 1.0 GLU:70-115 MG/DL 93 60 105 T. BILI: 0.3-1.2MG/DL 0.3 C.E 0.5 D. BILI: 0. 1-0. 4MG/DL 0.1 CA:9.0-11.0 MG/DL 9.7 10.1 12.1 P04:3.0-4.5 MG/DL 3.6 3. 3 4.3 URIC A:4.2-8.8MG/DL 64 4.9 54 T. PROT:6.0~8.5G/DL 7.0 6.6 ALB.:3.2-5.3 G/DL 4.7 4.4 4.4 35 AST:0-35 IU/L 3/ 39 56 23 ALT:0-30 IU/L 39 27 63 ALK PHOS:0-95 IU/L 48 NID 10 6.1 CHOL:151-268 MG/DL 191 118 100 LDH:0-200 IU/L 150 NO N.A CPK:0-160 U/L(male) 108 TG:20-190 MG/DL 96

174

page 9

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | A I D | 07 | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| | • | Screen - | Predrug 1 | Predrug 3 | Post 4 | tdrug 5 | Day |
|----------|------------|--------------------|--------------|-------------------|---------------------------|------------|------|
| TEST | NORMAL | 14Ju186 ddmmmyy | 19Jul 86 | 217M86 ddmmmyy | 227 <u>186</u> ddmmmyy | , | Date |
| MBC | 4500-11000 | 4500 | 5600 | 5500 | 6300 | 5700 | |
| RBC | 4.50~5.90 | 4.32 | 4.31 | 4.22 | 4.31 | 4.51 | |
| Ндь | 13.9-16.3 | 15.8 | 15.4 | 15.9 | 15.7 | 16.2 | |
| PCV | 41.0-53.0 | 45.5 | 45.6 | 45.4 | 46.4 | 47.8 | |
| Plt | 150-350 | 215 | 339 | 300 | 293 | 276 | |
| Bands | 2-6% | 3 | 5 | 2 | 6 | 3 | |
| Polys | 31-76% | 57 | 50 | 53 | 51 | 5 t | |
| Eos | 1-4% | O | 0 | 2 | 4 | 0 | |
| Bas | | 0 | O | 0 | 0 | 1 | } |
| Lymphs | 24-44% | 38 | 40 | <i>3</i> 8 | 30 | 32 | |
| Atyp Lym | | C | 6 | C | 0 | 2 | |
| Monos | 2-11% | 2 | 5 | 5 | 9 | 12 | |
| Other | | V | V | Ø | e | e | |
| | | | | | | | |
| Methem: | | 0.4 | | 0.2 | 0.0 | 0.0 | |
| G-6-PD | 7.4-9.4 | 8.2 | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |

page 10

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | A I D F M L | U7 | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug

Day

| | | • | | | | |
|---------------|------------|--------------------|---------|---------|---------|---------|
| TEST | NORMAL | 25Jul86 ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy | ddmmmyy |
| WBC | 4500-11000 | 8000 | | | | |
| RBC | 4.50-5.90 | 4.41 | | | | |
| Hgb | 13.9-16.3 | 16.0 | | | | |
| PCV | 41.0-53.0 | 46.7 | | | | |
| Plt | 150-350 | 248 | | | | |
| Bands | 2-6% | 1 | | | | |
| Polys | 31-76% | 61 | | _ | | |
| Eos | 1-4% | 30 | | | | |
| Bas | | 1 | | | | |
| Lymphs | 24-44% | 30 | | | | |
| Atyp Lym | | 0 | | | | |
| Monos | 2-11% | 7 | | | | |
| Other | | 6 | | | | · |
| | | | | | | |
| methem | | 0.3 | | | | |
| G-6-PD | 7.4-9.4 | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| ", | | | | | | |

| INVESTIGATOR'S NAME | | 1 | f i |
|---------------------|-------|----|------------------------------------|
| LIETMAN | A I D | 07 | PROTUCOL:DAMD 1.7-85-C-5133- 03 |

URINALYSIS VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Screen - | Predrug 1 | 4 | Postdrug 5 | 7 |
|-----------|--------------------|--------------|------|---------------|--------------------|
| | 16Jul86 dammmyy | | | qqwwwyy | ZSTul%6 ddmmmyy |
| Sp.Gr. | 1.016 | 1.008 | | | 1.006 |
| рн | 5.0 | 6.0 | | | 6.0 |
| Protein | Neg | heg | 1 | | heg |
| Glucose | neg | hig | | | reg |
| Ketones | neg | neg | | | his |
| Bili. | neg | neg | | | Nea |
| Jec. Bld. | heg | heg | | | neg |
| Cast/lpf | 0 | NI | | | O |
| wBC/hpf | 0 | | | _ | 0 |
| RBC/hpf | 0 | | | | 0 |
| Epi./hpf | 2-4 | | | | 0-/ |
| Crys/hpf | 0 | | - | | 0 |
| Bact/hpf | many | , | | | 0 |

ELECTROCARDIOGRAM

| Date ddmmmyy | Time 0-2400 | NORMAL check | ABNORMAL check | Describe abnormalities |
|--------------------|----------------|-----------------|-------------------|------------------------|
| 16JW86 21Jul 86 | 0645 | 7 | | |
| 21JW86 | 1202 | v' | | |
| 22Ju186 | | / | | |
| 23 Jul 86 | 6909 | ~ | | |
| 25Jul 86 | | | | |

Screen

33

177

page 12

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | A 1 D | 0-7 | PROTOCOL:DAMD 17-85-C-5133- 03 |

ADVERBE EXPERIENCES

| check | if | none | occurred | | ĺ |
|-------|----|------|----------|---|---|
| | | | | (| ı |

| | OVERSE EVENT | ONSET (Date/Time) | CESSATION Date/Time | | REL'N TO TEST DRUG | ACTION (Check) |
|------------|------------------|------------------------|--------------------------------------|----------|-----------------------|-------------------|
| # / | MIDE, Xb. Find | 21 Jul 76 dd mmm yy | /0 <u>Sap</u> <u>Po</u> dd mmm yy | Mild | DEF. | None |
| | , , | (0-2400) | (0-2400) | Mod | PROB. POSS. | Treatment |
| | | | | Sev | DEF. NOT | Stop test drug |
| * | Politic, contain | 23 Jul 73 dd mmm yy | OT hing of | Wild | DEF. | None |
| ((| j | (0-2400) | (0-2400) | ☐ Mod | PROB. POSS. | Treatment |
| | | | | Sev | DEF. NOT | Stop test drug |

COMMENTS (Indicate # and event)

| | | | | , | | | |
|--|---------------------|------------|------|---------------|----------------------------|--|--|
| INVESTI | GATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6 | 026 | | |
| LIETM | 17-85-C-5133- 03 | | | | | | |
| OUTCOME (To be completed for all subjects) | | | | | | | |
| X | Protocol co | ompleted | | | <u>25Ju 186</u> ddmmmyy | | |
| Premature termination of protocol ddmmmyy | | | | | | | |
| REASON FOR PREMATURE TERMINATION (Check appropriate category) Adverse Experience Died During Study Failure To Return For Follow-up Did Not Cooperate Protocol Violation Entry Violation Intercurrent Illness Administrative/Other If terminated early, explain briefly: | | | | | | | |

N.

3

X

K.

88

8

 $\tilde{\Sigma}$

888

C

80

び分

783

X M

M

8

| CONCOMITANT MEDICATIONS TOTAL DATE STOPPED Appain flush 100-1000 units 21 Jul 86 24 Jul 86 COMMENTS | INVESTIGATOR'S NAME | PT INITIAL | PT # COMPOU | ND: WREOZE | | | | |
|--|-------------------------|------------|----------------|------------|-----------------|--|--|--|
| DRUG NAME DRUG NAME DATE STOPPED TOTAL DATE STOPPED STOPPED 100-1000 units 21 Jul 86 24 Jul 84 COMMENTS | LIETMAN | AID | 07 PROTOC | DL:DAMD 17 | | | | |
| DRUG NAME DAILY DOSE STARTED STOPPED LOD-1000 Units 21 Jul 86 24 Jul 86 COMMENTS | CONCOMITANT MEDICATIONS | | | | | | | |
| COMMENTS | DRUG NAME | | | | DATE STOPPED | | | |
| 1 | hepasin flush | | 100-1000 units | 21 Jul 86 | 24 Jul 86 | | | |
| 1 | | | | | | | | |
| (please date and sign all comments) | | | | | | | | |

2

 \S

355

33

X

5.5

80

X

8

**

7

K

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | C L H | 08 | PROTOCOL:DAMD 17-85-C-5133- 03 |

PROTOCOL

| Study Day | Date ddmmmyy | Procedures |
|--------------|--------------|------------------------|
| 3 | 16 JM 86 | Screening laboratory |
| | 17 Jul 96 | History, Physical Exam |
| ٥ | 19 Jul 86 | Admission |
| 3 | 21 3 4 8 4 | Drug Administration |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 14 pages, for subject # 08.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Delie Miller M. D.

Investigator & signature

16, Dec, 24 dd/mmm/yy

| INVESTIGATOR'S NAME | PT | INI | TIAL | PT # | COMPOUND: WR6026 |
|---------------------|--------|-----|------------|------|-----------------------------------|
| LIETMAN | C F | LM | <u>H</u> _ | 08 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICAL HISTORY

Date of evaluation 17/Jul 186

Examiner Breut Petter.

Dr. B. G. Petty

print name

25/00/51 dd mmm yy

Age

X X

70

34 yrs

Sex

<u>M</u>_

Race

No Comments Yes Allergy Tobacco Use 42 PPD Alcohol Use Recreational Drug Use Medications past 2 weeks Experimental Drug Exposure Blood or plasma donor Prior Surgery V Eye, ear, nose, throat Endocrine (diabetes, thyroid) C-V (heart murmur, HBP) Pulmonary (cough, asthma) Hepatitis, gastro-intestinal GC Rxil & PEN-1 Genito-urinary Musculoskeletal Neuropsychiatric Other

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | | |
|---------------------|------------|------|-----------------------------------|--|--|
| LIETMAN | C L H | 08 | PROTOCOL:DAMD 17-85-C-5133- 03 | | |

8

88

X

88

XX XX

X

X

X

PHYSICAL EXAMINATION Date 17/34/86 dd mmm yy

| Temperature | Pulse | Respir | Blood | Pressure | Height (| Cm) | Wt. (kg) |
|-------------|---------|--------|-------|----------|----------|-----|----------|
| 3 4.7 c | _60/min | 10/min | 12 | 8182 | 178 | .0 | 7 6.0 |

| GENERAL EXAMINATION: | | | | | | |
|----------------------|------|------|---|--|--|--|
| (check) | Nor. | Abn. | Provide details of abnormalities | | | |
| Head/Neck | / | | | | | |
| EENT | ب | | mouth-fair dental hippiene type - It deep + wide aptic crees | | | |
| Chest, lungs | _ | | | | | |
| Heart | ~ | | | | | |
| Abdomen | ~ | | | | | |
| Genitalia | L | | Not done | | | |
| Rectal | | | Not done. | | | |
| Extremities | | | | | | |
| Skin | | / | Scan Circulateral Me | | | |
| Neurologic | / | | 0 | | | |

| NORMAL | χ | ABNORMAL | Describe | abnormalities: | |
|--------|-----|----------|----------|----------------|--|
| | . • | | | | |

Examiner

THE PROPERTY OF THE PROPERTY O

page 4

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | C L H F M L | 08 | PROTOCOL:DAMD 17-85-C-5133- 03 |

MEDICATION RECORD

STUDY: WR6026

| | | Start drug admin- istration (0-2400) | | | Bottle I.D. # |
|---|-----------|---|------|----|------------------|
| 3 | 21 Jul 86 | 0820 | N.A. | PO | |

DOSAGE (total) 60 mg

* Below Assay Sensitivity PLASMA CONCENTRATIO

| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo- globin |
|---------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|--------------|--------------------|
| | 0 | 21Jul86 | 0805 | 0805 | * | |
| | 0.25 | 21 Jul 86 | 0835 | 0835 | * | |
| | 0.50 | 21 Jul 86 | 0850 | 0850 | 763 | |
| | 0.75 | 21 Jul 86 | 0905 | 0905 | 13.2 | |
| | 1.0 | 21 Tul 86 | | 0920 | 14.2 | |
| | 1.25 | 21 Jul86 | 0935 | 0935 | 16.3 | |
| | 1.50 | 21 Jul 86 | 0950 | 0950 | 45.3 | |
| | 2.0 | 21 Jul 86 | | 1020 | C5.3 | |
| | 2.5 | 21 Jul 86 | 1050 | 1050 | GO.5 | |
| | 3.0 | 2548 | 1120 | 1120 | 64.4 | _ |
| | 3.5 | 21 Jul 86 | 1150 | 1150 | 47.3 | |
| | 4.0 | 21 Jul 86 | 1220 | 1220 | GG.9 | |
| - | 5.0 | 21 Jul 86 | 1320 | 1320 | <i>(</i> ي.و | |
| | 6.0 | 21 Jul 86 | 1420 | 1420 | <u>54.0</u> | |
| | 8.0 | 2/JW86 | 1620 | 1620 | 30.8 | |
| | 10.0 | 2/JW 86 | 1820 | 1821 | 276 | |
| | 12.0 | 21 Jul86 | | 42021 | 19.7 | |

page 5

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 | |
|---------------------|----------------|------|-----------------------------------|--|
| LIETMAN | C L H F M L | 08 | PROTOCOL:DAMD 17-85-C-5133- 03 | |

MEDICATION RECORD

STUDY: WR6026

85

88

8

X

| Study Day | Date ddmmmyy | Start drug admin- istration (0-2400) | End drug admin- istration (0-2400) | Route | Bottle I.D. # |
|--------------|--------------|---|---------------------------------------|-------|------------------|
| 3 | 21Jul86 | 0820 | NA | PO | |

* Below Assay Sens. tivity

DOSAGE (total) 60 mg

PLASMA CONCENTRATIONS

| page 2 | | | V = V = V = V | | | |
|-------------------------------------|-----------------------------|-----------------|-------------------------------------|----------------------------------|----------|--------------------------|
| Sample No. | Schedule Time (hours) | Date ddmmmyy | Scheduled Clock Time (0-2400) | Actual Clock Time (0-2400) | [WR6026] | Methemo- globin -% |
| | 24.0 | 225486 | 6820 | 0820 | 8.43 | |
| | 36.0 | 22Jul 86 | 2020 | 2020 | 6.60 | |
| | 48.0 | 63Jul 86 | 0920 | 0827 | * | |
| | 60.0 | 23 Jul 86 | 2020 | 2145 | * | |
| | 72.0 | 295486 | 0 8 20 | 0820 | * | |
| | 84.0 | \$24Jel86 | | 2026 | * | |
| | 96.0 | 25 Jul 86 | 0720 | 0820 | * | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| · · · · · · · · · · · · · · · · · · | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | 185 | | | , |

| INVESTIGATOR'S NAME | PT INITIAL PT # | COMPOUND: WR6026 |
|---------------------|-----------------|-----------------------------------|
| LIETMAN | C L H 08 | PROTOCOL:DAMD 17-85-C-5133- 03 |

URINE CONCENTRATIONS

STUDY:WR6026

X

| | | Start drug admin- istration (0-2400) | | | Bottle I.D. # |
|---|---------|---|------|----|------------------|
| 3 | 2174186 | 0820 | N.A. | PO | |

DOSAGE (total) 60 mg

| | | | | | | , | |
|---------------|---------------------------|------------------|--------------|-------------|----------------|--|-------------|
| Sample No. | Scheduled Collection Time | Start Collection | | End Coli | End Collection | | [WR6026] |
| | (hours) | ddmmmyy | 0-2400 | ddmmmyy | 0-2400 | Volume (ml) | mw. |
| UO1 | -24 TO -12 | 20Jul 86 | 0800 | 20Jul 86 | 2000 | 1285 | 0 |
| nos | -12 TO O | 20JU 86 | 2000 | 21 Jul 86 | 0800 | 1920 | |
| U03 | O TO 12 | 21Jul86 | 0800 | 2NU86 | 2000 | 1135 | 0317673 |
| U04 | 12 TO 24 | 215486 | 2000 | 22Jul 86 | 0800 | 1525 | 0.131506 |
| U05 | 24 TD 36 | 12 Jul 86 | 0800 | 22 Jul 86 | ೩೦ೲ | 2590 | 0.042949 |
| noe | 36 TQ 48 | 22 Jul 86 | 2000 | 23 Jul 86 | 0800 | 1615 | 0.0 14099 |
| U07 | 48 TD 60 | 23Jul86 | 0800 | 23 Jul 86 | 2000 | 5 () | 0.00 4174 |
| uoa | 60 TD 72 | 23 Jul 86 | 2000 | 24 Jul 86 | 2600 | .00 | 0.005197 |
| noa | 72 TO 84 | 24 Jul 86 | <i>0</i> 800 | 24 - 1.186 | 2000 | 1860 | 6 |
| U10 | 84 TO 96 | 24Jul 86 | 2000 | 25Jul 86 | රුපිර | 1970 | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| , | | | | | | | |
| | | | | | | | |
| | · | | | | | | _ |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | CLH | 08 | PROTOCOL:DAMD 17-85-C-5133- 03 |

CHEMISTRY VALUES

est 25% 25% 35% 35%

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Screen - | Predrug 1 | Predrug 2 | Pos ⁴ | tdrug 5 | Day |
|----------------------|-------------|--------------|--------------|------------------|------------|------|
| TEST:NORMAL | 167m/86 | 197486 | | ddmmmyy | J.J.W.P | Date |
| NA:135-148 MEQ/L | 143 | 140 | 136 | 140 | 146 | |
| K:3.5-5.0 MEQ/L | 45 | 4.6 | 4.0 | 4.6 | 4.3 | |
| CL:96-109 MEQ/L | 107 | 105 | .01 | 108 | 105 | |
| CO2:24-30 MEQ/L | 28 | 20 | 26 | 18 | 2~ | |
| SUN:12-25 MG/DL | 9 | 9 | 12 | 14 | 16 | |
| CREAT:0.4-1.5 MG/DL | 0.9 | 0.9 | 0.8 | 0.7 | 0.8 | |
| GLU:70-115 MG/DL | 101 | 64 | 70 | 87 | 77 | |
| T. BILI:0.3-1.2MG/DL | 0.5 | 0.2 | 0.5 | 0.2 | 0.3 | |
| D. BILI:0.1-0.4MG/DL | 0.2 | 0.0 | 0.1 | 0.0 | 0.0 | |
| CA:9.0-11.0 MG/DL | 9.5 | 9.8 | 9.9 | 9.2 | 9.9 | |
| PO4:3.0-4.5 MG/DL | 3.4 | 3.6 | 3,5 | 3.5 | 3.8 | |
| URIC A:4.2-8.8MG/DL | 6.7 | 7.2 | 5.9 | 7.4 | 72 | |
| T. PROT:6.0-8.5G/DL | 4.9 | 6.9 | 7.3 | 6.5 | 7.1 | |
| ALB.:3.2-5.3 G/DL | 4.7 | 4.8 | 4.8 | ND | 4.8 | |
| AST:0-35 IU/L | 21 | 17 | 21 | 16 | 18 | |
| ALT:0-30 IU/L | 18 | ત્રે ઉ | 1/ | /3 | 12 | |
| ALK PHOS:0-95 IU/L | 33 | 34 | 33 | 50 | 3/ | |
| CHOL:151-268 MG/DL | 217 | 238 | 154 | 209 | 275 | |
| LDH:0-200 IU/L | 4018113 | 133 | 189 | 1 55 | 129 | |
| CPK:0-160 U/L(male) | 116 | 98 | 85 | .79 | 78 | |
| TG:20-190 MG/DL | 127 | AIX | 22- | 1417 | 12/1 | |

| INVESTIGATOR'S NAME PT INITIAL | | PT # | COMPOUND: WR6026 |
|--------------------------------|----------------|------|-----------------------------------|
| LIETMAN | C L H F M L | 08 | PROTOCOL:DAMD 17-85-C-5133- 03 |

CHEMISTRY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug 7

| | 7 | | | | | Day |
|----------------------|----------------------|----------------------|----------|----------|---------|-----|
| TEST: NORMAL | 25JW86 ddmmmyy | 15 Aug 86 ddmmmyy | | ddmmmyy | ddmmmyy | Dat |
| NA:135-148 MEQ/L | 140 | ND | | | | |
| K:3.5-5.0 MEQ/L | 4.0 | NI | | | | |
| CL:96-109 MEQ/L | 103 | ND | | | | |
| CO2:24-30 MEQ/L | 21 | V . | | | |] |
| SUN:12-25 MG/DL | 14 | 9 | | | | |
| CREAT:0.4-1.5 MG/DL | 0.9 | 1.0 | | | | |
| GLU:70-115 MG/DL | 75 | 129 | | | | |
| T. BILI:0.3-1.2MG/DL | (Lab Accident) ND | 1.4 | | | | |
| D. BILI:0.1-0.4MG/DL | ND | 0.1 | | | | |
| CA:9.0-11.0 MG/DL | NP | 9.9 | | | | |
| PO4:3.0-4.5 MG/DL | ND | 3.7 | | | | |
| URIC A:4.2-8.8MG/DL | NP | 8.2 | | <u></u> | | |
| T. PROT:6.0-8.5G/DL | ND | 72 | | | | |
| ALB.:3.2-5.3 G/DL | MD | 4.9 | | <u></u> | | |
| AST:0-35 IU/L | M | 31 | | | | |
| ALT:0-30 IU/L | ND | 19 | | | | |
| ALK PHOS:0-95 IU/L | NS | 43 | | | | |
| CHOL:151-268 MG/DL | NĎ | 195 | | | | |
| LDH:0-500 IN/F | 192 | MD | | | | |
| CPK:0-160 U/L(male) | 88 | ND | | | | |
| TG:20-190 MG/DL | 185 | NO | | | | |
| L | | 10 | <u> </u> | <u> </u> | L | j |

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | C L H | 08 | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUESLaboratory: JOHNS HOPKINS HOSPITAL

| | • | Screen - | Predrug 1 | Predrug 3 | Post 4 | tdrug 5 | Day |
|----------|------------|--------------------|-----------------------------|---------------------|--------------------|------------------------------|------|
| TEST | NORMAL | 16Jul86 ddmmmyy | 19 <u>Ja</u> 186 ddmmmyy | 2/54/8/6 ddmmmyy | 2254/86 ddmmmyy | <u>23 لمات 23</u> ddmmmyy | Date |
| MBC | 4500-11000 | 6900 | 8000 | 8000 | 8300 | 4200 | |
| RBC | 4.50-5.90 | 5.16 | 5.14 | 5.27 | 4.79 | 5.07 | |
| Ндь | 13.9-16.3 | 14,6 | 14.4 | 15,2 | 13 6 | 11.4 | |
| PCV | 41.0-53.0 | 45,5 | 4 3,5 | 45.5 | 41,4 | 43 7 | |
| Plt | 150-350 | 262 | 295 | 302 | 290 | 294 | |
| Bands | 2-6% | 0 | 4 | 8 | 2 | 8 | |
| Polys | 31~76% | 989-65 | 47 | 41 | .58 | 49 | |
| Eos | 1-4% | 3 | 1 | 2 | J. | 2 | |
| Bas | | 0 | 0 | 1 | 0 | Ĉ | |
| Lymphs | 24-44% | 32 | 48 | 40 | 34 | 3+ | |
| Atyp Lym | | 6 | 0 | 0 | 0 | 1 | |
| Monos | 2-11% | 0 | 0 | 8 | 4 | 2) | |
| Other | | 0 | 0 | 0 | 0 | 0 | |
| | | | | | | | |
| Methem: | 0.0 - 1.5 | 0.3 | | 0.00.2 | 0.4 | 0.1 | |
| G-6-PD | 7.4-9.4 | 7.3 | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |

page 10

| INVESTIGATOR'S NAME PT INITIAL | | PT # | COMPOUND: WR6026 |
|--------------------------------|----------------|------|-----------------------------------|
| LIETMAN | C L H F M L | Œ | PROTOCOL:DAMD 17-85-C-5133- 03 |

HEMATOLOGY VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

Postdrug

7

Day

25/ul86 Date TEST NORMAL ddmmmyy ddmmmyy ddmmmyy ddmmmyy ddmmmyy WBC 4500-11000 200 RBC 4.50-5.90 5.17 13.9-16.3 Hgb 14.4 PCV 41.0-53.0 44.0 303 150-350 Plt Bands 2-6% 0 Polys 31-76% 48 Eos 1-4% I Bas 0 43 Lymphs 24-44% Atyp Lym 1 7 Monos 2-11% Other 0 mother 0.3 0.0-1.5 G-6-PD 7.4-9.4

| INVESTIGATOR'S NAME | PT INITIAL | # T9 | COMPOUND: WREDZE |
|---------------------|------------|------|-----------------------------------|
| LIETMAN | C L # | 08 | PROTUCOL:DAMD 17-85-C-5133- 03 |

URINALYSIS VALUES

Laboratory: JOHNS HOPKINS HOSPITAL

| • | Screen - | Screen Predrug - 1 | | Postdru 4 5 | |
|-----------|-------------|-----------------------|---------|----------------|--------------------|
| | 17Jul86 | 193 126 ddmmmyy | ggmmmyy | qqwwwyy | 25Ju/86 ddmmmyy |
| 5p.Gr. | 1.003 | 1.012 | | 1 | 1.008 |
| рН | 6.0 | 6.0 | | | 6.0 |
| Protein | Neg | Neg | | 1 | Nea |
| Glucose | Neg | Neg | | | nea |
| Ketones | Ned | Neg | | ! ! ! | Nea |
| Bili. | Ned | Neu | | (} ; | Neg |
| űcc. Ald. | Ned | Nea | | | nea |
| Cast/lpf | 1101 | النتوارا | | | 0 |
| wBC/hpf | 0-1 | | | | O |
| RBC/hpf | | | | | U |
| Epi./hpf | $\bigcup v$ | | | | 0 |
| Crys/hpf | 6 | | | | 0 |
| Bact/hpf | 0 | | | | 0 |

ELECTROCARDIOGRAM

| Date ddmmmyy | Time 0-2400 | NORMAL check | ABNORMAL check | Describe abnormalities |
|--------------------|----------------|-----------------|-------------------|------------------------|
| 17JW86 21J1 136 | 1446 | 1 | | |
| 38 W Cho | 1137 | V | | |
| adJul 86 | 0917 | V | | |
| 48 الملخة | 0910 | / | | |
| 25 Jul 86 | 0844 | V | | · |

screen

| INVESTIGATOR'S NAME | PT INITIAL | PT # | COMPOUND: WR6026 |
|---------------------|----------------|------|-----------------------------------|
| LIETMAN | C L H F M L | 08 | PROTOCOL:DAMD 17-85-C-5133- 03 |

ADVERSE EXPERIENCES

check if none occurred

| ∇ |
|------------------|
| IXI |
| $I \setminus XI$ |
| \mathbf{z} |
| <i>,</i> |

| 1 | DVERSE EVENT | ONSET (Date/Time) | CESSATION Date/Time | | REL'N TO TEST DRUG | ACTION (Check) |
|---|-----------------|-------------------|------------------------|-----------|-----------------------|-------------------|
| # | | | | | | |
| | | dd mmm yy | dd mmm yy | Mild | DEF. | None |
| | | | | | | |
| | | (0-2400) | (0-2400) | Mod | POSS. DEF. NOT | Treatment |
| | | | | | | |
| | | | | Se∨ | UNKNOWN | Stop test drug |
| * | | | | | | |
| | | dd mmm yy | dd mmm yy | Mild | DEF. PROB. | None |
| | | | | | | |
| | | (0-2400) | (0-2400) | Mod | Poss. | Treatment |
| | | | | | DEF. NOT | |
| | | | |] > Se | NUKNOMN | Stop test drug |

COMMENTS (Indicate # and event)

200

3

page 13

| INVESTIGATOR'S NAME PT INITIAL PT # COMPOUND: WR6026 | | | | | | | |
|--|-----------------------------------|----------------|------|---------------|----------------------------|--|--|
| LIETMA | Ν. | C L H F M L | 08 | PROTOCOL:DAMD | 17-85-C-5133- 03 | | |
| OUTCOME (To be completed for all subjects) | | | | | | | |
| K | Protocol co | ompleted | **** | | 25 <u>JU</u> 86 ddmmmyy | | |
| | Premature termination of protocol | | | | | | |
| REASON FOR PREMATURE TERMINATION (Check appropriate category) Adverse Experience Died During Study Failure To Return For Follow-up Did Not Cooperate Protocol Violation Entry Violation Intercurrent Illness Administrative/Other If terminated early, explain briefly: | | | | | | | |

| INVESTIGATOR'S NAME | PT INITIAL | - PT # | COMPOUND: WREOZE | | | | |
|---|---------------------|---------|------------------|-----------------|--------------------|--|--|
| LIETMAN | C L H | 08 | PROTOC | OL:DAMD 1 | .7-85-C-5133 03 | | |
| CONCOMITANT MEDICATIONS | | | | | | | |
| DRUG NAME | TOTAL DAILY DOSE | | DATE STARTED | DATE STOPPED | | | |
| heparin flush | | 100 100 | 00 unito | 31Jui 6 | 5Jul 86 | | |
| | | | | | | | |
| | | | | | | | |
| COMMENTS (please date and sign all comments) | | | | | | | |
| | | | | | | | |
| | | · | | | | | |

APPENDIX F Distribution List

1 Copy: Brian Schuster, M.D.

COL, MC

c/o Director

Walter Reed Army Institute of Research

ATTN: SGRD-UWI-F

Washington, D.C. 20307-5100

4 Copies: Commander

Walter Reed Army Institute of Research

ATTN: SGRD-UWI-D

Washington, D.C. 20307-5100

1 Copy: Commander

US Army Medical Research and Development Command

ATTN: SGRD-RMI-S Fort Detrick

Frederick, Maryland 21701-5012

1 Copy: Dean, School of Medicine

Uniformed Services University of the Health Sciences

4301 Jones Bridge Road Bethesda, Maryland 20014

1 Copy: Commandant

Academy of Health Sciences, U.S. Army

ATTN: AHS-CMD

Fort Sam Houston, Texas 78234

2 Copies: Administrator

Defense Technical Information Center

ATTN: DTIC-DDAC Cameron Station

Alexandria, Virginia 22304-6145